

Article

Robust Self-Testing of Four-Qubit Symmetric States

Daipengwei Bao , Xiaoqing Tan *, Qingshan Xu , Haozhen Wang and Rui Huang 

School of Information Science and Technology, Jinan University, Guangzhou 510632, China; baodaipengwei@stu2020.jnu.edu.cn (D.B.); xuqingshan1008@stu2018.jnu.edu.cn (Q.X.); wanghz0125@163.com (H.W.); huang6666@foxmail.com (R.H.)

* Correspondence: ttanxq@jnu.edu.cn

Abstract: Quantum verification has been highlighted as a significant challenge on the road to scalable technology, especially with the rapid development of quantum computing. To verify quantum states, self-testing is proposed as a device-independent concept, which is based only on the observed statistics. Previous studies focused on bipartite states and some multipartite states, including all symmetric states, but only in the case of three qubits. In this paper, we first give a criterion for the self-testing of a four-qubit symmetric state with a special structure and the robustness analysis based on vector norm inequalities. Then we generalize the idea to a family of parameterized four-qubit symmetric states through projections onto two subsystems.

Keywords: Bell inequality; self-testing; symmetric states; device independent

1. Introduction

In recent years, quantum technology has developed rapidly and is expected to gain new real-world applications in communication, simulation, sensing, and computing [1–4]. Quantum devices promise to effectively solve some problems that are difficult to deal with in the classical field [5,6]. However, it also brings a thorny problem. How do we verify the solutions? The task of ensuring the correct operations of quantum devices in terms of accuracy of output is known as quantum verification [7], which is attracting more attention.

A common quantum state verification technology was quantum state tomography (QST) [8] in the past. It has been implemented in systems with few components, but unfortunately, it becomes unfeasible for larger systems because the complexity grows exponentially with the system size. To solve this problem, another alternative technique called self-testing [9] was proposed. These two techniques could be used to verify the quantum systems.

Self-testing is a device-independent approach to verifying that the previously unknown quantum system state and uncharacterized measurement operators are to some degree close to the target state and measurements (up to local isometries) based only on observed statistics, without assuming the dimension of the quantum system. The device-independent (DI) approach [10] is important in practical quantum communications. One of the main applications of self-testing is quantum key distribution (QKD) [11,12], which is of great interest because of its high security. For the users, the quantum key distribution system is purchased from the device providers. However, if a device provider deliberately creates a “dishonest” quantum device, which does not perform key distribution according to the correct protocol, then the key distribution performed with such a device will be insecure. Therefore, it is imperative to test the trustworthiness of quantum cryptographic devices. Fortunately, based on the idea of self-testing quantum systems, it is possible to design device-independent quantum cryptography protocols. For example, in the device-independent QKD protocols, even if the device provider is not trusted, the user can still ensure that the keys generated by the device are secure. The essence is that the user self-tests the quantum device and uses its output as the key under the condition that the test is passed, and the key must be trusted in this case. In addition to quantum key distribution,



Citation: Bao, D.; Tan, X.; Xu, Q.; Wang, H.; Huang, R. Robust Self-Testing of Four-Qubit Symmetric States. *Entropy* **2022**, *24*, 1003. <https://doi.org/10.3390/e24071003>

Academic Editors: Giuliano Benenti, Shunlong Luo, Ming Li and Shao-Ming Fei

Received: 16 May 2022

Accepted: 18 July 2022

Published: 20 July 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

various protocols, such as random number generation [13], and entanglement witness [14], have been designed in a device-independent framework so far.

Let us consider a scenario where N distant observers share an unknown N -partite quantum state $|\Psi\rangle$. Each party can perform uncharacterized measurements $\{M_{a_i}^{x_i}\}$ on the state with their quantum devices, where i marks different parties, x_i marks different measurement settings for party i , and a_i marks the corresponding measurement outcomes. In a device-independent scenario, the process of measuring an unknown quantum state can be viewed as a black box for the N observers: they can only query their devices with possible measurement settings x_i , and to any query, the black box produces a corresponding outcome. As we do not assume the dimension of the quantum system, the dimension of the Hilbert space is not fixed. Without loss of generality, we assume that the unknown state is pure. There is no loss of generality because an extra system can be added to some of the parties, if necessary, to purify the state, and the purification of the state can be included in the black boxes. Similarly, we can further assume that the measurement operators are projective without loss of generality, as an auxiliary system in some known state can be added to the measured system to replace a general POVM on this system by a projective measurement on the extended system [9]. According to the postulates of quantum mechanics [15], the data they observe are given by

$$p(a_1, a_2, \dots, a_N | x_1, x_2, \dots, x_N) = \langle \Psi | M_{a_1}^{x_1} \otimes M_{a_2}^{x_2} \otimes \dots \otimes M_{a_N}^{x_N} | \Psi \rangle, \quad (1)$$

which is referred to as a *correlation* [16] based on the quantum nonlocality [17] of entangled states [18]. As the possibility to self-test quantum states and measurements usually relies on quantum nonlocality, only the entangled states can be device-independently verified by self-testing techniques. The self-testing problem consists of deciding if the knowledge of the *correlation* allows us to deduce the structure of the unknown quantum system.

Symmetric states [19] have been found useful in many quantum information tasks, such as measurement-based quantum computation (MBQC) [20], as they are not too entangled to be computationally universal. Due to the important role of symmetry in the field of quantum entanglement, it is important to explore the properties of symmetric states.

This paper is organized as follows. The basic definitions and preliminaries are given in Section 2. In Section 3, we prove analytically that a particular symmetric four-qubit state can be self-tested and give bounds that are robust to inevitable experimental errors. In addition, we show the self-testing of a family of parameterized four-qubit symmetric states, which are superpositions of four-qubit Dicke states through projections onto two subsystems in Section 4, and we give the conclusions in Section 5.

2. Basic Definitions and Preliminaries

In this section, we present the definitions of self-testing [21] and give the known results as several lemmas, which may be used as building blocks for our work.

Definition 1 (Self-testing). *A known correlation allows for self-testing the state $|\Psi'\rangle$ and measurements $\{M_{a_i}^{x_i}\}$; if any state and measurements $|\Psi\rangle$ and $\{M_{a_i}^{x_i}\}$ reproduce the correlation, there exists a local isometry Φ such that*

$$\begin{aligned} \Phi(|\Psi\rangle) &= |\text{junk}\rangle \otimes |\Psi'\rangle, \\ \Phi(M_{a_1}^{x_1} \otimes M_{a_2}^{x_2} \otimes \dots \otimes M_{a_N}^{x_N} |\Psi\rangle) &= |\text{junk}\rangle \otimes (M_{a_1}^{x_1} \otimes M_{a_2}^{x_2} \otimes \dots \otimes M_{a_N}^{x_N} |\Psi'\rangle), \end{aligned} \quad (2)$$

where the state $|\text{junk}\rangle$ is an auxiliary state which will be traced out and thus not taken into consideration.

The currently known self-testing protocols are mainly tailored for bipartite states [22–26]. We first review two-qubit self-testing. As given in [23,24], all pure two-qubit

entangled states can be self-tested by observing the maximum violation of the tilted CHSH inequality [27]

$$\mathcal{B}(\alpha, A_0, A_1, B_0, B_1) \equiv \alpha A_0 + A_0(B_0 + B_1) + A_1(B_0 - B_1) \leq 2 + \alpha, \tag{3}$$

where $0 \leq \alpha < 2$ and A_i and B_i are observables with outcomes ± 1 . The maximal violation is given by $b(\alpha) \triangleq \max_{\phi} \langle \phi | \mathcal{B}(\alpha, A_0, A_1, B_0, B_1) | \phi \rangle = \sqrt{8 + 2\alpha^2}$.

Lemma 1. Any pure two-qubit states in their Schmidt form $|\Psi_{\theta}\rangle = \cos \theta |00\rangle + \sin \theta |11\rangle$ can be self-tested by achieving the maximal quantum violation of the tilted CHSH inequality Equation (3). The corresponding measurements A_i and B_i for two distant parties, Alice and Bob, are set as

$$\begin{aligned} A_1 &= \sigma_z, B_1 = \cos \mu \sigma_z + \sin \mu \sigma_x, \\ A_2 &= \sigma_x, B_2 = \cos \mu \sigma_z - \sin \mu \sigma_x. \end{aligned} \tag{4}$$

Here, $\sin 2\theta = \sqrt{\frac{4-\alpha^2}{4+\alpha^2}}$ and $\mu = \arctan \sin 2\theta$.

Especially for the maximally entangled two-qubit states in the form $\frac{|00\rangle+|11\rangle}{\sqrt{2}}$, there exist another two criteria [25].

Lemma 2 (Mayers–Yao criterion). Consider five unknown dichotomic measurements $\{X_A, Z_A; X_B, Z_B, D_B\}$. If the following statistics are observed

$$\begin{aligned} \langle \Psi | Z_A Z_B | \Psi \rangle &= \langle \Psi | X_A X_B | \Psi \rangle = 1, \\ \langle \Psi | X_A Z_B | \Psi \rangle &= \langle \Psi | Z_A X_B | \Psi \rangle = 0, \\ \langle \Psi | Z_A D_B | \Psi \rangle &= \langle \Psi | X_A D_B | \Psi \rangle = \frac{1}{\sqrt{2}}, \end{aligned} \tag{5}$$

then up to a local isometry, the state $|\Psi\rangle$ is self-tested into the maximally entangled two-qubit state $\frac{|00\rangle+|11\rangle}{\sqrt{2}}$, and the measurements are the suitable complementary Pauli operators.

Lemma 3 (XOR game). Consider four unknown operators $\{A_0, A_1, B_0, B_1\}$ with binary outcomes ± 1 and let $E_{xy} \equiv \langle \Psi | A_x B_y | \Psi \rangle = \cos \alpha_{xy}$. The state $|\Psi\rangle$ can be self-tested into the maximally entangled two-qubit state $\frac{|00\rangle+|11\rangle}{\sqrt{2}}$ by winning the binary nonlocal XOR game defined by the figure of merit $\sum_{(x,y) \in (0,1)^2} f_{xy} E_{xy}$ if it satisfies $\alpha_{00} + \alpha_{10} = \alpha_{01} - \alpha_{11}$. The coefficients f_{xy} are constructed by

$$\begin{pmatrix} f_{00} \\ f_{01} \\ f_{10} \\ f_{11} \end{pmatrix} = \begin{pmatrix} \sin^{-1} \alpha_{00} \\ -\sin^{-1} (\alpha_{00} + \alpha_{10} + \alpha_{11}) \\ \sin^{-1} \alpha_{10} \\ \sin^{-1} \alpha_{11} \end{pmatrix}. \tag{6}$$

However, the self-testing of multipartite scenarios has not been fully explored. In this paper, we work on the four-qubit symmetric entangled states.

Definition 2 (Symmetric states). Symmetric quantum states preserve invariance under any permutation of their subsystems. We say that an n -partite state $|\Psi\rangle$ is symmetric if $P|\Psi\rangle = |\Psi\rangle$ for all $P \in S_n$, where S_n is the symmetric group of n elements. The n -qubit Dicke states $|S_{n,k}\rangle$ are typical examples of symmetric state, which are the equally weighted sums of all permutations of computational basis states with $n - k$ qubits being $|0\rangle$ and k being $|1\rangle$:

$$|S_{n,k}\rangle = \binom{n}{k}^{-1/2} \sum_{\text{Permutation}} \underbrace{|0\rangle|0\rangle \dots |0\rangle}_{n-k} \underbrace{|1\rangle|1\rangle \dots |1\rangle}_k. \tag{7}$$

Let $|\Psi\rangle$ be a state vector in an N -fold tensor product space $S_1 \otimes \dots \otimes S_N$, where $\dim S_1 = \dots = \dim S_N = d \geq 2$ and $N \geq 3$. As the generalization of the Schmidt decomposition given in [28], if $d = 2$, any multipartite states can be written in the expansion as

$$|\Psi\rangle = \sum_{i_1, i_2, \dots, i_N \in \{0,1\}} t^{i_1 i_2 \dots i_N} |i_1\rangle |i_2\rangle |i_3\rangle \dots |i_N\rangle, \tag{8}$$

where some coefficients satisfy

$$t^{011\dots 11} = t^{101\dots 11} = \dots = t^{111\dots 10} = 0, \tag{9}$$

and the rest $2^N - N$ orthogonal product states

$$\{\underbrace{|000\dots 00\rangle}_N, \underbrace{|000\dots 01\rangle}_{N-1}, \dots, \underbrace{|10\dots 00\rangle}_{N-1}, \dots, \underbrace{|001\dots 11\rangle}_{N-2}, \dots, \underbrace{|11\dots 100\rangle}_{N-2}, \underbrace{|111\dots 11\rangle}_N\} \tag{10}$$

can be seen as a set of local bases. To characterize the symmetric multi-qubit states, we only need to make the rest coefficients have properties

$$\begin{aligned} t^{000\dots 01} &= t^{00\dots 010} = \dots = t^{100\dots 00}, \\ &\vdots \\ t^{001\dots 11} &= t^{0101\dots 1} = \dots = t^{11\dots 100}. \end{aligned} \tag{11}$$

3. Self-Testing of a Four-Qubit Symmetric State

In this section, we focus on a four-qubit symmetric state with a special structure by using the known results. In the case of $N = 4$, as given in Equation (10), the set of local bases is

$$\{|0000\rangle, |0001\rangle, |0010\rangle, |0100\rangle, |1000\rangle, |0011\rangle, |0101\rangle, |0110\rangle, |1001\rangle, |1010\rangle, |1100\rangle, |1111\rangle\} \tag{12}$$

3.1. Self-Testing of a Specific Four-Qubit Symmetric State

The specific four-qubit symmetric state we consider is

$$|\Psi'_1\rangle = \frac{1}{2\sqrt{2}}(|0000\rangle + |0011\rangle + |0101\rangle + |0110\rangle + |1001\rangle + |1010\rangle + |1100\rangle + |1111\rangle)_{ABCD}, \tag{13}$$

which is shared by four distant observers, Alice, Bob, Charlie and David.

Rewrite the state as

$$\begin{aligned} |\Psi'_1\rangle &= \frac{1}{2\sqrt{2}}[\sqrt{2}|00\rangle_{AB} \otimes \frac{1}{\sqrt{2}}(|00\rangle + |11\rangle)_{CD} + \sqrt{2}|01\rangle_{AB} \otimes \frac{1}{\sqrt{2}}(|01\rangle + |10\rangle)_{CD} \\ &\quad + \sqrt{2}|10\rangle_{AB} \otimes \frac{1}{\sqrt{2}}(|01\rangle + |10\rangle)_{CD} + \sqrt{2}|11\rangle_{AB} \otimes \frac{1}{\sqrt{2}}(|00\rangle + |11\rangle)_{CD}]. \end{aligned} \tag{14}$$

The concept of partial measurements [29] is involved in our scheme, which appears very often in reality. A similar approach for quantum nonlocality characterization is given in [30], where quantum incompatibility is used to characterize nonlocality. According to the partial measurement postulate given in [29], if any two parties, without loss of generality, e.g., Alice and Bob, each measure in the σ_z basis, the remaining two parties share a maximally entangled two-qubit state $\frac{|00\rangle + |11\rangle}{\sqrt{2}}$ conditioned on the outcome “00” and “11”, respectively, which can be self-tested combining Lemma 2.

We construct the local isometry Φ as Figure 1. Here, H is the usual Hadamard gate. Obviously, if $Z_i = \sigma_z, X_i = \sigma_x$, we can extract the essential information on the unknown state into auxiliary systems. Inspired by this, Z_i and X_i should act analogously to the Pauli operators on $|\Psi'_1\rangle$ to guarantee the feasibility of the protocol. However, in order to make

the protocol device-independent, we cannot directly consider Z_i and X_i of each party as Pauli operators, but should construct them with the measurements $\{M_{a_i}^{X_i}\}$ properly. We sum the result up as below.




Figure 1. Swap circuit of the isometry Φ to self-test the target state $|\Psi'_1\rangle$.

Result 1. Consider four spatially separated parties, Alice, Bob, Charlie and David, each performing three measurements $\{X_s, Z_s, M_s\} (s \in \{A, B, C, D\})$ with binary outcomes on an unknown shared quantum state $|\Psi_1\rangle$. The target symmetric state $|\Psi'_1\rangle$ is self-tested if the statistics are observed as the following:

$$\begin{aligned} \langle P_A^0 P_B^0 P_C^0 P_D^0 \rangle &= \langle P_A^0 P_B^0 P_C^1 P_D^1 \rangle = \langle P_A^0 P_B^1 P_C^0 P_D^1 \rangle = \langle P_A^0 P_B^1 P_C^1 P_D^0 \rangle \\ &= \langle P_A^1 P_B^0 P_C^0 P_D^1 \rangle = \langle P_A^1 P_B^0 P_C^1 P_D^0 \rangle = \langle P_A^1 P_B^1 P_C^0 P_D^0 \rangle = \langle P_A^1 P_B^1 P_C^1 P_D^1 \rangle = \frac{1}{8}, \end{aligned} \quad (15)$$

$$\begin{cases} \langle P_i^0 P_j^0 X_k X_l \rangle = \langle P_i^0 P_j^0 Z_k Z_l \rangle = \frac{1}{4} \\ \langle P_i^0 P_j^0 X_k M_l \rangle = \langle P_i^0 P_j^0 Z_k M_l \rangle = \frac{1}{4\sqrt{2}} \\ \langle P_i^0 P_j^0 X_k Z_l \rangle = 0 \end{cases}, \quad \begin{cases} \langle P_A^1 P_B^1 X_C X_D \rangle = \langle P_A^1 P_B^1 Z_C Z_D \rangle = \frac{1}{4} \\ \langle P_A^1 P_B^1 X_C M_D \rangle = \langle P_A^1 P_B^1 Z_C M_D \rangle = \frac{1}{4\sqrt{2}} \\ \langle P_i^1 P_j^1 X_k Z_l \rangle = 0 \end{cases}, \quad (16)$$

where $(i, j, k, l) = \{(A, B, C, D), (A, C, B, D), (A, D, B, C), (B, C, A, D), (B, D, A, C), (C, D, A, B)\}$ and $P_s^0 \triangleq P_{Z_s=+1} = \frac{1+Z_s}{2}, P_s^1 \triangleq P_{Z_s=-1} = \frac{1-Z_s}{2}$, where $s \in \{A, B, C, D\}$ are projectors for the Z_s measurement.

Proof. To begin with, the output after the isometry given in Figure 1 is

$$\begin{aligned} |\tilde{\Psi}_1\rangle &= \Phi(|\Psi_1\rangle |0000\rangle_{A'B'C'D'}) \\ &= \sum_{a,b,c,d \in \{0,1\}} X_A^a X_B^b X_C^c X_D^d P_A^a P_B^b P_C^c P_D^d |\Psi_1\rangle |abcd\rangle. \end{aligned} \quad (17)$$

Observation Equation (15) implies that

$$\begin{aligned} &\langle P_A^0 P_B^0 P_C^0 P_D^0 \rangle + \langle P_A^0 P_B^0 P_C^1 P_D^1 \rangle + \langle P_A^0 P_B^1 P_C^0 P_D^1 \rangle + \langle P_A^0 P_B^1 P_C^1 P_D^0 \rangle \\ &+ \langle P_A^1 P_B^0 P_C^0 P_D^1 \rangle + \langle P_A^1 P_B^0 P_C^1 P_D^0 \rangle + \langle P_A^1 P_B^1 P_C^0 P_D^0 \rangle + \langle P_A^1 P_B^1 P_C^1 P_D^1 \rangle = 1, \end{aligned} \quad (18)$$

and thus $P_A^a P_B^b P_C^c P_D^d |\Psi_1\rangle = 0$ for other eight projectors. Based on the fact that $\langle \psi | \phi \rangle = 1$ implies $|\psi\rangle = |\phi\rangle$, observation of Equation (16) implies

$$\begin{cases} P_i^0 P_j^0 X_k |\Psi_1\rangle = P_i^0 P_j^0 X_l |\Psi_1\rangle, P_i^0 P_j^0 Z_k |\Psi_1\rangle = P_i^0 P_j^0 Z_l |\Psi_1\rangle \\ P_i^0 P_j^0 X_k |\Psi_1\rangle \perp P_i^0 P_j^0 Z_k |\Psi_1\rangle, P_i^0 P_j^0 X_l |\Psi_1\rangle \perp P_i^0 P_j^0 Z_l |\Psi_1\rangle \\ P_i^0 P_j^0 M_l |\Psi_1\rangle = \frac{P_i^0 P_j^0 X_l |\Psi_1\rangle + P_i^0 P_j^0 Z_l |\Psi_1\rangle}{\sqrt{2}} = \frac{P_i^0 P_j^0 X_k |\Psi_1\rangle + P_i^0 P_j^0 Z_k |\Psi_1\rangle}{\sqrt{2}} \end{cases}, \quad (19)$$

and

$$\begin{cases} P_A^1 P_B^1 X_C |\Psi_1\rangle = P_A^1 P_B^1 X_D |\Psi_1\rangle, P_A^1 P_B^1 Z_C |\Psi_1\rangle = P_A^1 P_B^1 Z_D |\Psi_1\rangle \\ P_A^1 P_B^1 X_C |\Psi_1\rangle \perp P_A^1 P_B^1 Z_C |\Psi_1\rangle, P_A^1 P_B^1 X_D |\Psi_1\rangle \perp P_A^1 P_B^1 Z_D |\Psi_1\rangle \\ P_A^1 P_B^1 M_D |\Psi_1\rangle = \frac{P_A^1 P_B^1 X_D |\Psi_1\rangle + P_A^1 P_B^1 Z_D |\Psi_1\rangle}{\sqrt{2}} = \frac{P_A^1 P_B^1 X_C |\Psi_1\rangle + P_A^1 P_B^1 Z_C |\Psi_1\rangle}{\sqrt{2}} \end{cases}. \quad (20)$$

Obviously, we have $(P_i^0 P_j^0 M_l)^2 |\Psi_1\rangle = P_i^0 P_j^0 M_l^2 |\Psi_1\rangle$. Since $X^2 = Z^2 = M^2 = I$, we have $P_i^0 P_j^0 |\Psi_1\rangle = \frac{P_i^0 P_j^0 (X_k + Z_k)^2 |\Psi_1\rangle}{2} = \frac{P_i^0 P_j^0 (X_l + Z_l)^2 |\Psi_1\rangle}{2}$. Hence, we obtain the following anti-commutation relation

$$\begin{aligned} P_i^0 P_j^0 X_k Z_k |\Psi_1\rangle &= -P_i^0 P_j^0 Z_k X_k |\Psi_1\rangle \\ P_i^0 P_j^0 X_l Z_l |\Psi_1\rangle &= -P_i^0 P_j^0 Z_l X_l |\Psi_1\rangle \end{aligned} \quad (21)$$

for all $(i, j, k, l) = \{(A, B, C, D), (A, C, B, D), (A, D, B, C), (B, C, A, D), (B, D, A, C), (C, D, A, B)\}$, and similarly,

$$\begin{aligned} P_A^1 P_B^1 X_C Z_C |\Psi_1\rangle &= -P_A^1 P_B^1 Z_C X_C |\Psi_1\rangle \\ P_A^1 P_B^1 X_D Z_D |\Psi_1\rangle &= -P_A^1 P_B^1 Z_D X_D |\Psi_1\rangle. \end{aligned} \quad (22)$$

All these properties of the operators will help to reduce the output Equation (17). By using Equation (21), $X_C X_D P_A^0 P_B^0 P_C^1 P_D^1 |\Psi_1\rangle$ is equal to $P_A^0 P_B^0 P_C^0 X_D P_D^0 X_D |\Psi_1\rangle$. As $P_A^0 P_B^0 X_C |\Psi_1\rangle = P_A^0 P_B^0 X_D |\Psi_1\rangle$ shown in Equation (19), this term becomes $P_A^0 P_B^0 P_C^0 P_D^0 |\Psi_1\rangle$. We can simplify the other five terms similarly. For the last term, we can obtain $P_A^1 P_B^1 P_C^0 P_D^0 |\Psi_1\rangle$ using Equations (20) and (22), which can also be simplified to $P_A^0 P_B^0 P_C^0 P_D^0 |\Psi_1\rangle$. As a reminder, there are eight terms equal to zero. Hence, the output Equation (17) is reduced to

$$|\Psi_1^*\rangle = P_A^0 P_B^0 P_C^0 P_D^0 |\Psi_1\rangle (|0000\rangle + |0011\rangle + |0101\rangle + |0110\rangle + |1001\rangle + |1010\rangle + |1100\rangle + |1111\rangle) \quad (23)$$

and can be normalized into the form of $|junk\rangle \otimes |\Psi_1'\rangle$, here $|junk\rangle = 2\sqrt{2} P_A^0 P_B^0 P_C^0 P_D^0 |\Psi_1\rangle$. \square

3.2. Robustness Analysis Based on the L₂ Norm

In this section, we give the analysis of robustness based on the vector norm inequality. Result 1 relies on the observation of Equations (15) and (16) exactly; however, which may be impossible in actual experiments due to the inevitable deviation from the ideal case. Suppose each observation in Equations (15) and (16) admits a deviation at most ϵ around the ideal value. We say that the self-testing of $|\Psi_1'\rangle$ is robust [31] if the isometry still extracts a state close to it and satisfies

$$\| |\tilde{\Psi}_1\rangle - |junk\rangle \otimes |\Psi_1'\rangle \| \leq f(\epsilon), \quad (24)$$

where $f(\epsilon) \rightarrow 0$ when $\epsilon \rightarrow 0$.

We show that

$$\| |\Psi_1\rangle - |junk\rangle \otimes |\Psi'_1\rangle \| \leq f(\epsilon) = 265.98\epsilon + 348.45\epsilon^{\frac{3}{4}} + 94.87\epsilon^{\frac{1}{2}} + 60.70\epsilon^{\frac{1}{4}} \quad (25)$$

in Appendix A, which proves the robustness of Result 1.

4. Self-Testing of a Family of Parameterized Four-Qubit Symmetric States

In this part, we consider a more general state

$$|\Psi'_2\rangle = \frac{1}{\sqrt{8+4t^2}} [|0000\rangle + t(|0001\rangle + |0010\rangle + |0100\rangle + |1000\rangle) + |0011\rangle + |0101\rangle + |0110\rangle + |1001\rangle + |1010\rangle + |1100\rangle + |1111\rangle]_{ABCD} \quad (26)$$

where $t > 0$ and $t \neq 1$. The parameterized state is a superposition of W state, GHZ state and $|S_{4,2}\rangle$ state, where the ratio of the coefficient of GHZ state and $|S_{4,2}\rangle$ state is a constant value, which is equal to $\frac{1}{\sqrt{3}}$. Rewrite the states as

$$\begin{aligned} |\Psi'_2\rangle &= \frac{1}{\sqrt{8+4t^2}} [\sqrt{2+2t^2} |00\rangle_{AB} \otimes \frac{1}{\sqrt{2+2t^2}} (|00\rangle + t|01\rangle + t|10\rangle + |11\rangle)_{CD} \\ &\quad + \sqrt{2+t^2} |01\rangle_{AB} \otimes \frac{1}{\sqrt{2+t^2}} (t|00\rangle + |01\rangle + |10\rangle)_{CD} \\ &\quad + \sqrt{2+t^2} |10\rangle_{AB} \otimes \frac{1}{\sqrt{2+t^2}} (t|00\rangle + |01\rangle + |10\rangle)_{CD} \\ &\quad + \sqrt{2} |11\rangle_{AB} \otimes \frac{1}{\sqrt{2}} (|00\rangle + |11\rangle)_{CD}]. \end{aligned} \quad (27)$$

Denote

$$\begin{aligned} |\psi_1\rangle &= \frac{1}{\sqrt{2+2t^2}} (|00\rangle + t|01\rangle + t|10\rangle + |11\rangle)_{CD}, \\ |\psi_2\rangle &= \frac{1}{\sqrt{2}} (|00\rangle + |11\rangle)_{CD}. \end{aligned} \quad (28)$$

The state $|\psi_1\rangle$ in its Schmidt form is

$$|\psi_1\rangle = \cos \beta |0'\rangle_C |0'\rangle_D + \sin \beta |1'\rangle_C |1'\rangle_D, \quad (29)$$

where $\cos \beta = \frac{1+t}{\sqrt{2+2t^2}}$, $\sin \beta = \frac{1-t}{\sqrt{2+2t^2}}$. Here, $\{|i'\rangle_C\}$, $\{|i'\rangle_D\}$, $i \in \{0, 1\}$ are the corresponding new bases for C and D . (See detail in Appendix C).

If $t = 1$, $|\psi_1\rangle$ is not an entangled state and the lack of nonlocality may result in the failure of the self-testing. Following the framework of [32], we intend to divide the four parties into two parts, and one of them performs local measurements on $|\Psi_2\rangle$. If we divide $ABCD$ randomly into groups that each have two parties, for example, AB and CD , as a result, the projection measurements may collapse the state shared by the remaining parts into some unknown pure bipartite entangled states. Then the remaining two parts should check whether the projected state they share violates maximally Equation (3) for the appropriate α . Without loss of generality, if A and B perform the measurement in the σ_z bases, $|\psi_1\rangle$ and $|\psi_2\rangle$ should be self-tested by C and D , respectively, and simultaneously conditioned on the outcomes "00" and "11".

Following the result given in Lemma 1, $|\psi_1\rangle$ can be self-tested by reaching the maximal violation of the tilted CHSH Bell inequality

$$b(\alpha) = \sqrt{8+2\alpha^2} = \frac{2\sqrt{2}(t+1)}{\sqrt{1+t^2}}, \quad (30)$$

where $\alpha = 2\sqrt{\frac{1-\sin^2 2\beta}{1+\sin^2 2\beta}}$ and the optimal measurement are set as Lemma 1 with $\tan \mu = \frac{1-t^2}{1+t^2}$. Meanwhile, $|\psi_2\rangle$ is still a maximally entangled two-qubit state under the same transformation of bases

$$|\psi_2\rangle = \frac{1}{\sqrt{2}}(|0'0'\rangle + |1'1'\rangle)_{CD}, \tag{31}$$

and hence, we can use the same measurement settings as $|\psi_1\rangle$. As the definition given in Lemma 3, $\alpha_{00} = \mu, \alpha_{01} = -\mu, \alpha_{10} = \frac{\pi}{2} - \mu, \alpha_{11} = -\frac{\pi}{2} - \mu$, and thus it will satisfy the condition $\alpha_{00} + \alpha_{10} = \alpha_{01} - \alpha_{11}$.

Define

$$f(t) = \begin{cases} 0, & t < 1 \\ 1, & t > 1 \end{cases}. \tag{32}$$

Then $|\psi_1\rangle$ can be self-tested by winning the XOR game and we give the criterion to self-test $|\Psi_2\rangle$ as the following Result 2.

Result 2 (See proof in Appendix B). Consider four spatially separated parties, Alice, Bob, Charlie and David, each performing five measurements with binary outcomes denoted as $A_i, B_j, C_k, D_l (i, j, k, l \in \{0, 1, 2, 3, 4\})$ on an unknown shared quantum state $|\Psi_2\rangle$. The target state $|\Psi_2\rangle$ is self-tested if the statistics are observed as the following

$$\begin{cases} \langle P_A^0 P_B^0 P_C^0 P_D^0 \rangle = \langle P_A^0 P_B^0 P_C^1 P_D^1 \rangle = \langle P_A^0 P_B^1 P_C^0 P_D^1 \rangle = \langle P_A^0 P_B^1 P_C^1 P_D^0 \rangle = \langle P_A^1 P_B^0 P_C^0 P_D^1 \rangle \\ \qquad \qquad \qquad = \langle P_A^1 P_B^0 P_C^1 P_D^0 \rangle = \langle P_A^1 P_B^1 P_C^0 P_D^0 \rangle = \langle P_A^1 P_B^1 P_C^1 P_D^1 \rangle = \frac{1}{8 + 4t^2} \\ \langle P_A^0 P_B^0 P_C^0 P_D^1 \rangle = \langle P_A^0 P_B^0 P_C^1 P_D^0 \rangle = \langle P_A^0 P_B^1 P_C^0 P_D^0 \rangle = \langle P_A^1 P_B^0 P_C^0 P_D^0 \rangle = \frac{t^2}{8 + 4t^2} \end{cases}, \tag{33}$$

$$\begin{cases} \langle P_M^0 P_N^0 \mathcal{B}(\alpha, Q_0, Q_1, R_2, R_3) \rangle = \frac{t^2 + 1}{2t^2 + 4} \sqrt{8 + 2\alpha^2} \\ \langle P_M^0 P_N^0 Q_0 (R_2 - R_3) \rangle = 0 \\ \frac{\langle P_M^0 P_N^0 R_2 \rangle}{2 \sin \mu} - \frac{\langle P_M^0 P_N^0 R_3 \rangle}{2 \sin \mu} = (-1)^{f(t)} \langle P_M^0 P_N^0 R_1 \rangle \end{cases}, \tag{34}$$

$$\begin{cases} \sum_{i \in (0,1), j \in (2,3)} f_{ij} \langle P_A^1 P_B^1 C_i Q_j \rangle = \frac{2}{(2 + t^2) \sin 2\mu} \\ \langle P_A^1 P_B^1 C_0 (D_2 - D_3) \rangle = 0 \\ \frac{\langle P_A^1 P_B^1 D_2 \rangle}{2 \sin \mu} - \frac{\langle P_A^1 P_B^1 D_3 \rangle}{2 \sin \mu} = (-1)^{f(t)} \langle P_A^1 P_B^1 D_1 \rangle \end{cases}, \tag{35}$$

$$\langle P_M^0 P_N^0 P_Q^0 P_R^0 R_0 \rangle = \frac{t}{4t^2 + 8}, \tag{36}$$

where $(M, N, Q, R) \in \{(A, B, C, D), (A, C, B, D), (A, D, B, C), (B, C, D, A), (B, D, A, C), (C, D, A, B)\}$, $P_s^0 \triangleq P_{Z_s=+1} = \frac{1+Z_s}{2}, P_s^1 \triangleq P_{Z_s=-1} = \frac{1-Z_s}{2}$, where $s \in \{A, B, C, D\}$ are projectors

for the Z_s measurement and $\sin \mu = \frac{|1-t|}{\sqrt{2+2t^2}}$, $\cos \mu = \frac{1+t}{\sqrt{2+2t^2}}$. At the same time, we find a proper construction of the local isometry Φ , where Z_s and X_s are based on the measurement settings

$$\begin{aligned} Z_A = A_1 &= (-1)^{f(t)} \frac{A_2 - A_3}{2 \sin \mu}, & X_A = A_0 &= \frac{A_2 + A_3}{2 \cos \mu}, \\ Z_B = B_1 &= (-1)^{f(t)} \frac{B_2 - B_3}{2 \sin \mu}, & X_B = B_0 &= \frac{B_2 + B_3}{2 \cos \mu}, \\ Z_C = C_1 &= (-1)^{f(t)} \frac{C_2 - C_3}{2 \sin \mu}, & X_C = C_0 &= \frac{C_2 + C_3}{2 \cos \mu}, \\ Z_D = D_1 &= (-1)^{f(t)} \frac{D_2 - D_3}{2 \sin \mu}, & X_D = D_0 &= \frac{D_2 + D_3}{2 \cos \mu}, \end{aligned} \tag{37}$$

and thus makes the protocol device-independent. In addition, each party may need another fifth measurements $A_4 = Z_A X_A, B_4 = Z_B X_B, C_4 = Z_C X_C, D_4 = Z_D X_D$ to obtain the observation of Equation (36). Since $\sigma_Z \sigma_X = i \sigma_Y$, the fifth measurements are feasible in practical experiments.

5. Conclusions

In this paper, we propose schemes to self-test a large family of four-qubit symmetric states. The target states we focus on are the superposition of the four-qubit Dicke states.

We first present a procedure for self-testing of a particular four-qubit symmetric state with a special structure, and this procedure makes use of the self-testing of the maximally entangled two-qubit state $\frac{|00\rangle + |11\rangle}{\sqrt{2}}$. At the same time, we prove that this protocol is robust against inevitable experimental errors based on norm inequality. In addition, we propose an approach to self-test a one-parameter family of four-qubit pure states through projections onto two systems. Here in our work, only the simplest Pauli measurements are used, which is quite helpful in the experiments.

It would also be of interest to work on a more general state with two parameters by using the swap method and semidefinite programming (SDP) [26] in the form

$$|\Psi\rangle = \cos \theta \cos \rho |GHZ\rangle + \cos \theta \sin \rho |S_{4,2}\rangle + \sin \theta |W\rangle, \tag{38}$$

where $\theta \in [0, \frac{\pi}{2}], \rho \in [0, \frac{\pi}{2}]$, which may provide better robustness than the analytical bounds. What is more, our work could potentially be generalized to a higher dimension scenario. These are reserved for our future work.

Author Contributions: Methodology, validation, investigation, resources, writing—original draft preparation, D.B.; writing—review and editing, project administration, funding acquisition, supervision, X.T.; discussion, Q.X., H.W. and R.H. All authors have read and agreed to the published version of the manuscript.

Funding: The research was partly funded by the Natural Science Foundation of Guangdong Province of China under Grant No. 2021A1515011440, the Major Program of Guangdong Basic and Applied Research under Grant No. 2019B030302008, the National Natural Science Foundation of China under Grant No. 62032009, and the Outstanding Innovative Talents Cultivation Funded Programs for Doctoral Students of Jinan University under Grant No. 2021CXB007.

Informed Consent Statement: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest.

Appendix A. Proof of the Robustness

In this section, we give the proof of Equation (25) based on the L_2 norm. Rewrite the norm Equation (24) as

$$\begin{aligned} \| |\tilde{\Psi}_1\rangle - |junk\rangle \otimes |\Psi'_1\rangle \| &= \| |\tilde{\Psi}_1\rangle - |\Psi_1^*\rangle + |\Psi_1^*\rangle - |junk\rangle \otimes |\Psi'_1\rangle \| \\ &\leq \| |\tilde{\Psi}_1\rangle - |\Psi_1^*\rangle \| + \| |\Psi_1^*\rangle - |junk\rangle \otimes |\Psi'_1\rangle \|. \end{aligned} \tag{A1}$$

Obviously, we need to find the upper bounds for $\|\lvert\Psi_1\rangle - \lvert\Psi_1^*\rangle\|$ and $\|\lvert\Psi_1^*\rangle - \lvert junk\rangle \otimes \lvert\Psi_1'\rangle\|$ respectively. Suppose each observation in Equations (15) and (16) has a deviation at most ϵ around the ideal value. Then we can obtain some inequalities, for instance

$$\begin{aligned} \frac{1}{8} - \epsilon \leq \langle P_A^0 P_B^0 P_C^0 P_D^0 \rangle \leq \frac{1}{8} + \epsilon, \quad \langle P_A^0 P_B^0 X_C X_D \rangle \geq \frac{1}{4} - \epsilon, \quad \langle P_A^0 P_B^0 Z_C Z_D \rangle \geq \frac{1}{4} - \epsilon, \\ \langle P_A^0 P_B^0 X_C Z_D \rangle \leq \epsilon, \quad \langle P_A^0 P_B^0 X_C M_D \rangle \geq \frac{1}{4\sqrt{2}} - \epsilon, \quad \langle P_A^0 P_B^0 Z_C M_D \rangle \geq \frac{1}{4\sqrt{2}} - \epsilon. \end{aligned} \tag{A2}$$

In addition, for convenience and rigorous of the derivation, we assume that

$$\begin{aligned} \langle P_A^0 P_B^0 P_C^0 P_D^1 \rangle \leq \epsilon, \langle P_A^0 P_B^0 P_C^1 P_D^0 \rangle \leq \epsilon, \langle P_A^0 P_B^1 P_C^0 P_D^0 \rangle \leq \epsilon, \langle P_A^1 P_B^0 P_C^0 P_D^0 \rangle \leq \epsilon, \\ \langle P_A^0 P_B^1 P_C^1 P_D^1 \rangle \leq \epsilon, \langle P_A^1 P_B^1 P_C^0 P_D^1 \rangle \leq \epsilon, \langle P_A^1 P_B^0 P_C^1 P_D^1 \rangle \leq \epsilon, \langle P_A^1 P_B^1 P_C^1 P_D^0 \rangle \leq \epsilon, \end{aligned} \tag{A3}$$

which may not direct the observation statistics. We can now write

$$\begin{aligned} & \| (P_A^0 P_B^0 X_C - P_A^0 P_B^0 X_D) \lvert\Psi_1\rangle \| \\ &= \sqrt{ \langle \Psi_1 \lvert P_A^0 P_B^0 X_C X_C P_A^0 P_B^0 + P_A^0 P_B^0 X_D X_D P_A^0 P_B^0 - 2 P_A^0 P_B^0 X_C X_D P_A^0 P_B^0 \lvert\Psi_1\rangle } \\ &= \sqrt{ \langle \Psi_1 \lvert P_A^0 P_B^0 X_C X_C P_A^0 P_B^0 \lvert\Psi_1\rangle + \langle \Psi_1 \lvert P_A^0 P_B^0 X_D X_D P_A^0 P_B^0 \lvert\Psi_1\rangle - 2 \langle \Psi_1 \lvert P_A^0 P_B^0 X_C X_D P_A^0 P_B^0 \lvert\Psi_1\rangle } \\ &\leq \sqrt{ \frac{1}{4} + 4\epsilon + \frac{1}{4} + 4\epsilon - 2(\frac{1}{4} - \epsilon) } = \sqrt{10\epsilon} = \epsilon_1, \end{aligned} \tag{A4}$$

and similarly,

$$\| (P_A^0 P_B^0 Z_C - P_A^0 P_B^0 Z_D) \lvert\Psi_1\rangle \| \leq \sqrt{10\epsilon}. \tag{A5}$$

In addition,

$$\begin{aligned} & \| P_A^0 P_B^0 \frac{X_C + Z_C}{\sqrt{2}} \lvert\Psi_1\rangle \| \\ &= \sqrt{ \frac{1}{2} \langle \Psi_1 \lvert P_A^0 P_B^0 X_C X_C P_A^0 P_B^0 + P_A^0 P_B^0 X_D X_D P_A^0 P_B^0 + 2 P_A^0 P_B^0 X_C Z_C P_A^0 P_B^0 \lvert\Psi_1\rangle } \\ &\leq \sqrt{ \frac{1}{2} [\frac{1}{4} + 4\epsilon + \frac{1}{4} + 4\epsilon + 2(\epsilon + \sqrt{10\epsilon(\frac{1}{4} + 4\epsilon)})] } = \sqrt{ \frac{1}{4} + 5\epsilon + \sqrt{10\epsilon(\frac{1}{4} + 4\epsilon)} }, \end{aligned} \tag{A6}$$

where $\langle \Psi \lvert P_A^0 P_B^0 X_C Z_C P_A^0 P_B^0 \lvert\Psi_1\rangle \leq \epsilon + \sqrt{10\epsilon(\frac{1}{4} + 4\epsilon)}$ from

$$\langle \Psi_1 \lvert P_A^0 P_B^0 X_C Z_C P_A^0 P_B^0 \lvert\Psi_1\rangle - \langle \Psi_1 \lvert P_A^0 P_B^0 X_C Z_D P_A^0 P_B^0 \lvert\Psi_1\rangle \leq \sqrt{10\epsilon(\frac{1}{4} + 4\epsilon)} \tag{A7}$$

by using the Cauchy–Schwarz inequality [33] and Equation (A5). Hence, we obtain

$$\begin{aligned} & \| P_A^0 P_B^0 M_D \lvert\Psi_1\rangle - P_A^0 P_B^0 \frac{X_C + Z_C}{\sqrt{2}} \lvert\Psi_1\rangle \| \\ &= \sqrt{ \langle \Psi_1 \lvert P_A^0 P_B^0 M_D M_D - \sqrt{2} P_A^0 P_B^0 (M_D X_C + M_D Z_C) \lvert\Psi_1\rangle + \left\| \frac{P_A^0 P_B^0 X_C + P_A^0 P_B^0 Z_C}{\sqrt{2}} \lvert\Psi_1\rangle \right\|^2 } \\ &\leq \sqrt{ \frac{1}{4} + 4\epsilon + \frac{1}{4} + 5\epsilon + \sqrt{10\epsilon(\frac{1}{4} + 4\epsilon)} - 2\sqrt{2} \times (\frac{1}{4\sqrt{2}} - \epsilon) } \\ &= \sqrt{ (9 + 2\sqrt{2})\epsilon + \sqrt{10\epsilon(\frac{1}{4} + 4\epsilon)} } = \epsilon'. \end{aligned} \tag{A8}$$

Since the norm of the projectors is equal to 1, we have

$$\begin{aligned}
 & \|P_A^0 P_B^0 (M_D)^2 |\Psi_1\rangle - P_A^0 P_B^0 M_D \frac{X_C + Z_C}{\sqrt{2}} |\Psi_1\rangle\| \\
 & \leq \|P_A^0 P_B^0 M_D\|_\infty \|P_A^0 P_B^0 M_D |\Psi_1\rangle - P_A^0 P_B^0 \frac{X_C + Z_C}{\sqrt{2}} |\Psi_1\rangle\| \\
 & \leq \|P_A^0\|_\infty \|P_B^0\|_\infty \|M_D\|_\infty \|P_A^0 P_B^0 M_D |\Psi_1\rangle - P_A^0 P_B^0 \frac{X_C + Z_C}{\sqrt{2}} |\Psi_1\rangle\| \tag{A9} \\
 & = \|P_A^0\|_\infty \|P_B^0\|_\infty \|P_M^0 - P_M^1\|_\infty \|P_A^0 P_B^0 M_D |\Psi_1\rangle - P_A^0 P_B^0 \frac{X_C + Z_C}{\sqrt{2}} |\Psi_1\rangle\| \\
 & \leq \|P_A^0\|_\infty \|P_B^0\|_\infty (\|P_M^0\|_\infty + \|P_M^1\|_\infty) \|P_A^0 P_B^0 M_D |\Psi_1\rangle - P_A^0 P_B^0 \frac{X_C + Z_C}{\sqrt{2}} |\Psi_1\rangle\| = 2\epsilon'.
 \end{aligned}$$

Similarly,

$$\|P_A^0 P_B^0 \frac{X_C + Z_C}{\sqrt{2}} M_D |\Psi_1\rangle - P_A^0 P_B^0 (\frac{X_C + Z_C}{\sqrt{2}})^2 |\Psi_1\rangle\| \leq 2\sqrt{2}\epsilon', \tag{A10}$$

which implies

$$\begin{aligned}
 & \|P_A^0 P_B^0 |\Psi_1\rangle - P_A^0 P_B^0 (\frac{X_C + Z_C}{\sqrt{2}})^2 |\Psi_1\rangle\| \leq (2 + 2\sqrt{2})\epsilon' \\
 \implies & \|P_A^0 P_B^0 X_C Z_C |\Psi_1\rangle + P_A^0 P_B^0 Z_C X_C |\Psi_1\rangle\| \leq 2(2 + 2\sqrt{2})\epsilon'.
 \end{aligned} \tag{A11}$$

Finally, since

$$\begin{cases} \|P_A^0 P_B^0 Z_C X_C |\Psi_1\rangle - P_A^0 P_B^0 Z_C X_D |\Psi_1\rangle\| \leq \sqrt{10\epsilon(\frac{1}{4} + 4\epsilon)} \\ \|P_A^0 P_B^0 X_D Z_C |\Psi_1\rangle - P_A^0 P_B^0 X_D Z_D |\Psi_1\rangle\| \leq \sqrt{10\epsilon(\frac{1}{4} + 4\epsilon)} \end{cases} \tag{A12}$$

$$\implies \|P_A^0 P_B^0 Z_C X_C |\Psi_1\rangle - P_A^0 P_B^0 X_D Z_D |\Psi_1\rangle\| \leq 2\sqrt{10\epsilon(\frac{1}{4} + 4\epsilon)}. \tag{A13}$$

Similarly,

$$\|P_A^0 P_B^0 X_C Z_C |\Psi_1\rangle - P_A^0 P_B^0 Z_D X_D |\Psi_1\rangle\| \leq 2\sqrt{10\epsilon(\frac{1}{4} + 4\epsilon)}, \tag{A14}$$

therefore we can obtain

$$\begin{aligned}
 & \|P_A^0 P_B^0 X_D Z_D |\Psi_1\rangle + P_A^0 P_B^0 Z_D X_D |\Psi_1\rangle\| \leq 2(2 + 2\sqrt{2})\epsilon' + 4\sqrt{10\epsilon(\frac{1}{4} + 4\epsilon)} \\
 & = 2(2 + 2\sqrt{2})\sqrt{(9 + 2\sqrt{2})\epsilon + \sqrt{10\epsilon(\frac{1}{4} + 4\epsilon)}} + 4\sqrt{10\epsilon(\frac{1}{4} + 4\epsilon)} \tag{A15} \\
 & \leq 2(2 + \sqrt{2})(10\epsilon)^{\frac{1}{4}} (\frac{9 + 2\sqrt{2} + 2\sqrt{10}}{\sqrt{10}} (\epsilon)^{\frac{1}{2}} + 1) + 4(\frac{\sqrt{10}}{2} (\epsilon)^{\frac{1}{2}} + 2\sqrt{10}\epsilon) = 2\epsilon_2.
 \end{aligned}$$

Hence, we obtain

$$\begin{cases} \|P_i^0 P_j^0 X_k |\Psi_1\rangle - P_i^0 P_j^0 X_l |\Psi_1\rangle\| \leq \epsilon_1 \\ \|P_i^0 P_j^0 Z_k |\Psi_1\rangle - P_i^0 P_j^0 Z_l |\Psi_1\rangle\| \leq \epsilon_1 \\ \|P_i^0 P_j^0 X_k Z_k |\Psi_1\rangle + P_i^0 P_j^0 Z_k X_k |\Psi_1\rangle\| \leq 2\epsilon_2' \\ \|P_i^0 P_j^0 X_l Z_l |\Psi_1\rangle + P_i^0 P_j^0 Z_l X_l |\Psi_1\rangle\| \leq 2\epsilon_2 \end{cases} \tag{A16}$$

and

$$\begin{cases} \|P_A^1 P_B^1 X_C |\Psi_1\rangle - P_A^1 P_B^1 X_D |\Psi_1\rangle\| \leq \epsilon_1 \\ \|P_A^1 P_B^1 Z_C |\Psi_1\rangle - P_A^1 P_B^1 Z_D |\Psi_1\rangle\| \leq \epsilon_1 \\ \|P_A^1 P_B^1 X_C Z_C |\Psi_1\rangle + P_A^1 P_B^1 Z_C X_C |\Psi_1\rangle\| \leq 2\epsilon_2 \\ \|P_A^1 P_B^1 X_D Z_D |\Psi_1\rangle + P_A^1 P_B^1 Z_D X_D |\Psi_1\rangle\| \leq 2\epsilon_2 \end{cases} \tag{A17}$$

where $(i, j, k, l) = \{(A, B, C, D), (A, C, B, D), (A, D, B, C), (B, C, A, D), (B, D, A, C), (C, D, A, B)\}$. In addition, we have

$$\begin{aligned} & \|X_D P_A^0 P_B^0 P_C^0 P_D^1 |\Psi_1\rangle |0001\rangle + X_C P_A^0 P_B^0 P_C^1 P_D^0 |\Psi_1\rangle |0010\rangle \\ & + X_B P_A^0 P_B^1 P_C^0 P_D^0 |\Psi_1\rangle |0100\rangle + X_A P_A^1 P_B^0 P_C^0 P_D^0 |\Psi_1\rangle |1000\rangle \\ & + X_B X_C X_D P_A^0 P_B^1 P_C^1 P_D^1 |\Psi_1\rangle |0111\rangle + X_A X_C X_D P_A^1 P_B^0 P_C^1 P_D^1 |\Psi_1\rangle |1011\rangle \\ & + X_A X_B X_D P_A^1 P_B^1 P_C^0 P_D^1 |\Psi_1\rangle |1101\rangle + X_A X_B X_C P_A^1 P_B^1 P_C^1 P_D^0 |\Psi_1\rangle |1110\rangle \| \\ \leq & \|X_D P_A^0 P_B^0 P_C^0 P_D^1 |\Psi_1\rangle |0001\rangle\| + \|X_C P_A^0 P_B^0 P_C^1 P_D^0 |\Psi_1\rangle |0010\rangle\| \\ & + \|X_B P_A^0 P_B^1 P_C^0 P_D^0 |\Psi_1\rangle |0100\rangle\| + \|X_A P_A^1 P_B^0 P_C^0 P_D^0 |\Psi_1\rangle |1000\rangle\| \\ & + \|X_B X_C X_D P_A^0 P_B^1 P_C^1 P_D^1 |\Psi_1\rangle |0111\rangle\| + \|X_A X_C X_D P_A^1 P_B^0 P_C^1 P_D^1 |\Psi_1\rangle |1011\rangle\| \\ & + \|X_A X_B X_D P_A^1 P_B^1 P_C^0 P_D^1 |\Psi_1\rangle |1101\rangle\| + \|X_A X_B X_C P_A^1 P_B^1 P_C^1 P_D^0 |\Psi_1\rangle |1110\rangle\| \\ = & |\langle P_A^0 P_B^0 P_C^0 P_D^1 \rangle| + |\langle P_A^0 P_B^0 P_C^1 P_D^0 \rangle| + |\langle P_A^0 P_B^1 P_C^0 P_D^0 \rangle| + |\langle P_A^1 P_B^0 P_C^0 P_D^0 \rangle| \\ & + |\langle P_A^0 P_B^1 P_C^1 P_D^1 \rangle| + |\langle P_A^1 P_B^0 P_C^1 P_D^1 \rangle| + |\langle P_A^1 P_B^1 P_C^0 P_D^1 \rangle| + |\langle P_A^1 P_B^1 P_C^1 P_D^0 \rangle| \leq 8\epsilon. \end{aligned} \tag{A18}$$

With a similar derivation in [34], we have $|\langle P_A^0 P_B^0 Z_C \rangle| \leq \epsilon_1 + \epsilon_2$ and $|\langle P_A^0 P_B^0 Z_D \rangle| \leq \epsilon_1 + \epsilon_2$, which implies that

$$\begin{aligned} & \|P_A^0 P_B^0 (1 + Z_C)(1 + Z_D)\| \\ = & 2\sqrt{\langle P_A^0 P_B^0 \rangle + \langle P_A^0 P_B^0 Z_C \rangle + \langle P_A^0 P_B^0 Z_D \rangle + \langle P_A^0 P_B^0 Z_C Z_D \rangle} \\ \leq & 2\sqrt{\frac{1}{4} + 4\epsilon + \epsilon_1 + \epsilon_2 + \epsilon_1 + \epsilon_2 + \frac{1}{4} + \epsilon}, \end{aligned} \tag{A19}$$

and thus

$$\begin{aligned} \left\| \frac{(1 + Z_A)(1 + Z_B)(1 + Z_C)(1 + Z_D)}{4\sqrt{2}} \right\| & = \left\| \frac{P_A^0 P_B^0 (1 + Z_C)(1 + Z_D)}{\sqrt{2}} \right\| \\ & \leq \sqrt{1 + 10\epsilon + 4(\epsilon_1 + \epsilon_2)} \leq 1 + 5\epsilon + 2(\epsilon_1 + \epsilon_2). \end{aligned} \tag{A20}$$

We now can write

$$\begin{aligned} \|\tilde{\Psi}_1 - |\Psi_1^*\rangle\| & \leq 8 \times (\epsilon_1 + \epsilon_2) + 8\epsilon = 8(\epsilon_1 + \epsilon_2 + \epsilon), \\ \|\Psi_1^* - |junk\rangle \otimes |\Psi_1'\rangle\| & \leq 5\epsilon + 2(\epsilon_1 + \epsilon_2), \end{aligned} \tag{A21}$$

which implies

$$f(\epsilon) = 13\epsilon + 10(\epsilon_1 + \epsilon_2) = 265.98\epsilon + 348.45\epsilon^{\frac{3}{4}} + 94.87\epsilon^{\frac{1}{2}} + 60.70\epsilon^{\frac{1}{4}}. \tag{A22}$$

Appendix B. Proof of the Self-Testing of a Family of Parameterized Four-Qubit Symmetric States

Observation Equation (33) implies that

$$\begin{aligned} &\langle P_A^0 P_B^0 P_C^0 P_D^0 \rangle + \langle P_A^0 P_B^0 P_C^0 P_D^1 \rangle + \langle P_A^0 P_B^0 P_C^1 P_D^0 \rangle + \langle P_A^0 P_B^1 P_C^0 P_D^0 \rangle + \langle P_A^1 P_B^0 P_C^0 P_D^0 \rangle \\ &+ \langle P_A^0 P_B^0 P_C^1 P_D^1 \rangle + \langle P_A^0 P_B^1 P_C^0 P_D^1 \rangle + \langle P_A^0 P_B^1 P_C^0 P_D^1 \rangle + \langle P_A^1 P_B^0 P_C^0 P_D^1 \rangle + \langle P_A^1 P_B^0 P_C^1 P_D^0 \rangle \\ &+ \langle P_A^1 P_B^1 P_C^0 P_D^0 \rangle + \langle P_A^1 P_B^1 P_C^1 P_D^1 \rangle = 1, \end{aligned} \tag{A23}$$

and thus $P_A^a P_B^b P_C^c P_D^d |\Psi_2\rangle = 0$ for other four projectors.

For convenience, we use $(M, N, Q, R) = (A, B, C, D)$ as an example to prove Result 2. Define the operators for party C and D as

$$\begin{aligned} X_C &= C_0, Z_C = C_1, Z'_C = C_0, X'_C = C_1, \\ X_D &= D_0, Z_D = D_1, Z'_D = \frac{D_2 + D_3}{2 \cos \mu}, X'_D = \frac{D_2 - D_3}{2 \sin \mu}. \end{aligned} \tag{A24}$$

The maximal violation of the tilted CHSH inequality as Equation (34) implies

$$P_A^0 P_B^0 Z'_C |\Psi_2\rangle = P_A^0 P_B^0 Z'_D |\Psi_2\rangle, \tag{A25a}$$

$$P_A^0 P_B^0 Z'_C X'_C |\Psi_2\rangle = -P_A^0 P_B^0 X'_C Z'_C |\Psi_2\rangle, \tag{A25b}$$

$$P_A^0 P_B^0 X'_C (I + Z'_D) |\Psi_2\rangle = \frac{1}{\tan \beta} P_A^0 P_B^0 X'_D (I - Z'_C) |\Psi_2\rangle. \tag{A25c}$$

Then we have

$$\begin{aligned} P_A^0 P_B^0 X_C |\Psi_2\rangle &= P_A^0 P_B^0 X_D |\Psi_2\rangle, \\ P_A^0 P_B^0 X_C Z_C |\Psi_2\rangle &= -P_A^0 P_B^0 Z_C X_C |\Psi_2\rangle \end{aligned} \tag{A26}$$

by Equations (A25a) and (A25b). The observation of Equation (34) implies

$$P_A^0 P_B^0 Z'_C |\Psi_2\rangle \perp P_A^0 P_B^0 X'_D |\Psi_2\rangle, \tag{A27}$$

and combined with the relation Equation (A25a) from the tilted CHSH inequality, we have

$$P_A^0 P_B^0 Z'_D |\Psi_2\rangle \perp P_A^0 P_B^0 X'_D |\Psi_2\rangle. \tag{A28}$$

We can write the $Z_D |\Psi_2\rangle$ in the subspace of $P_A^0 P_B^0$ as

$$P_A^0 P_B^0 Z_D |\Psi_2\rangle = (-1)^{f(t)} P_A^0 P_B^0 X'_D |\Psi_2\rangle \tag{A29}$$

by Equation (34) and thus we can define the vector $X_D |\Psi_2\rangle$ orthogonal to $Z_D |\Psi_2\rangle$ as

$$P_A^0 P_B^0 X_D |\Psi_2\rangle = P_A^0 P_B^0 Z'_D |\Psi_2\rangle. \tag{A30}$$

From Equations (A25a) and (A25c), we obtain

$$P_A^0 P_B^0 Z'_D X'_D |\Psi_2\rangle = -P_A^0 P_B^0 X'_D Z'_D |\Psi_2\rangle. \tag{A31}$$

Hence, we obtain

$$P_A^0 P_B^0 X_D Z_D |\Psi_2\rangle = P_A^0 P_B^0 Z'_D X'_D |\Psi_2\rangle = -P_A^0 P_B^0 X'_D Z'_D |\Psi_2\rangle = -P_A^0 P_B^0 Z_D X_D |\Psi_2\rangle. \tag{A32}$$

Similarly, we obtain the following relations

$$\begin{aligned}
 P_M^0 P_N^0 X_Q |\Psi_2\rangle &= P_M^0 P_N^0 X_R |\Psi_2\rangle, \\
 P_M^0 P_N^0 X_Q Z_Q |\Psi_2\rangle &= -P_M^0 P_N^0 Z_Q X_Q |\Psi_2\rangle, \\
 P_M^0 P_N^0 X_R Z_R |\Psi_2\rangle &= -P_M^0 P_N^0 Z_R X_R |\Psi_2\rangle
 \end{aligned}
 \tag{A33}$$

for all $(M, N, Q, R) = \{(A, B, C, D), (A, C, B, D), (A, D, B, C), (B, C, A, D), (B, D, A, C), (C, D, A, B)\}$. The maximal violation of the XOR game Equation (35) implies

$$\begin{aligned}
 P_A^1 P_B^1 Z'_C |\Psi_2\rangle &= P_A^1 P_B^1 Z'_D |\Psi_2\rangle, \\
 P_A^1 P_B^1 X'_C |\Psi_2\rangle &= P_A^1 P_B^1 X'_D |\Psi_2\rangle, \\
 P_A^1 P_B^1 Z'_C X'_C |\Psi_2\rangle &= -P_A^1 P_B^1 X'_C Z'_C |\Psi_2\rangle, \\
 P_A^1 P_B^1 Z'_D X'_D |\Psi_2\rangle &= -P_A^1 P_B^1 X'_D Z'_D |\Psi_2\rangle.
 \end{aligned}
 \tag{A34}$$

We can use a similar method as above and obtain

$$\begin{aligned}
 P_A^1 P_B^1 X_C |\Psi_2\rangle &= P_A^1 P_B^1 X_D |\Psi_2\rangle, \\
 P_A^1 P_B^1 X_C Z_C |\Psi_2\rangle &= -P_A^1 P_B^1 Z_C X_C |\Psi_2\rangle, \\
 P_A^1 P_B^1 X_D Z_D |\Psi_2\rangle &= -P_A^1 P_B^1 Z_D X_D |\Psi_2\rangle.
 \end{aligned}
 \tag{A35}$$

At last, the observation Equation (36) implies that

$$P_M^0 P_N^0 P_Q^0 P_R^0 X_R |\Psi_2\rangle = t P_M^0 P_N^0 P_Q^0 P_R^0 |\Psi_2\rangle
 \tag{A36}$$

for all $(M, N, Q, R) = \{(A, B, C, D), (A, C, B, D), (A, D, B, C), (B, C, A, D), (B, D, A, C), (C, D, A, B)\}$.

We construct the local isometry similar to Figure 1: just replace $|\Psi_1\rangle$ with $|\Psi_2\rangle$. The output after the isometry is

$$\begin{aligned}
 |\tilde{\Psi}_2\rangle &= \Phi(|\Psi_2\rangle |0000\rangle_{A'B'C'D'}) \\
 &= \sum_{a,b,c,d \in \{0,1\}} X_A^a X_B^b X_C^c X_D^d P_A^a P_B^b P_C^c P_D^d |\Psi_2\rangle |abcd\rangle.
 \end{aligned}
 \tag{A37}$$

By using Equation (A26), $X_D P_A^0 P_B^0 P_C^0 P_D^1 |\Psi_2\rangle$ is equal to $P_A^0 P_B^0 P_C^0 P_D^0 X_D |\Psi_2\rangle$. Combining with Equation (A36), one can simplify this term to $t P_A^0 P_B^0 P_C^0 P_D^0 |\Psi_2\rangle$. The third to fifth terms share a similar simplification process.

In addition, $X_C X_D P_A^0 P_B^0 P_C^1 P_D^1 |\Psi_2\rangle$ is equal to $P_A^0 P_B^0 P_C^0 X_C P_D^0 X_D |\Psi_2\rangle$ and then can be replaced by $P_A^0 P_B^0 P_C^0 P_D^0 |\Psi_2\rangle$ using Equation (A33). Terms from the seventh to eleventh are similar. For the last term, we can obtain $P_A^1 P_B^1 P_C^0 P_D^0 |\Psi_2\rangle$ using Equation (A35), which is then the same as the eleventh term. We remind that there are four terms equal to zero. Finally, the output is reduced to

$$\begin{aligned}
 |\Psi_2^*\rangle &= P_A^0 P_B^0 P_C^0 P_D^0 |\Psi_2\rangle (|0000\rangle + t|0001\rangle + t|0010\rangle + t|0100\rangle + t|1000\rangle \\
 &\quad + |0011\rangle + |0101\rangle + |0110\rangle + |1001\rangle + |1010\rangle + |1100\rangle + |1111\rangle),
 \end{aligned}
 \tag{A38}$$

which can be normalized to the form $|junk\rangle \otimes |\Psi_2'\rangle$, here $|junk\rangle = \sqrt{8 + 4t^2} P_A^0 P_B^0 P_C^0 P_D^0 |\Psi_2\rangle$. Then the unknown state $|\Psi_2\rangle$ is self-tested as $|\Psi_2'\rangle$, which proves that Result 2 holds with the required observations Equations (33)–(36). The protocol is also robust by a norm-inequality-based analysis similar to the Result 1 and the detailed derivation process is omitted here.

Appendix C. Relations between Pauli Operators and the Unknown Measurements

In this section, we give details of the relations between Pauli operators and the unknown measurements in Result 2 by Schmidt decomposition.

$$|\psi_1\rangle_{AB} = \frac{1}{\sqrt{2+2t^2}}(|00\rangle + t|01\rangle + t|10\rangle + |11\rangle). \tag{A39}$$

The coefficient matrix of $|\psi_1\rangle$ is

$$A = \frac{1}{\sqrt{2+2t^2}} \begin{pmatrix} 1 & t \\ t & 1 \end{pmatrix}, \tag{A40}$$

which has the Schmidt decomposition $A = USV$, where

$$S = \begin{pmatrix} \frac{1+t}{\sqrt{2+2t^2}} & 0 \\ 0 & \frac{|1-t|}{\sqrt{2+2t^2}} \end{pmatrix} \tag{A41}$$

and

$$\begin{cases} U = V = \begin{pmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \end{pmatrix} & \text{if } t < 1 \\ U = \begin{pmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \end{pmatrix}, V = \begin{pmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{pmatrix} & \text{if } t > 1 \end{cases}. \tag{A42}$$

Hence, if $t < 1$, we have

$$\begin{cases} |0'\rangle_A = \frac{1}{\sqrt{2}}(|0\rangle + |1\rangle)_A \\ |1'\rangle_A = \frac{1}{\sqrt{2}}(|0\rangle - |1\rangle)_A \end{cases}, \begin{cases} |0'\rangle_B = \frac{1}{\sqrt{2}}(|0\rangle + |1\rangle)_B \\ |1'\rangle_B = \frac{1}{\sqrt{2}}(|0\rangle - |1\rangle)_B \end{cases}. \tag{A43}$$

If $t > 1$,

$$\begin{cases} |0'\rangle_A = \frac{1}{\sqrt{2}}(|0\rangle + |1\rangle)_A \\ |1'\rangle_A = \frac{1}{\sqrt{2}}(|0\rangle - |1\rangle)_A \end{cases}, \begin{cases} |0'\rangle_B = \frac{1}{\sqrt{2}}(|0\rangle + |1\rangle)_B \\ |1'\rangle_B = \frac{1}{\sqrt{2}}(|1\rangle - |0\rangle)_B \end{cases}. \tag{A44}$$

Now we can consider the relation between operators Z' and X' with new bases and Pauli operators for part A,

$$\begin{aligned} Z'_A &= |0'\rangle_A \langle 0'|_A - |1'\rangle_A \langle 1'|_A = |0\rangle_A \langle 1|_A + |1\rangle_A \langle 0|_A = \sigma_x, \\ X'_A &= |0'\rangle_A \langle 1'|_A + |1'\rangle_A \langle 0'|_A = |0\rangle_A \langle 0|_A - |1\rangle_A \langle 1|_A = \sigma_z. \end{aligned} \tag{A45}$$

For part B, if $t < 1$,

$$Z'_B = \sigma_x, X'_B = \sigma_z, \tag{A46}$$

and if $t > 1$,

$$Z'_B = \sigma_x, X'_B = -\sigma_z. \tag{A47}$$

Hence, if the operators performed by each party are the same as Lemma 1 with new bases, they can be transformed into Pauli matrices

$$\begin{aligned}
 \sigma_Z = A_1 &= (-1)^{f(t)} \frac{A_2 - A_3}{2 \sin \mu}, \sigma_X = A_0 = \frac{A_2 + A_3}{2 \cos \mu}, \\
 \sigma_Z = B_1 &= (-1)^{f(t)} \frac{B_2 - B_3}{2 \sin \mu}, \sigma_X = B_0 = \frac{B_2 + B_3}{2 \cos \mu}, \\
 \sigma_Z = C_1 &= (-1)^{f(t)} \frac{C_2 - C_3}{2 \sin \mu}, \sigma_X = C_0 = \frac{C_2 + C_3}{2 \cos \mu}, \\
 \sigma_Z = D_1 &= (-1)^{f(t)} \frac{D_2 - D_3}{2 \sin \mu}, \sigma_X = D_0 = \frac{D_2 + D_3}{2 \cos \mu}.
 \end{aligned} \tag{A48}$$

References

1. Dowling, J.P.; Milburn, G.J. Quantum technology: The second quantum revolution. *Philos. Trans. R. Soc. A* **2003**, *361*, 1655–1674. [[CrossRef](#)] [[PubMed](#)]
2. Gisin, N.; Thew, R. Quantum communication. *Nat. Photon.* **2007**, *1*, 165–171. [[CrossRef](#)]
3. Georgescu, I.M.; Ashhab, S.; Nori, F. Quantum simulation. *Rev. Mod. Phys.* **2014**, *86*, 153–158. [[CrossRef](#)]
4. Degen, C.L.; Reinhard, F.; Cappellaro, P. Quantum sensing. *Rev. Mod. Phys.* **2017**, *89*, 035002. [[CrossRef](#)]
5. Yu, C.H.; Gao, F.; Liu, C.; Huynh, D.; Reynolds, M.; Wang, J. Quantum algorithm for visual tracking. *Phys. Rev. A* **2019**, *99*, 022301. [[CrossRef](#)]
6. Liu, H.; Wu, Y.; Wan, L.; Pan, S.; Qin, S.; Gao, F.; Wen, Q. Variational quantum algorithm for Poisson equation. *Phys. Rev. A* **2021**, *104*, 022418. [[CrossRef](#)]
7. Eisert, J.; Hangleiter, D.; Walk, N.; Roth, I.; Markham, D.; Parekh, R.; Chabaud, U.; Kashefi, E. Quantum certification and benchmarking. *Nat. Rev. Phys.* **2020**, *2*, 382–390. [[CrossRef](#)]
8. Cramer, M.; Plenio, M.B.; Flammia, S.T.; Somma, R.; Gross, D.; Bartlett, S.D.; Landon-Cardinal, O.; Poulin, D.; Liu, Y.-K. Efficient quantum state tomography. *Nat. Commun.* **2010**, *1*, 149. [[CrossRef](#)]
9. Mayers, D.; Yao, A. Self testing quantum apparatus. *Quant. Inf. Comput.* **2004**, *4*, 273–286. [[CrossRef](#)]
10. Gallego, R.; Brunner, N.; Hadley, C.; Acín, A. Device-independent tests of classical and quantum dimensions. *Phys. Rev. Lett.* **2010**, *105*, 230501. [[CrossRef](#)]
11. Long, G.; Liu, X. Theoretically efficient high-capacity quantum-key-distribution scheme. *Phys. Rev. A* **2002**, *65*, 032302. [[CrossRef](#)]
12. Kwek, L.; Cao, L.; Luo, W.; Wang, Y.; Sun, S.; Wang, X.; Liu, A. Chip-based quantum key distribution. *Assoc. Asia Pac. Phys. Soc. Bull.* **2021**, *31*, 15. [[CrossRef](#)]
13. Pironio, S.; Acín, A.; Massar, S.; de la Giroday, A.B.; Matsukevich, D.N.; Maunz, P.; Olmschenk, S.; Hayes, D.; Luo, L.; Manning, T.A.; et al. Random numbers certified by Bell’s theorem. *Nature* **2010**, *464*, 1021–1024. [[CrossRef](#)] [[PubMed](#)]
14. Bancal, J. Device-independent witnesses of genuine multipartite entanglement. *Phys. Rev. Lett.* **2011**, *106*, 250404. [[CrossRef](#)]
15. Nielsen, M.A.; Chuang, I.L. *Quantum Computation and Quantum Information*; Cambridge University Press: Cambridge, UK, 2000.
16. Bancal, J.-D.; Navascués, M.; Scarani, V.; Vértesi, T.; Yang, T.H. Physical characterization of quantum devices from nonlocal correlations. *Phys. Rev. A* **2015**, *91*, 022115. [[CrossRef](#)]
17. Popescu, S.; Rohrlich, D. Generic quantum nonlocality. *Phys. Rev. A* **1992**, *166*, 293–297. [[CrossRef](#)]
18. Horodecki, M.; Horodecki, P.; Horodecki, R. Quantum entanglement. *Rev. Mod. Phys.* **2009**, *81*, 865–942. [[CrossRef](#)]
19. Fannes, M.; Lewis, J.T.; Verbeure, A. Symmetric states of composite systems. *Lett. Math. Phys.* **1988**, *15*, 255–260. [[CrossRef](#)]
20. Briegel, H.J.; Browne, D.E.; Dür, W.; Raussendorf, R.; Van den Nest, M. Measurement-based quantum computation. *Nat. Phys.* **2009**, *5*, 19–26. [[CrossRef](#)]
21. Šupić, I.; Coladangelo, A.; Augusiak, R.; Acín, A. Self-testing multipartite entangled states through projections onto two systems. *New J. Phys.* **2018**, *20*, 083041. [[CrossRef](#)]
22. Coladangelo, A.; Goh, K.T.; Scarani, V. All pure bipartite entangled states can be self-tested. *Nat. Commun.* **2017**, *8*, 15485. [[CrossRef](#)] [[PubMed](#)]
23. Yang, T.H.; Navascués, M. Robust self-testing of unknown quantum systems into any entangled two-qubit states. *Phys. Rev. A* **2013**, *87*, 050102. [[CrossRef](#)]
24. Bamps, C.; Pironio, S. Sum-of-squares decompositions for a family of Clauser-Horne-Shimony-Holt-like inequalities and their application to self-testing. *Phys. Rev. A* **2015**, *91*, 052111. [[CrossRef](#)]
25. Wang, Y.; Wu, X.; Scarani, V. All the self-testings of the singlet for two binary measurements. *New J. Phys.* **2016**, *18*, 025021. [[CrossRef](#)]
26. Navascués, M.; Pironio, S.; Acín, A. A convergent hierarchy of semidefinite programs characterizing the set of quantum correlations. *New J. Phys.* **2008**, *10*, 073013. [[CrossRef](#)]
27. Acín, A.; Massar, S.; Pironio, S. Randomness versus nonlocality and entanglement. *Phys. Rev. Lett.* **2012**, *108*, 100402. [[CrossRef](#)]
28. Carteret, H.A.; Higuchi, A.; Sudbery, A. Multipartite generalization of the Schmidt decomposition. *J. Math. Phys.* **2000**, *41*, 7932–7939. [[CrossRef](#)]

29. Long, G. Collapse-in and collapse-out in partial measurement in quantum mechanics and its wise interpretation. *Sci. China Phys. Mech. Astron.* **2021**, *64*, 280321. [[CrossRef](#)]
30. Zhang, X.; Qu, R.; Chang, Z.; Quan, Q.; Gao, H.; Li, F.; Zhang, P. A geometrical framework for quantum incompatibility resources. *Assoc. Asia Pac. Phys. Soc. Bull.* **2022**, *32*, 17. [[CrossRef](#)]
31. Wu, X.; Cai, Y.; Yang, T.H.; Le, H.N.; Bancal, J.-D.; Scarani, V. Robust self-testing of the three-qubit W state. *Phys. Rev. A* **2014**, *90*, 042339. [[CrossRef](#)]
32. Li, X.; Wang, Y.; Han, Y.; Qin, S.; Gao, F.; Wen, Q. Self-Testing of Symmetric Three-Qubit States. *IEEE J. Sel. Areas Commun.* **2020**, *38*, 589–597. [[CrossRef](#)]
33. Bhatia, R.; Davis, C. A Cauchy-Schwarz inequality for operators with applications. *Linear Algebra Appl.* **1995**, *223–224*, 119–129. [[CrossRef](#)]
34. McKague, M.; Yang, T.H.; Scarani, V. Robust self-testing of the singlet. *J. Phys. A Math. Theor.* **2012**, *45*, 455304. [[CrossRef](#)]