

Supplementary Material

New Mild and Simple Approach to Isothiocyanates: A Class of Potent Anticancer Agents

Bingling Luo ^{1,2,†}, Jiankang Wang ^{1,2,†}, Xiaobing Li ¹, Wenhua Lu ¹, Jing Yang ¹, Yumin Hu ¹, Peng Huang ^{1,2} and Shijun Wen ^{1,2,*}



¹ Sun Yat-Sen University Cancer Center; State Key Laboratory of Oncology in South China; Collaborative Innovation Center for Cancer Medicine; Sun Yat-Sen University, 651 Dongfeng East Road, Guangzhou 510060, China; luobl@sysucc.org.cn (B.L.); lixiaobing0629@126.com (X.L.); luwenh@sysucc.org.cn (W.L.); yangjing@sysucc.org.cn (J.Y.); huym@sysucc.org.cn (Y.H.); huangpeng@sysucc.org.cn (P.H.)


² School of Pharmaceutical Sciences, Sun Yat-sen University, 132 Waihuan East Road, Guangzhou 510006, China; wangjk6@mail2.sysu.edu.cn (J.W.)


* Correspondence: wenshj@sysucc.org.cn; Tel.: +86-20-3994-3091


† The authors contributed equally to this work.


PEITC






























































































17 CDCl₃




1100
1000
900
800
700
600
500
400
300
200
100
0

—29.84
—36.48
—46.09

{76.84
77.16
77.48

—114.34
—120.15
127.91
129.48
133.39
138.66
141.77
—148.05
—151.65

17 CDCl₃



f1 (ppm)

10 20 30 40 50 60 70 80 90 100 110 120 130 140 150

