

Title: A Novel Brominated Alkaloid Securidine A, Isolated from the Marine Bryozoan *Securiflustra securifrons*

Supplementary materials: 1D NOE, 13C-NMR, HMBC, ME-HSQC, H2BC, COSY and ROESY correlations of compound Securidine A (**1**)

Authors: Priyanka Siranjeevi ^{1*a}, Kine Ø. Hansen ^{1*a}, Johan Isaksson ², Jeanette H. Andersen ¹ and Espen Hansen ¹.

¹ MARBIO, UiT – The Arctic University of Norway, Brevika, N-9037, Tromsø, Norway; raja.p.siranjeevi@uit.no (P.S); kine.o.hanssen@uit.no (K.Ø.H); jeanette.h.andersen@uit.no (J.H.A); espen.hansen@uit.no (E.H)

² Department of Chemistry, UiT – The Arctic University of Norway, Brevika, N-9037, Tromsø, Norway; johan.isaksson@uit.no (J.I)

* Correspondence: raja.p.siranjeevi@uit.no; Tel.: +47-77649268; kine.o.hanssen@uit.no; Tel.: +47-77649272

^a These authors contributed equally to this work.

Figure S1. 1D NOE spectrum of securidine A (**1**)


Figure S2. 13C-NMR spectrum of **1**.

Figure S3. Combined HSQC and HMBC spectra of **1**.

Figure S4. H2BC spectra of **1**


Figure S5. COSY Spectrum of **1**.

Figure S6. ROESY spectrum of **1**.



Chemical Formula: $C_{14}H_{21}BrN_4O_2$



Exact Mass: 356,08



^1H NMR (600 MHz, Methanol- d_4) δ 8.20 (s, 1H), 7.57 (s, 1H), 7.47 (d, $J = 2.3$ Hz, 1H), 7.22 (dd, $J = 8.6, 2.3$ Hz, 1H), 6.96 (d, $J = 8.8$ Hz, 1H), 3.84 (s, 3H), 3.41 (s, 4H), 3.20 (q, $J = 6.4$ Hz, 2H), 3.18 – 3.10 (m, 2H), 1.60 – 1.49 (m, 4H).

1D NOE + ES

methanol- d_3 , 298 K



^{13}C NMR (151 MHz, MeOD) δ 173.96, 158.77, 156.25, 134.52, 130.48, 130.25, 130.25, 113.10, 112.23, 56.58, 42.41, 42.01, 39.82, 27.52, 26.99.

CARBON

methanol- d_3 , 298 K

