

Communication

# One-pot Fluorination and Organocatalytic Robinson Annulation for Asymmetric Synthesis of Mono- and Difluorinated Cyclohexenones

Xin Huang <sup>1,\*</sup>, Weizhao Zhao <sup>1</sup>, Xiaofeng Zhang <sup>2</sup>, Miao Liu <sup>2</sup>, Stanley N. S. Vasconcelos <sup>3</sup> and Wei Zhang <sup>2,\*</sup>

<sup>1</sup> College of Chemistry and Life Sciences, Zhejiang Normal University, Jinhua 321004, China; zhao792129582@163.com

<sup>2</sup> Department of Chemistry, University of Massachusetts Boston, 100 Morrissey Boulevard, Boston, MA 02125, USA; Xiaofeng.Zhang002@umb.edu (X.Z.); liumiaomarcus@gmail.com (M.L.)

<sup>3</sup> Departamento de Farmácia, Universidade de São Paulo, Av. Prof. Lineu Prestes, 580 São Paulo, SP 05508-000, Brazil; stanleynsv@gmail.com

\* Correspondence: xin.huang@zjnu.cn (X.H.); wei2.zhang@umb.edu (W.Z.); Tel.: +1-617-287-6147 (W.Z.); Fax: +1-617-287-6030 (W.Z.)

Received: 4 August 2018; Accepted: 31 August 2018; Published: 4 September 2018



**Abstract:** A one-pot fluorination and organocatalytic Robinson annulation sequence has been developed for asymmetric synthesis of 6-fluorocyclohex-2-en-1-ones and 4,6-difluorocyclohex-2-en-1-ones. The reactions promoted by cinchona alkaloid amine afforded products bearing two or three stereocenters in good to excellent yields with up to 99% ee and 20:1 dr.

**Keywords:** organocatalysis; fluorination;  $\alpha$ -fluoro- $\beta$ -keto ester; Michael addition; Robinson annulation; cyclohexenone

## 1. Introduction

The development of new synthetic methods for organofluorine compounds is an active topic in organic and medicinal chemistry. Introduction of fluorine atom(s) could have significant impact on molecules' biological activity, bioavailability, and metabolic property [1–4]. Shown in Figure 1 are bioactive fluorinated cyclohexenones [5] including intermediate for antitumor agent COTC (Figure 1A) [6], aromatase inhibitor (Figure 1B) [7], and retinal protein bacteriorhodopsin (Figure 1C) [8].





Figure 1. Bio-active fluorinated cyclohexenones.

The  $\beta$ -ketoester scaffold is a versatile synthon for being both electrophilic and nucleophilic sites [9]. The asymmetric Michael addition reactions of  $\alpha$ -fluoro- $\beta$ -keto esters [10–15] and other monofluorinated nucleophiles [16–19] is an attractive topic. These nucleophiles have been used to react with nitroolefins [10,11,13,14,16,17], *N*-alkyl maleimides [12,15], chalcones [18], and  $\alpha,\beta$ -unsaturated


aldehydes [19]. Beside the Michael addition reactions, some Michael addition-initiated reactions involving  $\alpha$ -fluoro- $\beta$ -keto esters have also been reported [20–22]. Those reactions include Michael/aldol (Robinson) [20], Michael/Michael/aldol [21], and Michael/aza-Henry/lactamization sequences [22] for the construction of cyclohexenones, cyclohexanones, and 2-piperidinones bearing multiple stereocenters.

As part of our recent effort on the synthesis of organofluorine compounds using  $\alpha$ -fluoro- $\beta$ -keto esters [23–25], we have reported a one-pot fluorination/Robinson annulation sequence for fluorinated cyclohexenones **1** (Scheme 1A) [23], Robinson annulation/dehydrofluorination/aromatization sequence for phenols **2** (Scheme 1B) [24], and pyridines [25]. We envisioned that in the presence of an organocatalyst, such as cinchona alkaloid primary amine **cat-1**, the one-pot asymmetric synthesis could be developed for the preparation of monofluorocyclohexenones **3** and difluorocyclohexenones **4** (Scheme 1C).

### Previous work (one-pot reactions)



### This work (one-pot asymmetric reactions)




**Scheme 1.** Robinson annulation-based one-pot synthesis.

## 2. Results and Discussions

A number of six organocatalysts were screened for the asymmetric Robinson annulation reaction of chalcone **6a** and  $\alpha$ -fluoro- $\beta$ -ketoester **5a** (Table 1, entries 1–6). It was found that the reaction with 20 mol% of **cat-1** gave the desired product **3a** in 79% yield with 5:1 dr and 99% ee (entry 1), while other catalysts did not give good results. Acidity of the reaction system plays an important role in amine catalysis for lowering the LUMO energy of iminium ions [26]. Thus, an investigation of the reactions in the presence of different acids was carried out. The best result was achieved by adding  $\text{CF}_3\text{C}_6\text{H}_4\text{CO}_2\text{H}$  to afford **3a** in 89% yield with 9:1 dr and 99% ee (entry 10). We lowered the reaction temperature to  $-20\text{ }^\circ\text{C}$  which slightly increased the dr to 12:1, but significantly decreased the yield to 51% (entry 13). Because we failed to get the crystal of **3** for X-ray structure analysis, the absolute configuration of **3a** was deduced based on the information reported in the literature [27]. The *S*-amine catalyst promoted a

highly selective Re-face attack at the Michael acceptor. We believed that our reactions also catalyzed by the *S*-amine **cat-1** could have the same stereochemistry outcome.


**Table 1.** Screening of catalysts for the Robinson annulation <sup>a</sup>.



Entry	Cat	Additive	Yield (%) <sup>b</sup>	dr <sup>c</sup>	ee (%) <sup>d</sup>
1	<b>cat-1</b>	C <sub>6</sub> H <sub>5</sub> CO <sub>2</sub> H	79	5:1	99
2	<b>cat-2</b>	C <sub>6</sub> H <sub>5</sub> CO <sub>2</sub> H	NR	-	-
3	<b>cat-3</b>	C <sub>6</sub> H <sub>5</sub> CO <sub>2</sub> H	NR	-	-
4	<b>cat-4</b>	C <sub>6</sub> H <sub>5</sub> CO <sub>2</sub> H	NR	-	-
5	<b>cat-5</b>	C <sub>6</sub> H <sub>5</sub> CO <sub>2</sub> H	trace	-	-
6	<b>cat-6</b>	C <sub>6</sub> H <sub>5</sub> CO <sub>2</sub> H	27	3:1	85
7	<b>cat-1</b>	None	42	5:1	92
8	<b>cat-1</b>	AcOH	52	6:1	92
9	<b>cat-1</b>	TsOH	45	5:1	95
10	<b>cat-1</b>	CF <sub>3</sub> C <sub>6</sub> H <sub>4</sub> CO <sub>2</sub> H	89	9:1	99
11	<b>cat-1</b>	CCl <sub>3</sub> CO <sub>2</sub> H	75	8:1	90
12	<b>cat-1</b>	CF <sub>3</sub> CO <sub>2</sub> H	77	8:1	90
13 <sup>e</sup>	<b>cat-1</b>	CF <sub>3</sub> C <sub>6</sub> H <sub>4</sub> CO <sub>2</sub> H	51	12:1	99

<sup>a</sup> Reaction of 0.1 mmol of **5a** and 0.15 mmol of **6a** in 0.5 mL of MeCN. <sup>b</sup> Isolated yield. <sup>c</sup> Determined by <sup>1</sup>H-NMR of crude reaction mixture. <sup>d</sup> Determined by chiral HPLC. <sup>e</sup> Reaction was carried out at −20 °C.


The combination of fluorination and the Robinson annulation as a one-pot reaction for fluorinated cyclohexenone **3a** was then attempted (Table 2). The fluorination step does not need to be asymmetric because both stereocenters were generated during the subsequential Michael addition. The fluorination reaction of β-ketoester **8** with SelectFluor™ was conducted under microwave heating at 120 °C for 20 min without using a catalyst. After the completion of the fluorination, chalcone **6a**, CF<sub>3</sub>C<sub>6</sub>H<sub>4</sub>CO<sub>2</sub>H and catalyst **cat-1** were added to the reaction mixture at 25 °C for the Robinson annulation. But this one-pot fluorination/Robinson annulation only gave **3a** in <10% yield probably due to the effect of the acidic SelectFluor™ derivative in the reaction mixture. To address this issue, bases including Na<sub>2</sub>CO<sub>3</sub>, K<sub>2</sub>CO<sub>3</sub>, and Cs<sub>2</sub>CO<sub>3</sub> were used for the reaction. It was found that addition of 1.5 equiv. of Na<sub>2</sub>CO<sub>3</sub> gave **3a** in 82% yield with a good ee and dr (Table 2, entry 6).

**Table 2.** One-pot synthesis of 6-fluorocyclohex-2-en-1-ones **3**<sup>a</sup>.

Entry	Base (equiv)	Yield (%) <sup>b</sup>	dr <sup>c</sup>	ee (%) <sup>d</sup>
1	None	<10	-	-
2	Na <sub>2</sub> CO <sub>3</sub> (0.5)	42	9:1	99
3	K <sub>2</sub> CO <sub>3</sub> (0.5)	67	3:1	85
4	Cs <sub>2</sub> CO <sub>3</sub> (0.5)	85	1.5:1	52
5	Na <sub>2</sub> CO <sub>3</sub> (1.0)	57	9:1	99
6	Na <sub>2</sub> CO <sub>3</sub> (1.5)	82	9:1	99
7	Na <sub>2</sub> CO <sub>3</sub> (2.0)	83	9:1	99


<sup>a</sup> Reaction of 0.1 mmol of **8**, 0.15 mmol of SelectFlour™, 0.15 mmol of **6a** in 0.5 mL of MeCN. <sup>b</sup> Yield of isolated product **3a**. <sup>c</sup> Determined by <sup>1</sup>H-NMR. <sup>d</sup> Determined by HPLC on Venusil Chiral OD-H column with 90:10 hexane/*i*-PrOH.

With the optimized conditions in hand, the scope of the one-pot synthesis of 6-fluorocyclohexenones **3a–j** using different Michael acceptor **6** was investigated (Scheme 2). Substrates **6** with electron-donating (SMe, Ph) and electron-withdrawing (Br, CF<sub>3</sub>) groups on Ar<sup>1</sup> gave products **3b–f** in 82–90% yields with >96% ee and >9:1 dr. Reaction of a chalcone bearing a thiophene ring also gave product **3h** in a good yield and ee. The introduction of substituents on Ar<sup>2</sup> of **6** gave products **3i** and **3j** with decreased ee, probably due to an unfavorable stereoelectronic effect.

**Scheme 2.** One-pot synthesis of 6-fluorocyclohex-2-en-1-ones **3**.

We have recently reported a one-pot Robinson annulation/dehydrofluorination/aromatization sequence for fluorinated phenols **2** using  $\alpha$ -fluoro- $\beta$ -ketoesters and  $\alpha$ -fluoro- $\alpha,\beta$ -unsaturated ketones as substrates [24]. We envisioned that  $\alpha$ -fluoro- $\alpha,\beta$ -unsaturated ketones **6** could be used for asymmetric synthesis of 4,6-difluorocyclohexanones **4** under organocatalytic conditions. Indeed, one-pot reactions of  $\beta$ -ketoester **8** went smoothly to afford **4a–c** in 53–65% yields with 5:1–8:1 dr and 89–93% ee (Scheme 3).

However, all of these reactions afforded **4a–c** as decarboxylated products, even at low reaction temperatures ( $-30$ – $0$  °C). No aromatization products were observed under the reaction temperature without heating [24]. The products **4a–c** were found not to be stable during the workup and rotary vapor concentration of the crude product.



**Scheme 3.** One-pot synthesis of 4,6-difluorocyclohexanones **4**.

### 3. Conclusions

In summary, a one-pot fluorination and cinchona alkaloid amine-promoted organocatalytic Robinson sequence for asymmetric synthesis of fluorocyclohexenones bearing two stereocenters has been developed. Both stereocenters were established during the step of Michael addition to afford fluorinated cyclohexanones with up to 99% ee and 20:1 dr. Using fluorinated chalcones as Michael acceptors, 4,6-difluorocyclohexanones bearing three stereocenters were also synthesized stereoselectively and in good yields.

**Supplementary Materials:** Supplementary Materials are available online, General procedure; characterization of  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra of products, HPLC spectra of racemic and enantioenriched products.

**Author Contributions:** X.H. developed above reactions. W.Z., X.Z., M.L., and S.N.S.V. expanded the substrate scope. W.Z. conceived and supervised the project.

**Funding:** Joseph P. Healey Research Grant from UMass Boston.

**Acknowledgments:** We would like to thank the Centre for Green Chemistry at the University of Massachusetts Boston, Zhejiang Normal University to support this work.

**Conflicts of Interest:** The authors declare no conflicts of interest

### References

1. Mikami, K.; Itoh, Y.; Yamanaka, M. Fluorinated carbonyl and olefinic compounds: Basic character and asymmetric catalytic reactions. *Chem. Rev.* **2004**, *104*, 1–16. [[CrossRef](#)] [[PubMed](#)]
2. Smits, R.; Cadicamo, C.D.; Burger, K.; Kocsch, B. Synthetic strategies to  $\alpha$ -trifluoromethyl and  $\alpha$ -difluoromethyl substituted  $\alpha$ -amino acids. *Chem. Soc. Rev.* **2008**, *37*, 1727–1739. [[CrossRef](#)] [[PubMed](#)]
3. Wang, J.; Sánchez-Roselló, M.; Acena, J.L.; del Pozo, C.; So-rochinsky, A.E.; Fustero, S.; Soloshonok, V.A.; Liu, H. Fluorine in pharmaceutical industry: Fluorine-containing drugs introduced to the market in the last decade (2001–2011). *Chem. Rev.* **2014**, *114*, 2432–2506. [[CrossRef](#)] [[PubMed](#)]
4. Ojima, I. *Fluorine in Medicinal Chemistry and Chemical Biology*; Wiley-Blackwell: Chichester, UK, 2009.
5. Ho, T.L. *Carbocycle Construction in Terpene Synthesis*; Wiley-VCH: Weinheim, Germany, 1988.

6. Paolo, P.; Manuela, V.; Mario, V.; Antonella, I. 4,5,6,7-Tetrahydroindazole Derivatives as Antitumor Agents. WO Patent 2,000,069,846, 23 December 2000.
7. Roy, P.P.; Roy, K. Molecular docking and QSAR studies of aromatase inhibitor androstenedione derivatives. *J. Pharm. Pharmacol.* **2010**, *62*, 1717–1728. [[CrossRef](#)] [[PubMed](#)]
8. Liu, J.; Shelton, N.L.; Liu, R.S.H. Study of  $\alpha$ -crustacyanin utilizing halogenated canthaxanthins. *Org. Lett.* **2002**, *4*, 2521–2524. [[CrossRef](#)] [[PubMed](#)]
9. Govender, T.; Arvidsson, P.I.; Maguire, G.E.M.; Kruger, H.G.; Naicker, T. Enantioselective organocatalyzed transformations of  $\beta$ -Ketoesters. *Chem. Rev.* **2016**, *116*, 9375–9437. [[CrossRef](#)] [[PubMed](#)]
10. Oh, Y.; Kim, S.M.; Kim, D.Y. Organocatalytic synthesis of quaternary stereocenter bearing a fluorine atom: Enantioselective conjugate addition of  $\alpha$ -fluoro- $\beta$ -ketoesters to nitroalkenes. *Tetrahedron Lett.* **2009**, *50*, 4674–4676. [[CrossRef](#)]
11. Lu, Y.; Zou, G.; Zhao, G. Primary-secondary diamines catalyzed Michael reaction to generate chiral fluorinated quaternary carbon centers. *Tetrahedron* **2015**, *71*, 4137–4144. [[CrossRef](#)]
12. Huang, X.; Yi, W.B.; Ahad, D.; Zhang, W. Recyclable cinchona alkaloid catalyzed asymmetric Michael addition reaction. *Tetrahedron Lett.* **2013**, *54*, 6064–6066. [[CrossRef](#)]
13. Han, Y.; Luo, J.; Liu, C.; Lu, Y. Asymmetric generation of fluorine-containing quaternary carbons adjacent to tertiary stereocenters: Uses of fluorinated methines as nucleophiles. *Chem. Commun.* **2009**, *0*, 2044–2046. [[CrossRef](#)] [[PubMed](#)]
14. Li, H.; Zhang, S.; Yu, C.; Song, X.; Wang, W. Organocatalytic asymmetric synthesis of chiral fluorinated quaternary carbon containing  $\beta$ -ketoesters. *Chem. Commun.* **2009**, *0*, 2136–2138. [[CrossRef](#)] [[PubMed](#)]
15. Jiang, Z.; Pan, Y.; Zhao, Y.; Ma, T.; Lee, R.; Yang, Y.; Huang, K.W.; Wong, M.W.; Tan, C.H. Synthesis of a chiral quaternary carbon center bearing a fluorine atom: Enantio- and diastereoselective guanidine-catalyzed addition of fluorocarbon nucleophiles. *Angew. Chem. Int. Ed.* **2009**, *48*, 3627–3631. [[CrossRef](#)] [[PubMed](#)]
16. Cui, H.-F.; Li, P.; Wang, X.-W.; Zhu, S.-Z.; Zhao, G. Asymmetric Michael addition of  $\alpha$ -fluoro- $\alpha$ -phenylsulfonyl ketones to nitroolefins catalyzed by phenylalanine-based bifunctional thioureas. *J. Fluorine Chem.* **2012**, *133*, 120–126. [[CrossRef](#)]
17. Prakash, G.K.S.; Wang, F.; Stewart, T.; Mathew, T.; Olah, G.A.  $\alpha$ -Fluoro- $\beta$ -nitro(phenylsulfonyl)methane as a fluoromethyl pronucleophile: Efficient stereoselective Michael addition to chalcones. *Proc. Natl. Acad. Sci. USA.* **2009**, *106*, 4090–4094. [[CrossRef](#)] [[PubMed](#)]
18. Furukawa, A.; Shibata, N.; Mizuta, S.; Nakamura, S.; Toru, T.; Shiro, M. Catalytic enantioselective Michael addition of 1-fluorobis(phenylsulfonyl)methane to  $\alpha,\beta$ -unsaturated ketones catalyzed by cinchona alkaloids. *Angew. Chem. Int. Ed.* **2008**, *47*, 8051–8054. [[CrossRef](#)] [[PubMed](#)]
19. Ullah, F.; Zhao, G.L.; Deiana, L.; Zhu, M.; Dziedzic, P.; Ibrahim, I.; Hammar, P.; Sun, J.; Córdova, A. Enantioselective organocatalytic conjugate addition of fluorocarbon nucleophiles to  $\alpha,\beta$ -unsaturated aldehydes. *Chem. Eur. J.* **2009**, *15*, 10013–10017. [[CrossRef](#)] [[PubMed](#)]
20. Cui, H.F.; Yang, Y.Q.; Chai, Z.; Li, P.; Zheng, C.W.; Zhu, S.Z.; Zhao, G. Enantioselective synthesis of functionalized fluorinated cyclohexenones via Robinson annulation catalyzed by primary–secondary diamines. *J. Org. Chem.* **2010**, *75*, 117–122. [[CrossRef](#)] [[PubMed](#)]
21. Huang, X.; Liu, M.; Jasinski, J.P.; Peng, B.; Zhang, W. Recyclable organocatalysts for a one-pot asymmetric synthesis of 2-fluorocyclohexanols bearing six contiguous stereocenters. *Adv. Synth. Catal.* **2017**, *359*, 1919–1926. [[CrossRef](#)]
22. Huang, X.; Pham, K.; Zhang, X.-F.; Yi, W.-B.; Hyatt, J.; Tran, A.; Jasinski, J.P.; Zhang, W. Recyclable organocatalyst for one-pot asymmetric synthesis of fluorinated 2-Piperidinones bearing four contiguous stereogenic centers. *RSC Adv.* **2015**, *5*, 71071–71075. [[CrossRef](#)]
23. Yi, W.B.; Huang, X.; Cai, C.; Zhang, W. One-pot fluorination followed by Michael addition or Robinson annulation for preparation of  $\alpha$ -fluorinated carbonyl compounds. *Green Chem.* **2012**, *14*, 3185–3189. [[CrossRef](#)]
24. Qian, J.; Yi, W.B.; Huang, X.; Miao, Y.; Zhang, J.; Cai, C.; Zhang, W. One-pot synthesis of 3,5-disubstituted and polysubstituted phenols from acyclic precursors. *Org. Lett.* **2015**, *17*, 1090–1093. [[CrossRef](#)] [[PubMed](#)]
25. Song, Z.; Huang, X.; Yi, W.B.; Zhang, W. One-pot reactions for modular synthesis of polysubstituted and fused pyridines. *Org. Lett.* **2016**, *18*, 5640–5643. [[CrossRef](#)] [[PubMed](#)]

26. Brazier, J.B.; Tomkinson, N.C.O. Secondary and primary amine catalysts for iminium catalysis. In *Asymmetric Organocatalysis*; List, B., Ed.; Springer: Berlin/Heidelberg, Germany, 2010; Volume 291, pp. 281–347.
27. Halland, N.; Aburel, P.S.; Jørgensen, K.A. Highly enantio- and diastereoselective organocatalytic asymmetric domino Michael–aldol reaction of  $\beta$ -ketoesters and  $\alpha,\beta$ -unsaturated ketones. *Angew. Chem. Int. Ed.* **2004**, *43*, 1272–1277.

**Sample Availability:** Samples of the compounds are available from the authors.



© 2018 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).