

The Difference in Cytotoxic Activity between Two Optical Isomers of Gelsemine from *Gelsemium elegans* Benth. on PC12 Cells

Li Lin¹, Yan-Chun Liu^{1,2}, and Zhao-Ying Liu^{1,2,*}

¹ College of Veterinary Medicine, Hunan Agricultural University, Changsha 410128, Hunan, China

² Hunan Engineering Technology Research Center of Veterinary Drugs, Hunan Agricultural University, Changsha 410128, Hunan, China

* Correspondence: liu_zhaoying@hunau.edu.cn




Figure S1. The ^{13}C NMR spectrum of compound **1**.




Figure S2. The ^1H NMR spectrum of compound **1**.




Figure S3. The ^1H - ^1H COSY spectrum of compound **1**.




Figure S4. The HMBC spectrum of compound **1**.




Figure S5. The HSQC spectrum of compound **1**.




Figure S6. The DEPT135 spectrum of compound **1**.




Figure S7. The ^{13}C NMR spectrum of compound 2.




Figure S8. The ^1H NMR spectrum of compound 2.




Figure S9. The ¹H-¹H COSY spectrum of compound **2**.




Figure S10. The HMBC spectrum of compound **2**.




Figure S11. The HSQC spectrum of compound **2**.




Figure S12. The DEPT135 spectrum of compound **2**.




Figure S13. The ^{13}C NMR spectrum of compound 3.




Figure S14. The ^1H NMR spectrum of compound 3.