

1 Supplemental data

2 **Dissipation Kinetics and the Pre-Harvest Residue**
 3 **Limits of Acetamiprid and Chlorantraniliprole in**
 4 **Kimchi Cabbage using Ultra-Performance Liquid**
 5 **Chromatography-Tandem Mass Spectrometry**

6 Jonghwa Lee ^{1,2,†}, Byung Joon Kim ^{1,†}, Eunhye Kim ¹ and Jeong-Han Kim ^{1,*}


7 ¹ Department of Agricultural Biotechnology and Research Institute of Agriculture and Life Sciences, Seoul
 8 National University, Seoul, 08826, Republic of Korea; payton20@snu.ac.kr (B. K.); cocopam777@snu.ac.kr (E.
 9 K.)

10 ² Department of Veterinary and Animal Sciences, University of Massachusetts, Amherst, MA 01003, United
 11 States; jhlee006@gmail.com (J. L.)

12 * These authors contributed equally to this paper as first author.

13 * Correspondence: kjh2404@snu.ac.kr; Tel.: +82+2-880-4644


14 Received: date; Accepted: date; Published: date



15

16 **Figure S1.** Average weight of the kimchi cabbages which were measured in each sampling day
 17 ($n=3$).

18



19
20
21

Figure S2. Temperature and humidity of the field (Icheon) during the experimental period