

1

2

Supplementary material

3

Comparison of the partition efficiency of multiple phenolic compounds contained in propolis in different modes of acetonitrile-water based homogenous liquid-liquid extraction

7

8 **Wenbin Chen** ^{1,2,*}, **Xijuan Tu** ^{1,2}, **Dehui Wu** ¹, **Zhaosheng Gao** ¹, **Siyuan Wu** ¹, and **Shaokang**
9 **Huang** ¹


10 ¹ College of Bee Science, Fujian Agriculture and Forestry University, Fuzhou, China.

11 ² MOE Engineering Research Center of Bee Products Processing and Application, Fujian Agriculture and
12 Forestry University, Fuzhou, China.

13 * Correspondence: wbchen@fafu.edu.cn; Tel.: +86-591-83789482

14


15



16

17 **Figure S1.** Representative HPLC-DAD chromatogram ($\lambda=280$ nm) of phenolic standards. 1, caffeic
18 acid; 2, *p*-coumaric acid; 3, isoferulic acid; 4, dimethoxycinnamic acid; 5, cinnamic acid; 6, kaempferol;
19 7, caffeic acid phenethyl ester.


20



21

22 **Figure S2.** Extraction yields of investigated phenolic compounds under different addition amounts
 23 of a) NaCl and b) MgSO₄ in ACN-H₂O mixture (50:50, v/v). Error bars present the standard deviation
 24 (n=3).

25



26

27 **Figure S3.** Extraction yields of investigated phenolic compounds under different addition amounts
 28 of glucose in ACN-H₂O mixture (50:50, v/v). Error bars present the standard deviation (n=3).