

Supplementary Materials for

“Effect of spin contamination error and importance of open-shell structure for aggregation of Au atoms on a MgO surface”

Kohei Tada ^{1,*}, Tomohiro Maruyama ², Hiroaki Koga ³,
Mitsutaka Okumura ^{2,3}, and Shingo Tanaka ¹

¹ Department of Energy and Environment, National Institute of Advanced Industrial Science and Technology (AIST), 1-8-31, Midorigaoka, Ikeda, Osaka, 563-8577, Japan

² Department of Chemistry, Graduate School of Science, Osaka University, 1-1, Machikaneyama, Toyonaka, Osaka, 560-0043, Japan

³ Elements Strategy Initiative for Catalysts and Batteries (ESICB), Kyoto University, 1-30 Goryo Ohara, Nishikyo, Kyoto, Kyoto 615-8245, Japan

* Corresponding Author

phone: +81-72-751-8566

fax: +81-72-751-9714

k-tada@aist.go.jp

Contents

• Table S1	p. 3
• Figure S1	p. 4


Captions for Table S1 and Figure S1

Table S1 The nearest neighbor atoms of the ^2Au atom for all r .

Figure S1. Spin density distributions of the HS states. Isosurface values are 0.004 e/Bohr^3 .

Table S1 The nearest neighbor atoms of the ^2Au atom for all r.

r	LS	HS	BS
0.000	^1Au	^1Au	^1Au
0.025	^1Au	^1Au	^1Au
0.050	^1Au	^1Au	^1Au
0.075	^1Au	^1Au	^1Au
0.100	^1Au	^1Au	^1Au
0.125	^1Au	^1Au	^1Au
0.150	^1Au	^1Au	^1Au
0.175	^1Mg	^1Mg	^1Mg
0.200	^2O	^2O	^2O
0.225	^2O	^2O	^2O
0.250	^2O	^2O	^2O
0.275	^2O	^2O	^2O
0.300	^2O	^2O	^2O
0.325	^2Mg	^2Mg	^2Mg
0.350	^2Mg	^2Mg	^2Mg
0.375	^2Mg	^2Mg	^2Mg
0.400	^2Mg	^2Mg	^2Mg
0.425	^2Mg	^2Mg	^2Mg
0.450	^3O	^3O	^3O
0.475	^3O	^3O	^3O
0.500	^3O	^3O	^3O






Figure S1. Spin density distributions of the HS states. Isosurface values are 0.004 e/Bohr³.