

Article

Inhibitory Effect of Osthole from *Cnidium monnieri* on Tobacco Mosaic Virus (TMV) Infection in *Nicotiana glutinosa*

Ya-Han Chen ^{1,2}, Dong-Sheng Guo ², Mei-Huan Lu ^{2,3}, Jian-Ying Yue ¹, Yan Liu ⁴, Chun-Ming Shang ⁴, De-Rong An ^{2,*} and Ming-Min Zhao ^{1,*}

¹ College of Horticulture and Plant Protection, Inner Mongolia Agricultural University, Hohhot 010019, China; yhchen1018@nwfau.edu.cn (Y.-H.C.); yuejianying2018@163.com (J.-Y.Y.)

² College of Plant Protection and State Key Laboratory of Crop Stress Biology for Arid Areas, Northwest A&F University, Yangling 712100, China; gds1995908@163.com (D.-S.G.); lu_meihuan@sina.com (M.-H.L.)

³ Microbial Resources of Research Center, Microbiology Institute of Shaanxi, Xi'an 710043, China

⁴ Academy of Agriculture science in Baotou, Baotou 014010, China; liuyanww@126.com (Y.L.); chunmingsh@163.com (C.-M.S.)

* Correspondence: anderong323@163.com (D.-R.A.); Mingminzh@163.com (M.-M.Z.); Tel.: +86-158-2909-7529 (D.-R.A.); +86-157-7136-0659 (M.-M.Z.); Fax: +86-029-8708-2710 (D.-R.A.); +86-0471-638-5801 (M.-M.Z.)

Supplementary Materials




Figure S1. ¹H-NMR of osthole.




Figure S2. C-NMR of osthole.