

Lignin Precipitation and Fractionation from OrganoCat Pulping to Obtain Lignin with Different Sizes and Chemical Composition

Dennis Weidener ^{1,2,3}, **Arne Holtz** ^{3,4}, **Holger Klose** ^{1,3,5}, **Andreas Jupke** ⁴, **Walter Leitner** ^{2,6}
and **Philipp M. Grande** ^{1,3,*}

- ¹ Institute of Bio- and Geosciences, Plant Sciences, Forschungszentrum Jülich GmbH, 52425 Jülich, Germany
 - ² Institute of Technical and Macromolecular Chemistry (ITMC), RWTH Aachen University, Worringer Weg 1, 52074 Aachen, Germany
 - ³ Bioeconomy Science Center (BioSC), c/o Forschungszentrum Jülich, 52425 Jülich, Germany
 - ⁴ Fluid Process Engineering (AVT.FVT), RWTH Aachen University, Forckenbeckstraße 51, 52074 Aachen, Germany
 - ⁵ Institute of Biology I, RWTH Aachen University, Worringer Weg 3, 52074 Aachen, Germany
 - ⁶ Max-Planck-Institute of Chemical Energy Conversion, Stiftstraße 34-36, 45470 Mülheim an der Ruhr, Germany
- * Correspondence: p.grande@fz-juelich.de; Tel.: +49-2461-616341; Fax: +49-2461-612492




Figure S1: Expansion of HSQC NMR spectrum of OrganoCat lignin, A1 (carbohydrate region), A2 (lignin region) and A3 (aliphatic region), B investigated linkages and monomer units.