

Supplementary Materials: Confinement effects on glass-forming aqueous dimethyl sulfoxide solutions

Dominik Demuth, Melanie Reuhl, Moritz Hopfenmüller, Nail Karabas, Simon Schoner, and Michael Vogel

- 1 Figure S1 shows temperature-dependent dielectric loss spectra $\epsilon''(\nu)$ of D₂O-DMSO-*h*₆ in
- 2 lysozyme for two compositions of the solution, explicitly, for water mole fractions of $c_w = 66\%$ and
- 3 $c_w = 33\%$. Figure S2 displays temperature-dependent dielectric loss spectra $\epsilon''(\nu)$ of D₂O-DMSO-*h*₆
- 4 solution in ficoll at mass fractions of $w_s = 70\%$ and $w_s = 40\%$.




Figure S1. Dielectric loss spectra $\epsilon''(\nu)$ of D₂O-DMSO-*h*₆ in lysozyme for two compositions of the solution in the temperature range 140–300 K: (a) D₂O:DMSO ratio of 2:1 and (b) D₂O:DMSO ratio of 1:2.




Figure S2. Dielectric loss spectra $\epsilon''(\nu)$ of 66 mol% D₂O-DMSO-*h*₆ solution in ficoll at mass fractions of (a) $w_s = 70\%$ and (b) $w_s = 40\%$. The covered temperature ranges are indicated.