

Supplementary Materials:

Neo-aplysiatoxin A; a new aplysiatoxin derivative from Okinawan cyanobacterium *Moorea producens*

Mioko Kawaguchi, Masayuki Satake, Bo-Tao Zhang, Yue-Yun

Xiao, Masayuki Fukuoka, Hajime Uchida, Hiroshi Nagai

Figure S1. ESI-HRMS spectrum of neo-aplysiatoxin A in negative ion mode

Figure S2. ^1H NMR spectrum of neo-aplysiatoxin A in methanol- d_4

Figure S3. ^{13}C NMR spectrum of neo-aplysiatoxin A in methanol- d_4

Figure S4. ^1H - ^1H COSY NMR spectrum of neo-aplysiatoxin A in methanol- d_4

Figure S5. ^1H - ^{13}C HSQC spectrum of neo-aplysiatoxin A in methanol- d_4

Figure S6. ^1H - ^{13}C HMBC spectrum of neo-aplysiatoxin A in methanol- d_4

Figure S7. NOESY spectrum of neo-aplysiatoxin A in methanol- d_4




Figure S1. ESI-HRMS spectrum of neo-aplysiatoxin A in negative ion mode




Figure S2. ¹H NMR spectrum of neo-aplysiatoxin A in methanol-*d*₄




Figure S3. ^{13}C NMR spectrum of neo-aplysiatoxin A in methanol- d_4




Figure S4. ^1H - ^1H COSY NMR spectrum of neo-aplysiatoxin A in methanol- d_4




Figure S5. ^1H - ^{13}C HSQC spectrum of neo-aplysiatoxin A in methanol- d_4




Figure S6. ^1H - ^{13}C HMBC spectrum of neo-aplysiatoxin A in methanol- d_4




Figure S7. NOESY spectrum of neo-aplysiatoxin A in methanol- d_4