

Supplementary Materials

Lamalbid, chlorogenic acid, and verbascoside as the tools for standardization of *Lamium album* flowers-development and validation of HPLC-DAD method

Monika E. Czerwińska ^{1,*}, Eleonora Kalinowska², Dominik Popowski ¹, and Agnieszka Bazyłko ¹

¹ Department of Pharmacognosy and Molecular Basis of Phytotherapy, Medical University of Warsaw, 1 Banach street, 02-097 Warsaw, Poland; mczerwinska@wum.edu.pl (M.E.C);

dominik.popowski@wum.edu.pl (D.P.); agnieszka.bazylko@wum.edu.pl (A.B.)

² Students' Scientific Association at the Department of Pharmacognosy and Molecular Basis of Phytotherapy, Medical University of Warsaw, 1 Banach street, 02-097 Warsaw, Poland; eleonora_kalinowska@op.pl

* Correspondence: mczerwinska@wum.edu.pl; Tel+48-22-572-0953



Lamalbid (1)

Figure S1. HPLC-DAD chromatogram registered for standard of lamalbid (1).

Scheme S1. Analysis of residual sum of squares for lamalbid (1).

Chlorogenic acid (3)





Figure S2. HPLC-DAD chromatogram registered for standard of chlorogenic acid (3).

Scheme S2. Analysis of residual sum of squares for chlorogenic acid (3).





Rutin (4)

Figure S3. HPLC-DAD chromatogram registered for standard of rutin (4).

Scheme S3. Analysis of residual sum of squares for rutin (4).



Verbascoside (6)

Figure S4. HPLC-DAD chromatogram registered for standard of verbascoside (6).

Scheme S4. Analysis of residual sum of square for verbascoside (6).

Quercetin malonylhexoside (7)





Figure S5. HPLC-DAD chromatogram registered for standard of quercetin malonylhexoside (7).

Scheme S5. Analysis of residual sum of squares for quercetin malonylhexoside (7).




Tiliroside (8)

Figure S6. HPLC-DAD chromatogram registered for standard of tiliroside (8).

Scheme S6. Analysis of residual sum of squares for tiliroside (8).