

## Supplemental Information

# Ursolic Acid Enhances the Sensitivity of MCF-7 and MDA-MB-231 Cells to Epirubicin by Modulating the Autophagy Pathway


Zhennan Wang <sup>1,2,†</sup>, Pingping Zhang <sup>2,†</sup>, Huan Jiang <sup>1</sup>, Bing Sun <sup>1</sup>, Huaizhi Luo <sup>1</sup> and Aiqun Jia <sup>1,2,\*</sup>

<sup>1</sup> School of Environmental and Biological Engineering, Nanjing University of Science and Technology, Nanjing 210094, China; 15050525636@163.com (Z.W.); huanjiang929@foxmail.com (H.J.); yangwenleg@163.com (B.S.); rohizer@163.com (H.L.)


<sup>2</sup> State Key Laboratory of Marine Resource Utilization in South China Sea, School of Pharmaceutical Sciences, Hainan University, Haikou 570228, China; bbsuju2@163.com

\* Correspondence: ajia@hainanu.edu.cn; Tel.: +86-898-66254967; Fax: +86-898-66254967

† These authors contributed equally to this work.



**Figure S1.** ELISA analysis of the caspase3 activity in MCF-7 (A) and MAD-MB-231 (B) cells. \*  $P < 0.05$ , \*\*  $P < 0.01$ , \*\*\*  $P < 0.005$ .



**Figure S2.** LC3-II protein assembly process.