

Supplementary Materials for CYP2C9 manuscript

Development and Validation of an HPLC-UV Method for the Quantification of 4'-Hydroxydiclofenac Using Salicylic Acid: Future Applications for Measurement of In Vitro Drug–Drug Interaction in Rat Liver Microsomes

Hassan Salhab * and James Barker

School of Life Sciences, Pharmacy and Chemistry, Kingston University, Kingston upon Thames, London KT1 2EE, UK; j.barker@kingston.ac.uk

* Correspondence: k1542808@kingston.ac.uk; Tel.: +447984974741




Figure S1. Calibration curve of 4'-hydroxydiclofenac for intra-assay and inter-assay Day 1.




Figure S2. Calibration curve of 4'-hydroxydiclofenac for inter-assay Day 2.




Figure S3. Calibration curve of 4'-hydroxydiclofenac for inter-assay Day 3.




Figure S4. HPLC chromatogram of salicylic acid (100 μ M).




Figure S5. HPLC chromatogram of 4'-hydroxydiclofenac (100 μM).




Figure S6. HPLC chromatogram of Diclofenac (200 μM).




Figure S7. Calibration curve of 4'-hydroxydiclofenac stability test day 1.




Figure S8. Calibration curve of 4'-hydroxydiclofenac stability test day 2.




Figure S9. Calibration curve of 4'-hydroxydiclofenac stability test day 3.