

## Supplementary Materials

# Effect of Kaempferol and Its Glycoside Derivatives on Antioxidant Status of HL-60 Cells Treated with Etoposide

**Magdalena Kluska<sup>1</sup>, Michał Juszczak<sup>1</sup>, Jerzy Żuchowski<sup>2</sup>, Anna Stochmal<sup>2</sup> and Katarzyna Woźniak<sup>1\*</sup>**

<sup>1</sup> Department of Molecular Genetics, Faculty of Biology and Environmental Protection, University of Łódź, 90-236 Łódź, Poland

<sup>2</sup> Department of Biochemistry and Crop Quality, Institute of Soil Science and Plant Cultivation, State Research Institute, 24-100 Puławy, Poland

\* Correspondence: katarzyna.wozniak@biol.uni.lodz.pl; Tel.: +48-42-635-47-76; Fax: +48-42-635-44-84

**Table S1.** Gene expression changes in HL-60 cells treated with etoposide and kaempferol or its glycoside derivatives.

Treatment/Concentration	Relative gene expression (fold change)				
	<i>NFE2L2</i>	<i>HO-1</i>	<i>NQO1</i>	<i>SOD1</i>	<i>SOD2</i>
Ctrl	1 ± 0.16	1 ± 0.11	1 ± 0.16	1 ± 0.16	1 ± 0.27
K 10 µg/mL	0.77 ± 0.25	1.39 ± 0.18 **	1.6 ± 50.16 **	1.22 ± 0.1	1.2 ± 0.11
K 50 µg/mL	0.98 ± 0.15	2.52 ± 0.42 ***	2.3 ± 0.27 ***	1.42 ± 0.23 *	1.54 ± 0.25 *
P 2 10 µg/mL	0.85 ± 0.07	0.79 ± 0.17	1.18 ± 0.52	1.5 ± 0.28 *	1.08 ± 0.33
P 2 50 µg/mL	0.9 ± 0.27	1.09 ± 0.14	1.32 ± 0.13 *	1.18 ± 0.13	1.25 ± 0.09
P 5 10 µg/mL	0.91 ± 0.13	0.99 ± 0.12	1.24 ± 0.32	1.99 ± 0.24 ***	1.57 ± 0.35 *
P 5 50 µg/mL	0.96 ± 0.07	1.1 ± 0.29	1.3 ± 0.15 *	1.14 ± 0.15	1.1 ± 0.14
P 7 10 µg/mL	0.92 ± 0.14	1 ± 0.24	1.27 ± 0.3	1.3 ± 0.33	1.12 ± 0.23
P 7 50 µg/mL	1.08 ± 0.42	1.06 ± 0.25	1.75 ± 0.2 **	1.72 ± 0.14 ***	1.65 ± 0.23 *
E 1 µM	1.13 ± 0.27	4.87 ± 0.18 ***	1.14 ± 0.26	1.01 ± 0.1	1.01 ± 0.09
E 1 µM + K 10 µg/mL	0.98 ± 0.28	9.49 ± 0.51 *** ###	1.51 ± 0.51	1.68 ± 0.52 * #	1.72 ± 0.53 * #
E 1 µM + K 50 µg/mL	1.06 ± 0.15	9.33 ± 0.35 *** ###	1.84 ± 0.38 ** #	1.41 ± 0.36	1.72 ± 0.37 * ##
E 1 µM + P 2 10 µg/mL	1.08 ± 0.44	4.83 ± 0.28 ***	1.36 ± 0.37	1.21 ± 0.09 #	0.83 ± 0.55
E 1 µM + P 2 50 µg/mL	1.28 ± 0.17	6.62 ± 0.28 *** ###	1.21 ± 0.37	1.46 ± 0.09 ** ###	1.89 ± 0.55 * #
E 1 µM + P 5 10 µg/mL	1.07 ± 0.16	5.45 ± 0.1 *** ###	1.28 ± 0.24	1.11 ± 0.08	0.9 ± 0.32
E 1 µM + P 5 50 µg/mL	1.09 ± 0.11	9.97 ± 0.42 *** ###	1.69 ± 0.41 *	1.34 ± 0.35	1.59 ± 0.4 #
E 1 µM + P 7 10 µg/mL	1.22 ± 0.16	5.29 ± 0.17 *** #	1.3 ± 0.13 *	1.43 ± 0.18 * ##	1.18 ± 0.15
E 1 µM + P 7 50 µg/mL	1.19 ± 0.23	5.49 ± 0.22 *** ##	1.55 ± 0.12 ** #	1.28 ± 0.17 #	1.38 ± 0.13 * ##

*NFE2L2* - nuclear factor erythroid 2-related factor 2, *HO-1* - heme oxygenase-1, *NQO1* - NAD(P)H:quinone oxidoreductase 1, *SOD1* - superoxide dismutase 1, *SOD2* - superoxide dismutase 2. Relative expression of genes in HL-60 cells incubated for 24 h at 37 °C with 10-50 µg/mL kaempferol (K), P2, P5, P7 and/or 1 µM etoposide (E). The table shows mean results ± SD, n = 4; \* p < 0.05, \*\* p < 0.01, \*\*\* p < 0.001 vs. control (Ctrl); # p < 0.05, ## p < 0.01, ### p < 0.001 vs. etoposide (E).