

Synthesis and anticancer activity of 1,3,4-thiadiazole with 3-methoxyphenyl substituent.

Sara Janowska¹, Dmytro Khylyuk¹, and Monika Wujec^{1,*}

¹Department of Organic Chemistry, Faculty of Pharmacy, Medical University,
4a Chodzki Str., 20-093 Lublin, Poland;

Table of contents:

1. ¹ H NMR spectra of thiosemicarbazide derivatives (SC1-SC6).....	2
2. ¹ H NMR spectra of 1,3,4-thiadiazole derivatives (STC1-STC6).....	8
3. ¹³ C NMR spectra of thiosemicarbazide derivatives (SC1-SC6).....	13
4. ¹³ C NMR spectra of 1,3,4-thiadiazole derivatives (STC1-STC6).....	19




Figure S1. The ^1H NMR of compound SC1.




Figure S2. The ^1H NMR of compound SC2.




Figure S3. The ^1H NMR of compound SC3.




Figure S4. The ^1H NMR of compound SC4.




Figure S5. The ^1H NMR of compound SC5.




Figure S6. The ^1H NMR of compound SC6.




Figure S7. The ^1H NMR of compound SCT1.




Figure S8. The ^1H NMR of compound SCT2.




Figure S9. The ¹H NMR of compound SCT4.




Figure S10. The ^1H NMR of compound SCT5.




Figure S11. The ^1H NMR of compound SCT6.




Figure S12. The ^{13}C NMR of compound SC1.




Figure S13. The ^{13}C NMR of compound SC2.




Figure S14. The ^{13}C NMR of compound SC3.




Figure S15. The ^{13}C NMR of compound SC4.




Figure S16. The ^{13}C NMR of compound SC5.




Figure S17. The ^{13}C NMR of compound SC6.




Figure S18. The ^{13}C NMR of compound SCT1.




Figure S19. The ^{13}C NMR of compound SCT2.




Figure S20. The ^{13}C NMR of compound SCT4.




Figure S21. The ^{13}C NMR of compound SCT5.




Figure S22. The ^{13}C NMR of compound SCT6.

