

Supplementary Materials

Stimuli-Responsive Polypeptide Nanoparticles for Enhanced DNA delivery

Olga Korovkina ¹, Dmitry Polyakov ², Viktor Korzhikov-Vlakh ^{1,3} and Evgenia Korzhikova-Vlakh ^{3*}

¹ Saint-Petersburg State University, Institute of Chemistry, Universitetsky pr. 26, 198504 St. Petersburg, Russia

² Institute of Experimental Medicine, Acad. Pavlov Street 12, 197376 St. Petersburg, Russia

³ Institute of Macromolecular Compounds, Russian Academy of Sciences, Bolshoy pr. 31, 199004, St. Petersburg, Russia

* Correspondence: vlakh@hq.macro.ru (E.K.-V.)



(a)



(b)

Figure S1. The initial gel-electrophoresis images to the marked ones shown in Figure 6 of the main text: (a) and (b) correspond to (a) and (b) of Figure 6. The description is provided in the Figure 6 and its legend, as well as in the text of section 2.4.




Figure S2. The initial gel-electrophoresis images to the marked ones shown in Figure 8 of the main text: top and bottom images **(a)** correspond to (a) and (b) images of Figure 8, respectively; top and bottom images **(b)** correspond to (c) and (d) images of Figure 8, respectively. The description is provided in the Figure 8 and its legend, as well as in the text of section 2.5.