

Supplementary Information

Synthesis and antimicrobial activity of new Mannich bases with piperazine moiety

Sara Janowska^{1*}, Sylwia Andrzejczuk², Piotr Gawryś³ and Monika Wujec^{1*}

¹Department of Organic Chemistry, Faculty of Pharmacy, Medical University of Lublin, 4a Chodzki Street, 20-093 Lublin, Poland

²Department of Pharmaceutical Microbiology, Faculty of Pharmacy, Medical University of Lublin, 1 Chodzki Street, 20-093 Lublin, Poland

³Students Research Group at the Department of Organic Chemistry

*corresponding authors: monika.wujec@umlub.pl, sara.janowska@umlub.pl

Table of contents:

1. ¹ H NMR spectra	2
2. ¹³ C NMR spectra	6




Figure S1. The ^1H NMR of compound 5.




Figure S2. The ^1H NMR of compound 6.




Figure S3. The ^1H NMR of compound 7.




Figure S4. The ^1H NMR of compound 8.




Figure S5. The ^1H NMR of compound 9.




Figure S6. The ^{13}C NMR of compound 5.




Figure S7. The ^{13}C NMR of compound 6.




Figure S8. The ^{13}C NMR of compound 7.




Figure S9. The ^{13}C NMR of compound 8.




Figure S10. The ^{13}C NMR of compound 9.