

Talaromyces sp. Ethyl Acetate Crude Extract as Potential Mosquitocide to Control *Culex pipiens quinquefasciatus*

Junhui Chen ^{1,†}, Zhiyong Xu ^{2,†}, Yangqing Liu ³, Feiying Yang ¹, Limei Guan ¹, Jian Yang ¹, Jianghuai Li ¹, Guodong Niu ^{1,4}, Jun Li ^{1,4} and Liang Jin ^{1,*}

¹ Institute of Biological Resources, Jiangxi Academy of Sciences, Nanchang 330929, China; allenchen0426@gmail.com (J.C.); 18059141865@163.com (F.Y.); glmnh@126.com (L.G.); jemappelleyangjian@zju.edu.cn (J.Y.); jeremy_leakey@sina.com (J.L.); gniu@fiu.edu (G.N.); lij@fiu.edu (J.L.)

² Institute of Applied Chemistry, Jiangxi Academy of Sciences, Nanchang 330929, China; zhiyongxuconfident@hotmail.com

³ Nanchang Center for Disease Control and Prevention, Nanchang 330100, China; nccdclq@163.com

⁴ Department of Biological Sciences, Florida International University, Miami, FL 33199, USA

* Correspondence: jinliang079@163.com; Tel.: +86-183-7917-4620

† These authors contributed equally to this work.

Figure S1. ¹H NMR (400 MHz, CDCl₃) spectrum of vermistatin -----2

Figure S2. ¹³C NMR (100 MHz, CDCl₃) spectrum of vermistatin -----3

Figure S3. ¹H NMR (400 MHz, CDCl₃) spectrum of dihydrovermistatin -----3

Figure S4. ¹³C NMR (100 MHz, CDCl₃) spectrum of dihydrovermistatin -----4




Figure S1. ¹H NMR (400 MHz, CDCl₃) spectrum of J1




Figure S2. ¹³C NMR (100 MHz, CDCl₃) spectrum of J1




Figure S3. ¹H NMR (400 MHz, CDCl₃) spectrum of J2




Figure S4. ^{13}C NMR (100 MHz, CDCl_3) spectrum of J2