

Article

Polyoxypregnane Glycosides from Root of *Marsdenia tenacissima* and Inhibited Nitric Oxide Levels in LPS Stimulated RAW 264.7 Cells

Zhi Na ^{1,*}, Pianchou Gongpan ¹ and Qingfei Fan ^{2,*} 

¹ Key Laboratory of Tropical Plant Resources and Sustainable Use, Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences, Mengla 666303, China

² College of Science, Yunnan Agricultural University, Kunming 650201, China

* Correspondence: nazhi@xtbg.org.cn (Z.N.); fanqingfei@xtbg.ac.cn (Q.F.)

Abstract: Six new polyoxypregnane glycosides, marstenacisside F1–F3 (1–3), G1–G2 (4–5) and H1 (6), as well as 3-O-β-D-glucopyranosyl-(1→4)-6-deoxy-3-O-methyl-β-D-allopyranosyl-(1→4)-β-D-oleandropyranosyl-11α,12β-di-O-benzoyl-tenacigenin B (7), were isolated from the roots of *Marsdenia tenacissima*. Their structures were established by an extensive interpretation of their 1D and 2D NMR and HRESIMS data. Compounds 1–7 were tenacigenin B derivatives with an oligosaccharide chain at C-3. This was the first time that compound 7 had been isolated from the title plant and its ¹H and ¹³C NMR data were reported. Compounds 4 and 5 were the first examples of C₂₁ steroid glycoside bearing unique β-glucopyranosyl-(1→4)-β-glucopyranose sugar moiety. All the isolated compounds were evaluated for anti-inflammatory activity by inhibiting nitric oxide (NO) production in the lipopolysaccharide-induced RAW 264.7 cells. The results showed that marstenacisside F1 and F2 exhibited significant NO inhibitory activity with an inhibition rate of 48.19 ± 4.14% and 70.33 ± 5.39%, respectively, at 40 μM, approximately equal to the positive control (L-NMMA, 68.03 ± 0.72%).

Keywords: *Marsdenia tenacissima*; *Asclepiadaceae*; polyoxypregnane glycoside; anti-inflammatory



Citation: Na, Z.; Gongpan, P.; Fan, Q.

Polyoxypregnane Glycosides from Root of *Marsdenia tenacissima* and Inhibited Nitric Oxide Levels in LPS Stimulated RAW 264.7 Cells.

Molecules **2023**, *28*, 886. <https://doi.org/10.3390/molecules28020886>

Academic Editor: Jih-Jung Chen

Received: 7 December 2022

Revised: 9 January 2023

Accepted: 12 January 2023

Published: 16 January 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Marsdenia tenacissima (Roxb.) Moon (*Asclepiadaceae*), a perennial climber, is distributed mainly in the southwest of China and other parts of tropical and subtropical Asia. The stems and roots of *M. tenacissima* are traditional Chinese medicine and Dai herbal medicine, respectively. The dried stems of *M. tenacissima*, known as “tongguanteng”, have been used in the treatment of asthma, cancer, and trachitis [1]. The roots of this plant, known as “Dai-Bai-Jie”, have been widely used as a Dai herbal medicine by Dai people living in Laos, Myanmar and the Yunnan province of China due to the root’s pharmacological functions of relieving pain, clearing heat, decreasing swelling and detoxification, etc. [2]. There have been many more chemical investigations on the stems than the roots of *M. tenacissima*. Previous phytochemical studies on the stems had revealed this plant as an extremely rich source of C₂₁ steroid glycosides [3–14]. Although “Dai-Bai-Jie” has been widely used as a Dai herbal medicine, few phytochemical studies on the roots of this plant have been reported so far [15–17]. These studies showed that the main chemical composition of roots was also the same as the stems, i.e., C₂₁ steroid glycosides. These compounds were only screened for anti-HIV activity and were necessary for anti-inflammatory activity, because the traditional usage for “Dai-Bai-Jie” was the treatment of inflammatory-associated diseases. Inflammation is a response of the organism to injury related to physical or chemical noxious stimuli or microbiological toxins, which is involved in multiple pathologies such as arthritis, asthma, multiple sclerosis, colitis, inflammatory bowel diseases, and atherosclerosis [18]. It can be speculated that the presence of key chemical constituents with effective

anti-inflammatory activity had led to the extensive clinical application of “Dai-Bai-Jie” in traditional ethnomedicine, so we were interested in clarifying the relationship between the constituents and anti-inflammatory activity of this plant.

In order to search for more novel natural products, particularly those with potential anti-inflammatory activity, from the roots of *M. tenacissima*, a systematic phytochemical study was carried out on their 95% ethanol extract. As a result, six new polyoxypregnane glycosides, named marstenacisside F1–F3 (1–3), G1–G2 (4–5), and H1 (6), as well as 3-*O*- β -D-glucopyranosyl-(1 \rightarrow 4)-6-deoxy-3-*O*-methyl- β -D-allopyranosyl-(1 \rightarrow 4)- β -D-oleandropyranosyl-11 α ,12 β -di-*O*-benzoyl-tenacigenin B (7), were isolated (Figure 1). Compound 7 was isolated from the title plant, and its ^1H and ^{13}C NMR data were reported, for the first time. In the present paper, we describe the isolation and structure elucidation of these compounds, and we also evaluate the anti-inflammatory activity of the isolated compounds in terms of the inhibitory effect on NO production in LPS-induced RAW 264.7 cells.




Figure 1. Structures of pregnane glycosides (1–7) isolated from *M. tenacissima*.

2. Results and Discussion

2.1. Structural Elucidation

Compound 1 displayed a sodium adduct ion at m/z 877.4344 $[\text{M} + \text{Na}]^+$ in its HRESIMS spectrum (Figure S1), and its molecular formula was determined as $\text{C}_{47}\text{H}_{66}\text{O}_{14}$ (calcd for $\text{C}_{47}\text{H}_{66}\text{NaO}_{14}$, 877.4345). There were three methyl signals at δ_{H} 1.18 (3H, s, CH_3 -18), 1.09 (3H, s, CH_3 -19), and 2.26 (3H, s, CH_3 -21), and three methine protons bearing secondary alcoholic functions at δ_{H} 3.60 (m, H-3), 5.58 (t, $J = 10.2$ Hz, H-11) and 5.25 (d, $J = 10.2$ Hz, H-12) in the ^1H NMR spectrum of 1. The combination of the ^1H and ^{13}C NMR data (Figures S2 and S3) indicated a C_{21} steroidal skeleton for 1. By comparison with C_{21} steroids isolated from the title plant, the ^{13}C NMR data (Table 1) of 1 were similar to those of 11,12-diester of tenacigenin B [19]. The signals at δ_{H} 6.55 (q, $J = 6.8$ Hz), 1.46 (m), and 1.44 (s), and δ_{C} 167.4, 138.1, 128.5, 14.2, and 11.6, indicated the existence of a tigloyl (Tig) group. The long-range correlation from δ_{H} 5.58 (H-11) to δ_{C} 167.4 (Tig-C-1) in the HMBC spectrum (Figure 2) was

observed, which indicated the Tig group was assigned at C-11. At the same time, there was a benzoyl (Bz) group in **1**, due to the existence of a series of NMR signals at δ_{H} 7.89 (d, $J = 7.4$ Hz), 7.51 (t, $J = 7.4$ Hz), and 7.38 (t, $J = 7.4$ Hz), and δ_{C} 166.1, 133.1, 129.7 (2C), 129.4 and 128.3 (2C). The Bz group was demonstrated to be attached at C-12 by the HMBC correlation of H-12 (δ_{H} 5.25) to the carbonyl carbon (δ_{C} 166.1) of the benzoyl group. In the NOESY spectrum of **1** (Figure 3), crossing peaks between H-11 and CH₃-18 (δ_{H} 1.18) and between H-12 and H-9 (δ_{H} 2.08) revealed that H-11 and H-12 were in β -orientation and α -orientation, respectively. Furthermore, the C-17 side chain was in an α -orientation as supported by the NOESY correlations between H-17 (δ_{H} 2.95) and CH₃-18 (δ_{H} 1.18), and between H-12 (δ_{H} 5.25) and CH₃-21 (δ_{H} 2.26) (Figure S7). Thus, the aglycone of **1** was determined to be 11 α -O-tigloyl-12 β -O-benzoyl-tenacigenin B.

Table 1. ¹³C NMR data of the aglycones of compounds **1–7** in pyridine-*d*₅ (125 MHz, δ in ppm).

Position	1 ^a	2 ^a	3 ^a	4	5	6	7
1	37.4	37.6	38.5	37.8	38.1	37.9	37.8
2	29.1	29.1	28.9	29.8	29.6	29.8	29.7
3	76.4	76.4	76.6	77.8	77.9	76.0	75.9
4	34.7	34.8	34.5	35.2	35.1	35.2	35.2
5	44.0	44.0	44.5	44.0	43.9	43.9	43.9
6	26.7	26.7	27.0	27.2	27.1	27.2	27.3
7	31.8	31.9	32.3	25.2	25.2	25.1	25.3
8	66.9	66.9	66.0	66.8	66.8	66.7	66.9
9	51.3	51.3	54.3	51.9	51.8	51.7	51.8
10	39.1	39.2	39.3	39.4	39.4	39.4	39.5
11	68.8	69.5	68.6	69.1	68.8	69.0	69.9
12	75.2	75.5	74.2	75.2	75.2	74.9	75.5
13	46.1	46.2	47.3	44.0	46.1	46.1	46.5
14	71.5	71.6	71.6	71.7	71.6	71.7	71.9
15	26.8	26.8	27.7	32.2	32.1	32.1	32.1
16	25.1	25.1	25.4	27.1	27.1	27.1	27.2
17	59.8	59.9	60.3	59.7	60.0	59.9	59.9
18	16.6	16.7	17.5	16.9	17.0	16.9	16.9
19	12.7	12.8	12.9	13.2	13.2	13.1	13.2
20	211.1	211.1	212.6	210.3	210.0	210.1	210.4
21	30.3	30.2	32.7	30.3	29.9	30.0	30.2
11-O	Tig	Bz	HPA	Tig	mBu	HPA	Bz
1	167.4	166.2	172.0	167.2	175.5	171.5	166.5
2	128.5	130.0	41.1	129.2	41.5	41.4	130.6
3	138.1	129.5	125.3	138.1	26.6	124.6	129.9
4	11.6	128.1	130.7	12.2	11.9	131.3	128.7
5	14.2	132.9	115.6	14.3	15.7	116.4	133.5
6		128.1	155.2			158.2	128.7
7		129.5	115.6			116.4	129.9
8			130.7			131.3	
12-O	Bz	Bz		Ac	Ac	Ac	Bz
1	166.1	166.1		170.7	170.8	170.8	166.3
2	129.4	129.0		20.5	20.9	20.3	130.6
3	129.7	129.5					129.8
4	128.3	128.1					128.7
5	133.1	132.8					133.3
6	128.3	128.1					128.7
7	129.7	129.5					129.8

^a Measured in CDCl₃.




Figure 2. Key HMBC and ^1H - ^1H COSY correlations of **1**.




Figure 3. Key ROESY correlations of the aglycone of **1**.

In the NMR spectra of **1**, there were two anomeric proton signals at δ_{H} 4.58 (dd, $J = 9.8, 1.8$ Hz) and 4.79 (d, $J = 8.1$ Hz) and two carbon signals at δ_{C} 96.9 and 99.1. The above evidence proved that the sugar moiety of **1** contained two units. The coupling constants (8.1 and 9.8 Hz) of the two anomeric protons indicated that both glycosidic linkages were β -oriented. At the same time, there were characteristic proton signals of two methyls at δ_{H} 1.36 (d, $J = 5.5$ Hz) and 1.25 (d, $J = 6.0$ Hz); two methoxyl groups at δ_{H} 3.66 (s) and 3.37 (s) in the ^1H NMR spectrum of **1**. The evidence of two methoxyl groups located at C-3 in each of the two sugar moiety was deduced from the long-range correlations between the methoxyl group at δ_{H} 3.37 (s) and the carbon signal at δ_{C} 78.8, and between another methoxyl group at δ_{H} 3.66 (s) and the carbon signal at δ_{C} 81.0. Subsequently, the two sugar units were identified as 6-deoxy-3-methoxy sugars, which generally occur in *M. tenacissima* [1]. Based on the HSQC, HMBC, and ^1H - ^1H COSY spectra (Figure S4–S6), the NMR spectra of each sugar were fully assigned (Tables 2 and 3). The two sugar units were then determined as oleandropyranosyl (Ole) and 6-deoxy-3-*O*-methyl-allopyranose (Allo), respectively [20]. Meanwhile, the oleandrose was the inner sugar and 6-deoxy-3-*O*-methyl allose was the outer, which was supported by the HMBC correlations of δ_{H} 4.79 (Allo-H-1) with δ_{C} 79.1 (Ole-C-4), and δ_{H} 4.58 (Ole-H-1) with δ_{C} 76.4 (aglycone-C-3). Consequently, the sugar moiety of **1** was determined as pachybiose. Compared to the previously reported data of tenacigenin B [21], changes in the chemical shift in aglycone of **1**, i.e., C-2 (−3.2 ppm), C-3 (+5.6 ppm), and C-4 (−3.8 ppm) were observed, which suggested that the sugar moiety was linked to the C-3 hydroxyl of the aglycone. Thus, compound **1** was elucidated as 3-*O*-6-deoxy-3-*O*-methyl- β -D-allopyranosyl (1→4)- β -D-oleandropyranosyl-11 α -*O*-tigloyl-12 β -*O*-benzoyl-tenacigenin B and named marstenacisside F1.

Table 2. ^{13}C NMR data of the sugar moieties of compounds 1–7 in pyridine- d_5 (125 MHz, δ in ppm).

Position	1 ^a	2 ^a	3 ^a	4	5	6	7
	Ole	Ole	Ole	Glc-1	Glc-1	Ole	Ole
1	96.9	97.0	97.0	101.2	101.1	97.5	97.4
2	36.1	36.1	36.1	71.4	71.5	37.9	37.8
3	78.8	78.8	78.8	78.0	78.0	79.7	79.6
4	79.1	79.1	79.1	84.7	84.7	83.4	83.3
5	71.4	71.4	71.4	78.0	77.9	71.9	71.9
6	18.6	18.6	18.6	62.6	62.6	19.1	19.0
3-OMe	55.6	55.6	55.7			57.3	57.3
	Allo	Allo	Allo	Glc-2	Glc-2	Allo	Allo
1	99.1	99.1	99.2	106.7	106.7	102.0	101.9
2	71.8	71.8	71.8	77.1	77.1	72.7	72.7
3	81.0	81.0	81.0	78.3	78.3	83.2	83.2
4	72.8	72.8	72.8	71.4	71.5	83.4	83.3
5	71.3	71.3	71.3	78.9	78.9	69.6	69.5
6	17.9	17.9	17.9	62.6	62.6	18.4	18.3
3-OMe	62.0	61.9	62.0			61.7	61.7
						Glc	Glc
1						106.6	106.6
2						75.5	75.5
3						78.4	78.4
4						72.0	72.0
5						78.5	78.4
6						63.0	63.0

^a Measured in CDCl_3 .**Table 3.** ^1H NMR data of the sugar moieties of compounds 1–7 in pyridine- d_5 (500 MHz, δ in ppm, J in Hz).

Position	1 ^a	2 ^a	3 ^a	4	5	6	7
	Ole	Ole	Ole	Glc-1	Glc-1	Ole	Ole
1	4.58 dd (9.8, 1.8)	4.58 dd (9.8, 1.9)	4.58 dd (9.7, 1.7)	5.01 d (7.6)	5.08 d (7.7)	4.78 d (8.9)	4.73 d (9.4)
2	1.47 m	1.45 m	1.49 m	4.31 m	4.31 m	1.26 m	1.33 m
	2.30 m	2.30 m	2.30 m			2.41 m	2.36 m
3	3.39 m	3.38 m	3.38 m	4.25 m	4.23 m	3.61 m	3.57 m
4	3.33 m	3.31 m	3.34 m	4.09 m	4.13 m	3.59 m	3.52 m
5	3.33 m	3.31 m	3.34 m	3.92 m	3.92 m	3.64 m	3.52 m
6	1.36 d (5.5)	1.32 d (5.5)	1.37 d (5.5)	4.33 m 4.50 m	4.34 m 4.52 m	1.67 d (5.8)	1.57 d (5.0)
3-OMe	3.37 s	3.35 s	3.37 s			3.50 s	3.47 s
	Allo	Allo	Allo	Glc-2	Glc-2	Allo	Allo
1	4.79 d (8.1)	4.77 d (8.3)	4.79 d (8.3)	5.22 d (7.8)	5.25 d (7.8)	5.27 d (8.1)	5.24 d (8.1)
2	3.47 m	3.46 m	3.47 m	4.11 m	4.12 m	3.84 m	3.79 m
3	3.79 t (3.0)	3.78 t (3.0)	3.78 t (3.0)	4.35 m	4.37 m	4.47 t (2.5)	4.45 m
4	3.17 m	3.17 m	3.17 m	4.21 m	4.21 m	3.73 dd (9.6, 2.5)	3.72 dd (9.4, 2.0)
5	3.55 m	3.54 m	3.55 m	3.93 m	3.94 m	4.27 m	4.23 m
6	1.25 d (6.0)	1.24 d (6.4)	1.25 d (6.1)	4.33 m 4.45 m	4.34 m 4.43 m	1.63 d (6.2)	1.61 d (6.2)
3-OMe	3.66 s	3.65 s	3.65 s			3.81 s	3.80 s
						Glc	Glc
1						4.97 d (7.8)	4.95 d (7.7)
2						4.02 m	4.00 m
3						4.26 m	4.22 m
4						4.22 m	4.19 m
5						4.01 m	3.97 m
6						4.36 dd (11.6, 5.4)	4.36 dd (11.6, 5.1)
						4.54 dd (11.6, 2.3)	4.53 d (11.6)

^a Measured in CDCl_3 .

The ^1H and ^{13}C NMR spectroscopic data ascribed to the sugar moieties of 2–3 are consistent with those of 1 (Tables 2 and 3), so they should contain the same sugar moiety as 1.

Compound 2 possessed a molecular formula of $\text{C}_{49}\text{H}_{64}\text{O}_{14}$, determined by HRESIMS ion at m/z 899.4190 $[\text{M} + \text{Na}]^+$ (calcd for $\text{C}_{49}\text{H}_{64}\text{NaO}_{14}$, 899.4188) (Figure S8). The NMR data of 2 showed a pattern analogous to 1, except for an ester group (Figures S9 and S10). In the ^{13}C NMR spectrum of 2, there were signals of two benzoyl groups on the aglycone of 2.

Meanwhile, the tigloyl unit signals of **1** were absent in **2**. An extra Bz group was positioned at C-11, which was deduced from the HMBC correlation from δ_{H} 5.77 (H-11) to δ_{C} 166.2 (Bz₁-C-1) (Figure S13). Due to the HMBC correlation between δ_{H} 4.58 (Ole-H-1) and δ_{C} 76.4 (aglycone-C-3), the glycosidation site was located at C-3 of the aglycone. Consequently, **2** was defined as 3-O-6-deoxy-3-O-methyl- β -D-allopyranosyl(1 \rightarrow 4)- β -D-oleandropyranosyl-11 α ,12 β -di-O-benzoyl-tenacigenin B, and named marstenacisside F2 (Figures S11–S14).

Compound **3** showed a quasi-molecular ion peak at m/z 825.4040 [$M + \text{Na}$]⁺, in accordance with the molecular formula C₄₃H₆₂O₁₄ (calcd for C₄₃H₆₂NaO₁₄, 825.4032) (Figure S15). In the ¹H NMR spectrum of **3** (Figure S16), due to the absence of an ester group at C-12, there was a higher-field shift signal at δ_{H} 3.27 (1H, d, $J = 9.7$ Hz), compared to the δ_{H} 5.35 (1H, d, $J = 10.2$ Hz) in the ¹H NMR spectrum of **2**. Consequently, the aglycone of **3** was a monoester of tenacigenin B. The signals at δ_{H} 7.09 and 6.75 (d, $J = 8.6$ Hz, each 2H) and the ¹³C NMR (Figure S17) singlet at δ_{C} 155.2 indicated that **3** contained a 4-hydroxyphenyl group. In the HMBC spectrum (Figure S20), the aromatic resonances at δ_{C} 125.3 and 130.7 were correlated with the signal at δ_{H} 3.54 (m, 2H), and 7.09 correlated with the δ_{C} 41.1. Moreover, the HMBC spectrum showed a correlation between δ_{H} 3.54 and δ_{C} 172.0. Therefore, a methylene group (δ_{H} 3.54, δ_{C} 41.1) was located between the 4-hydroxyphenyl moiety and the carbonyl group (δ_{C} 172.0). Hence, **3** contained a (4-hydroxyphenyl) acetyl (HPA) group. The HPA group was assigned at C-11 by HMBC correlation of the proton at δ_{H} 5.05 (H-11) with the carbonyl carbon at δ_{C} 172.0 of the HPA group. The glycosidation site was deduced from the HMBC cross peaks between δ_{H} 4.58 (Ole-H-1) and δ_{C} 76.6 (aglycone-C-3). Therefore, **3** was defined as 3-O-6-deoxy-3-O-methyl- β -D-allopyranosyl (1 \rightarrow 4)- β -D-oleandropyranosyl-11 α -O-(4-hydroxyphenyl) acetyl-tenacigenin B and named marstenacisside F3 (Figures S18–S21).

Compound **4** had a molecular formula of C₄₀H₆₀O₁₇, as determined by HRESIMS [$M + \text{Na}$]⁺ ion at m/z 835.3714 (calcd for C₄₀H₆₀NaO₁₇, 835.3723) (Figure S22). The proton signals of δ_{H} 1.90 (3H, s), 6.86 (1H, qq, $J = 7.1, 1.4$ Hz), 1.80 (3H, s), and 1.61 (3H, d, $J = 7.1$ Hz) and the carbon signals of δ_{C} 170.7, 20.5, 167.2, 129.2, 138.1, 12.2, and 14.3 indicated the presence of an acetyl (Ac) and a tigloyl group in the aglycone of **4**. The ¹H NMR data of the aglycone moiety of **4** were very close to those of **1**, except for the presence of the signal for an acetyl group at δ_{H} 1.90 (3H, s) and the absence of the protons signals for the benzoyl group. HMBC cross-peaks between δ_{H} 5.76 (H-11) and δ_{C} 167.2 (Tig-C-1), and between δ_{H} 5.40 (H-12) and δ_{C} 170.7 (Ac-C-1) (Figure S27), indicated that a tigloyl and acetyl group were located at C-11 and C-12, respectively. Accordingly, the aglycone of **4** was identified as 11 α -O-tigloyl-12 β -O-acetyl-tenacigenin B.

In the NMR spectra of **4** (Figures S23 and S24), there were two anomeric proton signals at δ_{H} 5.01 (d, $J = 7.6$ Hz) and 5.22 (d, $J = 7.8$ Hz) and two carbon signals at δ_{C} 101.2 and 106.7. The above facts proved that the sugar moiety of **4** contained two units. Furthermore, the ¹³C NMR spectra displayed two terminal oxygenated methylene groups at δ_{C} 62.6. On the basis of the above evidence and compared with previously reported data [22], the two sugar units were identified as glucopyranoses. According to the coupling constants of anomeric protons (7.6 and 7.8 Hz), the linkages of two sugar units were in β -configuration. The linkage of two sugar moiety could be β -glucopyranosyl-(1 \rightarrow 4)- β -glucopyranoside deduced from the long-range correlations between δ_{H} 5.22 (Glc₂-H-1) and δ_{C} 84.7 (Glc₁-C-4). Relative to the previously reported values of 11 α -O-tigloyl-12 β -O-acetyl-tenacigenin B [23], changes in the chemical shift in aglycone of **4**, i.e., C-2 (−1.8 ppm), C-3 (+8.4 ppm), and C-4 (−2.9 ppm) were detected, which suggested that the sugar moiety was attached at the C-3 of the aglycone. Thus, compound **4** was finally elucidated as 3-O- β -D-glucopyranosyl-(1 \rightarrow 4)- β -D-glucopyranosyl-11 α -O-tigloyl-12 β -O-acetyl-tenacigenin B and named marstenacisside G1 (Figures S25–S28).

The ¹H and ¹³C NMR data of the sugar moiety of **5** were well in agreement with those of **4**. Accordingly, **5** had the same sugar moiety as **4**.

Compound **5** exhibited a molecular formula of C₄₀H₆₂O₁₇ based on the HRESIMS ion [$M + \text{Na}$]⁺ at m/z 837.3883 (calcd for C₄₀H₆₂NaO₁₇, 837.3879) (Figure S29). The NMR data of **5** (Figures S30 and S31) showed a similar pattern to **4**, except for an ester substitution.

In the ^1H NMR spectrum of **5**, there were proton signals of an acetyl group at δ_{H} 2.02 (3H, s) and extra signals of 2-methylbutyryl ester units at δ_{H} 0.84 (3H, t, $J = 7.5$ Hz), 1.04 (3H, d, $J = 7.0$ Hz), 1.34 (1H, m), 1.68 (1H, m), and 2.26 (1H, m). Meanwhile, the signals of the tigloyl unit of **4** were absent in **5**. The HMBC cross peak between the carbonyl carbon at δ_{C} 175.5 of the 2-methylbutyryl group and the proton signal at δ_{H} 5.68 (H-11) disclosed a 2-methylbutyryl group located at C-11 (Figure S34). Likewise, the HMBC correlation of the carbonyl carbon at δ_{C} 170.8 of the acetyl unit with the proton signal at δ_{H} 5.36 (H-12) revealed the existence of an acetyl group at C-12. The glycosidation site was deduced from the long-range coupling of the δ_{H} 5.08 (Glc₁-H-1) with δ_{C} 77.9 (agclone-C-3). Accordingly, compound **5** was established as 3-*O*- β -D-glucopyranosyl-(1 \rightarrow 4)- β -D-glucopyranosyl-11 α -*O*-2-methylbutyryl-12 β -*O*-acetyl-tenacigenin B and was named marstenacisside G2 (Figures S32–S35).

Compound **6** gave a molecular formula of $\text{C}_{51}\text{H}_{74}\text{O}_{20}$ based on the HRESIMS (m/z 1029.4659 $[\text{M} + \text{Na}]^+$, calcd for $\text{C}_{51}\text{H}_{74}\text{NaO}_{20}$, 1029.4666) (Figure S36). ^{13}C NMR data analysis indicated that the aglycone moiety of **6** differs from the aglycone moiety of **3** by the presence of an extra acetyl group (δ_{C} 170.8 and 20.3). The positions of the diester groups were deduced by the HMBC correlations between δ_{H} 5.68 (H-11) and δ_{C} 171.5 (HPA-C-1), and between δ_{H} 5.36 (H-12) and δ_{C} 170.8 (Ac-C-1) (Figure S41). Therefore, the aglycone structure of **6** was identified as 11 α -*O*-(4-hydroxyphenyl) acetyl-12 β -*O*-acetyl-tenacigenin B.

The ^1H and ^{13}C NMR spectra of **6** (Figures S37 and S38) exhibited three anomeric proton signals at δ_{H} 5.27 (1H, d, $J = 8.1$ Hz), 4.97 (1H, d, $J = 7.8$ Hz), and 4.78 (1H, d, $J = 8.9$ Hz) and three anomeric carbon signals at δ_{C} 102.0, 106.6 and 97.5, suggesting the existence of three sugar units in the molecule. Due to the large coupling constants of anomeric protons, the linkages of three sugar units were in β -configuration. At the same time, there were characteristic proton signals of two methyls at δ_{H} 1.67 (d, $J = 5.8$ Hz) and 1.63 (d, $J = 6.2$ Hz); two methoxyl groups at δ_{H} 3.81 (s) and 3.50 (s); and two ABM spin protons at δ_{H} 4.36 (dd, $J = 11.6, 5.4$ Hz), 4.54 (dd, $J = 11.6, 2.3$ Hz) in the ^1H NMR spectrum of **6**. On the basis of the above evidence and compared with previously reported data [24,25], the sugar units were identified as oleandrose, 6-deoxy-3-*O*-methyl-allose, and glucoses. The connectivity of the sugars was established by the HMBC correlations between δ_{H} 4.97 (Glc-H-1) and δ_{C} 83.4 (Allo-C-4); between δ_{H} 5.27 (Allo-H-1) and δ_{C} 83.4 (Ole-C-4). As a result, the sugar moiety was determined as β -D-glucopyranosyl-(1 \rightarrow 4)-6-deoxy-3-*O*-methyl- β -D-allopyranosyl-(1 \rightarrow 4)- β -D-oleandropyranoside, which was identical to the neocondurangotriose in the compounds that were also isolated from *M. tenacissima* [17]. Furthermore, the glycosidation site was deduced from the long-range correlations between δ_{H} 4.78 (Ole-H-1) and δ_{C} 76.0 (aglycone-C-3). Hence, the structure of **6** was established as 3-*O*- β -D-glucopyranosyl-(1 \rightarrow 4)-6-deoxy-3-*O*-methyl- β -D-allopyranosyl-(1 \rightarrow 4)- β -D-oleandropyranosyl-11 α -*O*-(4-hydroxyphenyl) acetyl-12 β -*O*-acetyl-tenacigenin B and named marstenacisside H1 (Figures S39–S42).

Compound **7** had a molecular formula of $\text{C}_{55}\text{H}_{74}\text{O}_{19}$ determined by HRESIMS ion $[\text{M} + \text{Na}]^+$ at m/z 1061.4718 (calcd for $\text{C}_{55}\text{H}_{74}\text{NaO}_{19}$, 1061.4722) (Figure S43). Compound **7** was predicted to be novel pregnane glycoside in a crude extract of *Marsdenia tenacissima* by means of LC-ESI-MSⁿ [26], and the compound was not isolated from the crude extract. This is the first time that compound **7** has been isolated from the title plant and its ^1H and ^{13}C NMR data were reported. The structure of **7** was elucidated as 3-*O*- β -D-glucopyranosyl-(1 \rightarrow 4)-6-deoxy-3-*O*-methyl- β -D-allopyranosyl-(1 \rightarrow 4)- β -D-oleandropyranosyl-11 α ,12 β -di-*O*-benzoyl-tenacigenin B, on the basis of 1D, 2D NMR and HRESI data (Figures S44–S49).

2.2. NO Inhibitory Evaluations

Compounds **1–7** were screened for anti-inflammatory activity by inhibiting NO production in LPS-induced RAW 264.7 cells. Compounds **1** and **2** showed significant NO inhibitory activity with an inhibition rate of $48.19 \pm 4.14\%$ and $70.33 \pm 5.39\%$, respectively, at 40 μM , approximately equal to the positive control (L-NMMA, $68.03 \pm 0.72\%$) (Figure 4 and Table 4). The effects of compounds **1–7** on cell viability are shown in Figure S50. All

compounds showed dose-dependent NO inhibitory activity. Only **1** and **2** showed significant NO inhibitory activity, and other compounds did not show the activity. The above facts may be related to the structure of compounds **1** and **2**. There were tigloyl and/or benzoyl groups at C-11 and C-12 of **1** and **2**, and the sugar moiety was pachybiose. Although **7** also had two benzoyl groups at C-11 and C-12, the sugar moiety was neocondurangotriose.




Figure 4. Inhibitory effects of compounds **1**–**7** on NO production in LPS-induced RAW 264.7 cells. Experiments were performed in triplicate, and the data are presented as the mean \pm SD. ** $p < 0.01$, *** $p < 0.001$ vs. LPS group. NS, not significant by *T*-test.

Table 4. Inhibitory effects of compounds **1**–**7** on NO production in LPS-induced RAW 264.7 cells.

Compound	Concentration (μ M)	NO Inhibition Rate (%)
1	40	48.19 \pm 4.14
2	40	70.33 \pm 5.39
3	40	−4.09 \pm 7.28
4	40	−0.86 \pm 1.59
5	40	0.80 \pm 1.91
6	40	−5.57 \pm 1.15
7	40	7.13 \pm 5.00
L-NMMA ^a	40	68.03 \pm 0.72

^a Positive control.

3. Materials and Methods

3.1. General Experimental Procedures

The UV spectra were collected on a Shimadzu UV-2401 PC spectrophotometer (Shimadzu Corp: Kyoto, Japan). Optical rotations were measured on an Autopol VI polarimeter; The IR spectra were determined on a Nicolet iS10 spectrometer with KBr pellets. HRESIMS was recorded on an Agilent 1290 UPLC/6540 Q-TOF mass spectrometer (Agilent, Palo Alto, CA, USA). All NMR spectra were acquired on a Bruker Avance III 500 spectrometer. Semi-preparative HPLC was performed on a Waters HPLC system consisting of a 1525 binary pump and a 2487 detector, equipped with a YMC-pack ODS-A column (250 \times 10 mm, YMC Co., Ltd., Kyoto, Japan). Silica gel (200–300 mesh, Qingdao Marine Chemical Co., Ltd., Qingdao, China), Lichroprep RP-18 gel (40–63 μ m, Merck, Darmstadt, Germany), Sephadex LH-20 gel (GE Healthcare, Sweden), and MCI gel (75–150 μ m, Mitsubishi Chemical Co., Tokyo, Japan) were used to perform column chromatography.

3.2. Plant Material

The roots of *Marsdenia tenacissima* were collected from Simao, Yunnan Province, China in January 2020. A voucher specimen (No. 20200101) was deposited in the authors' research group.

3.3. Extraction and Isolation

The dried powder roots of *Marsdenia tenacissima* (2.5 kg) were percolated with 95% ethanol at room temperature three times (3 days each time) and then concentrated under reduced pressure to give concentrated extract. The concentrated extract was efficiently partitioned with ethyl acetate (EtOAc). The EtOAc fraction (68.6 g) was separated by an MCI gel CHP 20P column, eluted with MeOH–H₂O (*v/v*, 30:70, 50:50, 80:20, 95:5) to provide four portions (Fr. A–D). Fr. B (28.1 g) was subjected to silica gel CC eluted with CH₂Cl₂–MeOH (25:1–3:1) to obtain five fractions (Fr. B.1–5). Fr. B.2 (2.8 g) was chromatographed over a Sephadex LH-20 column, eluting with MeOH to give four fractions (Fr. B.2.1–4). Fr. B.2.2 (256 mg) was further purified by semi-preparative HPLC using MeOH/H₂O (70:30, 3 mL/min) to give compounds **3** (12 mg, *t_R* = 12.3 min) and **6** (9 mg, *t_R* = 19.6 min). Fr. B.4 (2.1 g) was separated by ODS MPLC (MeOH–H₂O, 60:40 to 100:0) to yield five fractions (Fr. B.4.1–5). Fr. B.4.3 (202 mg) was further separated by semi-preparative with MeOH/H₂O (80:20, 3 mL/min) to yield compounds **4** (11 mg, *t_R* = 8.3 min) and **5** (8 mg, *t_R* = 9.6 min). Fr. C (35.7 g) was subjected to silica gel CC eluted with CH₂Cl₂–MeOH (50:1–4:1) to obtain five fractions (Fr. C.1–5). Fr. C.3 (3.2 g) was chromatographed over a Sephadex LH-20 column, eluting with CH₂Cl₂–MeOH (1:1) to give four fractions (Fr. C.3.1–4). Fr. C.3.3 (306 mg) was further purified by semi-preparative HPLC using MeOH/H₂O (75:25, 3 mL/min) to afford compounds **7** (11 mg, *t_R* = 12.3 min), **1** (9 mg, *t_R* = 16.6 min), and **2** (10 mg, *t_R* = 20.7 min).

3.4. Compound Characterization Data

Marstenacisside F1 (**1**): white amorphous powder; $[\alpha]_D^{23} + 19.6$ (c 0.15, MeOH); UV (MeOH) λ_{\max} (log ϵ): 196 (4.46), 226 (4.26), 273 (3.01) nm; IR (KBr): ν_{\max} 3436, 2929, 1719, 1451, 1367, 1281, 1164, 1071, 988, 711 cm⁻¹; ¹H NMR (CDCl₃) data of aglycone moiety of **1**: δ 1.09 (3H, s, 19-CH₃), 1.18 (3H, s, 18-CH₃), 1.44 (3H, brs, Tig-H-5), 1.46 (3H, m, Tig-H-4), 2.08 (1H, m, H-9), 2.26 (3H, s, 21-CH₃), 2.99 (1H, d, *J* = 6.4 Hz, H-17 β), 3.60 (1H, m, H-3), 5.25 (1H, d, *J* = 10.2 Hz, H-12 α), 5.58 (1H, t, *J* = 10.2 Hz, H-11 β), 6.55 (1H, q, *J* = 6.8 Hz, Tig-H-3), 7.38 (2H, t, *J* = 7.4 Hz, Bz-H-4, 6), 7.51 (1H, t, *J* = 7.4 Hz, Bz-H-5), 7.89 (2H, d, *J* = 7.4 Hz, Bz-H-3, 7); HRESIMS: *m/z* 877.4344 [M + Na]⁺ (calcd for C₄₇H₆₆NaO₁₄, 877.4345); for ¹³C NMR data of the aglycone moiety of **1** see Table 1. For ¹H and ¹³C NMR data of the sugar moiety of **1** see Tables 2 and 3.

Marstenacisside F2 (**2**): white amorphous powder; $[\alpha]_D^{23} + 26.9$ (c 0.18, MeOH); UV (MeOH) λ_{\max} (log ϵ): 196 (4.49), 230 (4.33), 274 (3.34) nm; IR (KBr): ν_{\max} 3436, 2932, 1721, 1451, 1367, 1283, 1162, 1070, 988, 708 cm⁻¹; ¹H NMR (CDCl₃) data of aglycone moiety of **2**: δ 1.14 (3H, s, 19-CH₃), 1.21 (3H, s, 18-CH₃), 2.21 (1H, m, H-9), 2.27 (3H, s, 21-CH₃), 3.01 (1H, d, *J* = 7.3 Hz, H-17 β), 3.60 (1H, m, H-3), 5.35 (1H, d, *J* = 10.2 Hz, H-12 α), 5.77 (1H, t, *J* = 10.2 Hz, H-11 β), 7.18 (2H, t, *J* = 7.4 Hz, Bz₂-H-4, 6), 7.22 (2H, t, *J* = 7.4 Hz, Bz₁-H-4, 6), 7.33 (1H, t, *J* = 7.4 Hz, Bz₂-H-5), 7.37 (1H, t, *J* = 7.4 Hz, Bz₁-H-5), 7.75 (4H, d, *J* = 7.4 Hz, Bz₁-H-3, 7, Bz₂-3, 7); HRESIMS: *m/z* 899.4190 [M + Na]⁺ (calcd for C₄₉H₆₄NaO₁₄, 899.4188); for ¹³C NMR data of the aglycone moiety of **2** see Table 1. For ¹H and ¹³C NMR data of the sugar moiety of **2** see Tables 2 and 3.

Marstenacisside F3 (**3**): white amorphous powder; $[\alpha]_D^{19} - 4.9$ (c 0.10, MeOH); UV (MeOH) λ_{\max} (log ϵ): 196 (4.16), 224 (3.69), 278 (3.08) nm; IR (KBr): ν_{\max} 3445, 2933, 1703, 1619, 1367, 1164, 1068, 989, 610 cm⁻¹; ¹H NMR (CDCl₃) data of aglycone moiety of **3**: δ 1.03 (3H, s, 19-CH₃), 1.14 (3H, s, 18-CH₃), 1.85 (1H, m, H-9), 2.26 (3H, s, 21-CH₃), 3.01 (1H, t, *J* = 6.2 Hz, H-17 β), 3.27 (1H, d, *J* = 9.7 Hz, H-12 α), 3.54 (2H, m, HPA-2), 3.66 (1H, m, H-3), 5.05 (1H, d, *J* = 9.7 Hz, H-11 β), 6.75 (2H, d, *J* = 8.6 Hz, HPA-H-5, 7), 7.09 (2H, d, *J* = 8.6 Hz, HPA-H-4, 8); HRESIMS: *m/z* 825.4040 [M + Na]⁺ (calcd for C₄₃H₆₂NaO₁₄, 825.4032); for ¹³C NMR data of the aglycone moiety of **3** see Table 1. For ¹H and ¹³C NMR data of the sugar moiety of **3** see Tables 2 and 3.

Marstenacisside G1 (**4**): white amorphous powder; $[\alpha]_D^{23} + 2.1$ (c 0.10, MeOH); UV (MeOH) λ_{\max} (log ϵ): 196 (3.93), 216 (3.94) nm; IR (KBr): ν_{\max} 3428, 2934, 1706, 1619, 1368, 1269, 1077, 1031 cm⁻¹; ¹H NMR (pyridine-*d*₅) data of aglycone moiety of **4**: δ 1.20 (3H, s, 19-CH₃), 1.28 (3H, s, 18-CH₃), 1.61 (3H, d, *J* = 7.1 Hz, Tig-H-5), 1.80 (3H, brs, Tig-H-4), 2.05

(1H, m, H-9), 1.90 (3H, s, Ac-H-2), 2.23 (3H, s, 21-CH₃), 2.90 (1H, d, $J = 6.7$ Hz, H-17 β), 3.88 (1H, m, H-3), 5.40 (1H, d, $J = 10.2$ Hz, H-12 α), 5.76 (1H, t, $J = 10.2$ Hz, H-11 β), 6.86 (1H, qq, $J = 7.1, 1.4$ Hz, Tig-H-3); HRESIMS: m/z 835.3714 [M + Na]⁺ (calcd for C₄₀H₆₀NaO₁₇, 835.3723); for ¹³C NMR data of the aglycone moiety of 4 see Table 1. For ¹H and ¹³C NMR data of the sugar moiety of 4 see Tables 2 and 3.

Marstenacisside G2 (5): $[\alpha]_D^{22} + 13.3$ (c 0.10, MeOH); UV (MeOH) λ_{max} (log ϵ): 196 (3.83), 228 (3.48), 274 (2.72) nm; IR (KBr): ν_{max} 3426, 2936, 1736, 1632, 1367, 1247, 1076, 1030 cm⁻¹; ¹H NMR (pyridine-*d*₅) data of aglycone moiety of 5: δ 0.84 (3H, t, $J = 7.5$ Hz, mBu-H-4), 1.04 (3H, d, $J = 7.0$ Hz, mBu-H-5), 1.13 (3H, s, 19-CH₃), 1.25 (3H, s, 18-CH₃), 1.34 and 1.68 (2H, m, mBu-H-3), 2.02 (3H, s, Ac-H-2), 2.00 (1H, m, H-9), 2.26 (1H, m, mBu-H-2), 2.24 (3H, s, 21-CH₃), 2.87 (1H, d, $J = 6.9$ Hz, H-17 β), 3.84 (1H, m, H-3), 5.36 (1H, d, $J = 10.2$ Hz, H-12 α), 5.68 (1H, t, $J = 10.2$ Hz, H-11 β); HRESIMS: m/z 837.3883 [M + Na]⁺ (calcd for C₄₀H₆₂NaO₁₇, 837.3879); for ¹³C NMR data of the aglycone moiety of 5 see Table 1. For ¹H and ¹³C NMR data of the sugar moiety of 5 see Tables 2 and 3.

Marstenacisside H1 (6): white amorphous powder; $[\alpha]_D^{22} - 1.6$ (c 0.10, MeOH); UV (MeOH) λ_{max} (log ϵ): 197 (4.38), 225 (3.98), 277 (3.57) nm; IR (KBr): ν_{max} 3445, 2935, 1738, 1516, 1367, 1253, 1071, 991 cm⁻¹; ¹H NMR (pyridine-*d*₅) data of aglycone moiety of 6: δ 1.13 (3H, s, 18-CH₃), 1.17 (3H, s, 19-CH₃), 1.77 (3H, s, Ac-H-2), 2.07 (1H, d, $J = 10.1$ Hz, H-9), 2.24 (3H, s, 21-CH₃), 2.86 (1H, d, $J = 6.7$ Hz, H-17 β), 3.82 (1H, m, H-3), 5.36 (1H, d, $J = 10.2$ Hz, H-12 α), 5.68 (1H, t, $J = 10.2$ Hz, H-11 β), 7.11 (2H, d, $J = 8.4$ Hz, HPA-H-5, 7), 7.29 (2H, d, $J = 8.6$ Hz, HPA-H-4, 8); HRESIMS: m/z 1029.4659 [M + Na]⁺ (calcd for C₅₁H₇₄NaO₂₀, 1029.4666); for ¹³C NMR data of the aglycone moiety of 6 see Table 1. For ¹H and ¹³C NMR data of the sugar moiety of 6 see Tables 2 and 3.

3-*O*- β -D-glucopyranosyl-(1 \rightarrow 4)-6-deoxy-3-*O*-methyl- β -D-allopyranosyl-(1 \rightarrow 4)- β -D-oleandropyranosyl-11 α ,12 β -di-*O*-benzoyl-tenacigenin B (7): white amorphous powder; $[\alpha]_D^{21} + 20.8$ (c 0.23, MeOH); UV (MeOH) λ_{max} (log ϵ): 201 (4.18), 230 (4.24), 274 (3.16) nm; IR (KBr): ν_{max} 3427, 2934, 1721, 1452, 1367, 1281, 1071, 710 cm⁻¹; ¹H NMR (pyridine-*d*₅) data of aglycone moiety of 7: δ 1.29 (3H, s, 18-CH₃), 1.31 (3H, s, 19-CH₃), 2.09 (1H, m, H-9), 2.32 (3H, s, 21-CH₃), 2.97 (1H, d, $J = 6.3$ Hz, H-17 β), 3.78 (1H, m, H-3), 5.78 (1H, d, $J = 10.2$ Hz, H-12 α), 6.13 (1H, t, $J = 10.2$ Hz, H-11 β), 7.16 (2H, t, $J = 7.4$ Hz, Bz₂-H-4, 6), 7.22 (2H, t, $J = 7.4$ Hz, Bz₁-H-4, 6), 7.24 (1H, t, $J = 7.4$ Hz, Bz₂-H-5), 7.43 (1H, t, $J = 7.4$ Hz, Bz₁-H-5), 7.94 (2H, d, $J = 7.4$ Hz, Bz₂-H-3, 7), 7.99 (2H, d, $J = 7.4$ Hz, Bz₁-H-3, 7); HRESIMS: m/z 1061.4718 [M + Na]⁺ (calcd for C₅₅H₇₄NaO₁₉, 1061.4722); for ¹³C NMR data of the aglycone moiety of 7 see Table 1. For ¹H and ¹³C NMR data of the sugar moiety of 7 see Tables 2 and 3.

3.5. Cell Culture and Nitric Oxide Inhibitory Assay

The macrophage RAW 264.7 cells (passage number was 10–13) were obtained from Cell Bank of Chinese Academy of Sciences. The RAW 264.7 cells were plated in 96-well plates (1.5×10^5 cells/well) and treated with different isolate concentrations (dissolved in DMSO) of 10, 20, and 40 μ M, respectively, followed by stimulation with 1 μ g/mL LPS (Sigma, St. Louis, MO, USA) for 18 h [27]. Griess reagents (Sigma, St. Louis, MO, USA) were used to measure NO production. The optical density (OD) was determined at a 570 nm wavelength, with L-NMMA as a positive control [28]. Three independent experiments were carried out in triplicate. The cell viability was evaluated by the MTT assay [29].

4. Conclusions

Six new polyoxypregnane glycosides, marstenacisside F1–F3 (1–3), G1–G2 (4–5), and H1 (6), as well as 3-*O*- β -D-glucopyranosyl-(1 \rightarrow 4)-6-deoxy-3-*O*-methyl- β -D-allopyranosyl-(1 \rightarrow 4)- β -D-oleandropyranosyl-11 α ,12 β -di-*O*-benzoyl-tenacigenin B (7), were isolated from the ethanolic extract of the roots of *Marsdenia tenacissima* by modern chromatographic techniques and characterized by comprehensive spectroscopic data. Their structures were tenacigenin B derivatives with an oligosaccharide chain at C-3. Compounds 4 and 5 were the first examples of C₂₁ steroid glycoside bearing unique β -glucopyranosyl-(1 \rightarrow 4)- β -glucopyranose sugar moiety. Compound 7 was isolated from the title plant for the first

time, and its ^1H and ^{13}C NMR data were reported. The patterns of compounds 1–7 were consistent with those of compounds previously isolated from this plant. All isolates were evaluated for anti-inflammatory activity by inhibiting the production of NO stimulated by LPS in RAW 264.7 cells, with L-NMMA as a positive control. Among those compounds, compounds 1 and 2 exhibited significant NO inhibition at 40 μM .

Supplementary Materials: The following supporting information can be downloaded at <https://www.mdpi.com/article/10.3390/molecules28020886/s1>; Figures S1–S49: HRESIMS, 1D and 2D NMR of compounds 1–7. Figure S50: Effects of compounds 1–7 on cell viability.

Author Contributions: Z.N. designed the experiment, performed the isolation and identification of all the compounds, and also wrote this paper; P.G. contributed to the nitric oxide inhibition assay, data analysis, and wrote this section; Q.F. provided comments and suggestions on structure elucidation and reviewed the manuscript. All authors have read and agreed to the published version of the manuscript.

Funding: This work was financially supported by the Chinese Academy of Sciences (CAS) 135 Program (XTBG-F02) and Yunnan Fundamental Research Projects (grant NO. 202101AT070057).

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: All the data in this research were presented in the manuscript and Supplementary Material.

Conflicts of Interest: The authors declare no conflict of interest.

Sample Availability: Samples of compounds 1–7 are available from the authors.

References

1. Wang, P.L.; Yang, J.; Zhu, Z.F.; Zhang, X.J. *Marsdenia tenacissima*: A Review of Traditional Uses, Phytochemistry and Pharmacology. *Am. J. Chin. Med.* **2018**, *46*, 1449–1480. [[CrossRef](#)] [[PubMed](#)]
2. Li, H.T.; Kang, L.P.; Guo, B.L.; Zhang, Z.L.; Guan, Y.H.; Pang, X.; Peng, C.Z.; Ma, B.P.; Zhang, L.X. Original plant identification of Dai nationality herb “Daibaijie”. *Chin. J. Chin. Mater. Med.* **2014**, *39*, 1525–1529.
3. Qiu, S.X.; Luo, S.Q.; Lin, L.Z.; Cordell, G.A. Further polyoxypregnane glycosides from *Marsdenia tenacissima*. *Phytochemistry* **1996**, *41*, 1385–1388. [[PubMed](#)]
4. Xia, Z.H.; Xing, W.X.; Mao, S.L.; Lao, A.N.; Uzawa, J.; Yoshida, S.; Fujimoto, Y. Pregnane glycosides from the stems of *Marsdenia tenacissima*. *J. Asia Nat. Prod. Res.* **2004**, *6*, 79–85. [[CrossRef](#)] [[PubMed](#)]
5. Wang, X.L.; Li, Q.F.; Yu, K.B.; Peng, S.L.; Zhou, Y.; Ding, L.S. Four new pregnane glycosides from the stems of *Marsdenia tenacissima*. *Helv. Chim. Acta* **2006**, *89*, 2738–2744. [[CrossRef](#)]
6. Liu, J.; Yu, Z.B.; Ye, Y.H.; Zhou, Y.W. A new C₂₁ steroid glycoside from *Marsdenia tenacissima*. *Chin. Chem. Lett.* **2008**, *19*, 444–446. [[CrossRef](#)]
7. Lei, Y.S.; Li, Z.L.; Yang, S.S.; Liu, Z.L.; Hua, H.M. C₂₁ steroids from the stems of *Marsdenia tenacissima*. *Acta Pharm. Sin.* **2008**, *43*, 509–512.
8. Huang, X.D.; Liu, T.; Wang, S. Two new polyoxypregnane glycosides from *Marsdenia tenacissima*. *Helv. Chim. Acta* **2009**, *92*, 2111–2117. [[CrossRef](#)]
9. Wang, X.L.; Peng, S.L.; Ding, L.S. Further polyoxypregnane glycosides from *Marsdenia tenacissima*. *J. Asia Nat. Prod. Res.* **2010**, *12*, 654–661. [[CrossRef](#)]
10. Zhang, H.; Tan, A.M.; Zhang, A.Y.; Chen, R.; Yang, S.B.; Huang, X. Five new C₂₁ steroidal glycosides from the stems of *Marsdenia tenacissima*. *Steroids* **2010**, *75*, 176–183. [[CrossRef](#)]
11. Xia, Z.H.; Mao, S.L.; Lao, A.N.; Uzawa, J.; Yoshida, S.; Fujimoto, Y. Five new pregnane glycosides from the stems of *Marsdenia tenacissima*. *J. Asia Nat. Prod. Res.* **2011**, *13*, 477–485. [[CrossRef](#)] [[PubMed](#)]
12. Yao, S.; To, K.K.W.; Wang, Y.Z.; Yin, C.; Tang, C.P.; Chai, S.; Ke, C.Q.; Lin, G.; Ye, Y. Polyoxypregnane Steroids from the Stems of *Marsdenia tenacissima*. *J. Nat. Prod.* **2014**, *77*, 2044–2053. [[CrossRef](#)] [[PubMed](#)]
13. Khang, P.V.; Xu, T.; Xu, L.X.; Zhou, N.N.; Hu, L.H.; Wang, R.; Ma, L. Steroidal glycosides from *Marsdenia tenacissima*. *Phytochem. Lett.* **2015**, *12*, 54–58. [[CrossRef](#)]
14. Yao, S.; To, K.K.W.; Ma, L.; Yin, C.; Tang, C.P.; Chai, S.; Ke, C.Q.; Lin, G.; Ye, Y. Polyoxypregnane steroids with an open-chain sugar moiety from *Marsdenia tenacissima* and their chemoresistance reversal activity. *Phytochemistry* **2016**, *126*, 47–58. [[CrossRef](#)] [[PubMed](#)]

15. Pang, X.; Kang, L.P.; Yu, H.S.; Zhao, Y.; Han, L.F.; Zhang, J.; Xiong, C.Q.; Zhang, L.X.; Yu, L.Y.; Ma, B.P. New polyoxypregnane glycosides from the roots of *Marsdenia tenacissima*. *Steroids* **2015**, *93*, 68–76. [[CrossRef](#)] [[PubMed](#)]
16. Pang, X.; Kang, L.P.; Fang, X.M.; Yu, H.S.; Han, L.F.; Zhao, Y.; Zhang, L.X.; Yu, L.Y.; Ma, B.P. C₂₁ steroid derivatives from the Dai herbal medicine Dai-Bai-Jie, the dried roots of *Marsdenia tenacissima*, and their screening for anti-HIV activity. *J. Nat. Med.* **2017**, *72*, 166–180. [[CrossRef](#)] [[PubMed](#)]
17. Pang, X.; Kang, L.P.; Fang, X.M.; Zhao, Y.; Yu, H.S.; Han, L.F.; Li, H.T.; Zhang, L.X.; Guo, B.L.; Yu, L.Y.; et al. Polyoxypregnane glycosides from the roots of *Marsdenia tenacissima* and their anti-HIV activities. *Planta Med.* **2017**, *83*, 126–134. [[CrossRef](#)]
18. Guzik, T.J.; Korbout, R.; Adamek-Guzik, T. Nitric oxide and superoxide in inflammation. *J. Physiol. Pharmacol.* **2003**, *54*, 469–487.
19. Deng, J.; Liao, Z.X.; Chen, D.F. Three new polyoxypregnane glycosides from *Marsdenia tenacissima*. *Helv. Chim. Acta* **2005**, *88*, 2675–2682. [[CrossRef](#)]
20. Deng, J.; Liao, Z.X.; Chen, D.F. Marsdenosides A–H, polyoxypregnane glycosides from *Marsdenia tenacissima*. *Phytochemistry* **2005**, *66*, 1040–1051. [[CrossRef](#)]
21. Deng, J.; Liao, Z.X.; Chen, D.F. Two new C₂₁ steroids from *Marsdenia tenacissima*. *Chin. Chem. Lett.* **2005**, *16*, 487–490.
22. Li, Q.F.; Wang, X.L.; Ding, L.S.; Zhang, C. Polyoxypregnanes from the stems of *Marsdenia tenacissima*. *Chin. Chem. Lett.* **2007**, *18*, 831–834. [[CrossRef](#)]
23. Luo, S.Q.; Lin, L.Z.; Cordell, G.A.; Xue, L.; Johnson, M.E. Polyoxypregnane glycosides from *Marsdenia tenacissima*. *Phytochemistry* **1993**, *34*, 1615–1620.
24. Ma, B.X.; Fang, T.Z. Novel saponins hainaneosides A and B isolated from *Marsdenia hainanensis*. *J. Nat. Prod.* **1997**, *60*, 134–138. [[CrossRef](#)]
25. Sahu, N.P.; Panda, N.; Mandal, N.B.; Banerjee, S.; Koike, K.; Nikaid, O.T. Polyoxypregnane glycosides from the flowers of *Dregea volubilis*. *Phytochemistry* **2002**, *61*, 383–388. [[CrossRef](#)]
26. McGarvey, B.D.; Liao, H.; Ding, K.Y.; Wang, X.L. Dereplication of known pregnane glycosides and structural characterization of novel pregnanes in *Marsdenia tenacissima* by high-performance liquid chromatography and electrospray ionization-tandem mass spectrometry. *J. Mass. Spectrom.* **2012**, *47*, 687–693. [[CrossRef](#)]
27. Cheng, C.S.; Zou, Y.; Peng, J. Oregano Essential Oil Attenuates RAW264.7 Cells from Lipopolysaccharide-Induced Inflammatory Response through Regulating NADPH Oxidase Activation-Driven Oxidative Stress. *Molecules* **2018**, *23*, 1857. [[CrossRef](#)]
28. Reif, D.W.; McCreedy, S.A. N-nitro-L-arginine and N-monomethyl-L-arginine exhibit a different pattern of inactivation toward the three nitric oxide synthases. *Arch. Biochem. Biophys.* **1995**, *320*, 170–176. [[CrossRef](#)]
29. Jin, K.K.; Hyun, C.G. 4-Hydroxy-7-Methoxycoumarin inhibits inflammation in LPS-activated RAW264.7 macrophages by suppressing NF- κ B and MAPK activation. *Molecules* **2020**, *25*, 4424–4433.

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.