

The Synthesis, Structural Characterization, and DFT

Calculation of a New Binuclear Gd(III) Complex with 4-Acetylphenoxyacetic Acid and 1,10-Phenanthroline Ligands and Its Roles in Catalytic Activity

Ying Liu ¹, Xiao Tang ², Xi-Hai Yan ¹, Li-Hua Wang ³, Xi-Shi Tai ^{1,*}, Mohammad Azam ^{4,*} and Dong-Qiu Zhao ^{5,*}

¹ College of Chemistry and Chemical Engineering, Weifang University, Weifang 261061, China;
ydliuying@163.com (Y.L.); yan7899@126.com (X.-H.Y.)

² College of Science, Institute of Materials Physics and Chemistry, Nanjing Forestry University, Nanjing 210037, China; xiaotang@njfu.edu.cn

³ College of Biology and Oceanography, Weifang University, Weifang 261061, China; wanglihua929@163.com

⁴ Department of Chemistry, College of Science, King Saud University, Riyadh 11451, Saudi Arabia

⁵ School of Physics and Electric Engineering, Anyang Normal University, Anyang 455000, China

* Correspondence: taixs@wfu.edu.cn (X.-S.T.); azam_res@yahoo.com (M.A.); dqzhao@aynu.edu.cn (D.-Q.Z.); Tel.: +86-536-8785286 (X.-S.T.); Fax: +86-536-8785286 (X.-S.T.)




Figure s1. The calculated absorption spectrum of the ligands