

Microsolvation of a Proton by Ar Atoms: Structures and Energetics of Ar_nH^+ Clusters

María Judit Montes de Oca-Estévez ^{1,2,†} and Rita Prosmiti ^{1,*}

¹ Institute of Fundamental Physics (IFF-CSIC), CSIC, Serrano 123, 28006 Madrid, Spain;
juditmontesdeoca@iff.csic.es

² Atelgraphics S.L., Mota de Cuervo 42, 28043 Madrid, Spain

* Correspondence: rita@iff.csic.es; Tel.: +34-91-5616800 (ext. 442292)












† Doctoral Programme in Theoretical Chemistry and Computational Modelling, Doctoral School, Universidad Autónoma de Madrid, 28049 Madrid, Spain.













SUPPLEMENTARY MATERIAL



Table S1: Binding energies (in cm^{-1}) from CCSD(T)/AV6Z and MP2/AVQZ calculations for $[\text{Ar}_n\text{H}]^+$ complexes, with $n=1-3$ and 1-9, respectively, $(V_{2B}/V_{3B})V_{4B}$ and previous theoretical work [8]. The corresponding optimized structures (side and top views) obtained from the CCSD(T)/AV6Z calculations for $n=1-3$ and V_{4B} potential for $n=3-24$ are also depicted.

$[\text{Ar}_n\text{H}]^+$		This work		Ref. [8]	
n		MP2/AVQZ	$(V_{2B}/V_{3B})V_{4B}$	-MP2/AVTZ	
	Side view	Top view	CCSD(T)/AV6Z		
1			-32441.82	-	-31180
			-32799.69		
2			-5794.70	(-3097/-)	-5739
			-5353.75		
3			-767.20	(-5295/-861)	-718
			-712.73		
4			-848.09	(-5231/-867)-831.17	-796
5			-833.81	-838.77	-784
6			-833.41	-846.66	-777

7			-911.34 -944.05 -855
8			-605.46 -632.33 -557
9			-630.05 -727.03 -
10			- -736.31 -
11			- -746.36 -

12			- - -825.62 - -
13			- - -992.76 - -
14			- - -638.93 - -
15			- - -741.55 - -
16			- - -747.65 - -
17			- - -757.01 - -

18		
19		
20		
21		
22		
23		

24			—	-513.60	-