

Supplementary Materials

Cholesteryl Phenolipids as Potential Biomembrane Antioxidants

Vânia Costa ¹, Marlene Costa ¹, Francisca Arques ¹, Mariana Ferreira ¹, Paula Gameiro ¹, Dulce Geraldo ², Luís S. Monteiro ², Fátima Paiva-Martins ^{1,*}

¹ REQUIMTE-LAQV, Department of Chemistry and Biochemistry, Faculty of Sciences, University of Porto, Portugal, mpmartin@fc.up.pt

² Chemistry Centre, University of Minho, Gualtar, 4710-057 Braga, Portugal; monteiro@quimica.uminho.pt

* Correspondence: mpmartin@fc.up.pt




Figure S1 – ^1H -NMR spectra (400 MHz, $(\text{CD}_3)_2\text{CO}$) of cholesteryl caffeate.




Figure S2 – ^{13}C -NMR spectra (100 MHz, $(\text{CD}_3)_2\text{CO}$) of cholesteryl caffeate.




Figure S3 – ^1H -NMR spectra (400 MHz, CDCl_3) of cholesteryl dihydrocaffeate.




Figure S4 – ^{13}C -NMR spectra (100 MHz, CDCl_3) of cholesteryl dihydrocaffeate.




Figure S5 – ^1H -NMR spectra (400 MHz, CDCl_3) of cholesteryl homoprotocatechuate.




Figure S6 – ^{13}C -NMR spectra (100 MHz, CDCl_3) of cholesteryl homoprotocatechuate.

