

Supplementary Materials

Increased trypanocidal activity of the salinomycin derivative ironomycin is due to ROS production and iron uptake impairment


Dietmar Steverding ^{1,*}, Stuart A. Rushworth ¹, Georgina R. Hurle ², Michał Antoszczak ³, Michał Sulik ³, Adam Huczynski ³ and Kevin M. Tyler ²

¹ Bob Champion Research and Education Building, Norwich Medical School, University of East Anglia, Norwich NR4 7UQ, United Kingdom; D.Steverding@uea.ac.uk (D.S.); S.Rushworth@uea.ac.uk (S.A.R.)

² BioMedical Research Centre, Norwich Medical School, University of East Anglia, Norwich NR4 7TJ, United Kingdom; G.Hurle@uea.ac.uk (G.R.H.); K.Tyler@uea.ac.uk (K.M.T.)

³ Department of Medical Chemistry, Faculty of Chemistry, Adam Mickiewicz University, Uniwersytetu Poznańskiego 8, 61–614 Poznań, Poland; michal.antoszczak@amu.edu.pl (M.A.); michal.sulik@amu.edu.pl (M.S.); adam.huczynski@amu.edu.pl (A.H.)

* Correspondence: D.Steverding@uea.ac.uk



Scheme S1. Reduction of RhoNox-1 by iron(II) to rhodamine B.




Figure S1. ¹H-NMR (top) and ¹³C-NMR (bottom) spectra of RhoNox-1 in CDCl₃.