

Supplementary Information

Table S1. Primers used for RT-PCR.

Primer	Sequences (forward and reverse)	Distribution	AT (°C)	Product size (bp)	Extension time (s)
<i>BTG1</i>	5'-CCCTTCCTGGGCATCGT-3'	L10240.1	60	97	34
	5'-CGGCGTCGTCATCATCC-3'	645–741			
<i>PI3K</i>	5'-CACCGCATTTGTCGT-3'	NM_001242466	60	225	34
	5'-CTCCCACTTCTACGC-3'	4016–4240			
<i>Akt</i>	5'-GGGACAGAGGAGCAAGGT-3'	NM_005163	60	208	34
	5'-CGACAGCGGAAAGGTAA-3'	2656–2863			
<i>Bcl xl</i>	5'-TTCCCAGAAAGGATACAGC-3'	Z23115	60	92	34
	5'-GGGTCTCCATCTCCGATT-3'	185–276			
<i>survivin</i>	5'-CTTGGCCCAGTGTTTCTT-3'	DQ227257	60	124	34
	5'-GCTTCCAGTCCCTCCCT-3'	159–282			
<i>VEGF</i>	5'-AGGAGGAGGGCAGAATC-3'	NM_001171630	60	258	34
	5'-ATGTGCTGGCCTTGGT-3'	1131–1388			
<i>MMP-2</i>	5'-GAGCACTCCCAAGACCCT-3'	NM_001127891	60	255	34
	5'-AGTCCGCCAAATGAACC-3'	1398–1652			
<i>GAPDH</i>	5'-CAATGACCCCTTCATTGACC-3'	NM_002046.3	60	135	34
	5'-TGGAAGATGGTGATGGGATT-3'	201–335			

AT, annealing temperature; *GADPH*, glyceraldehyde-3-phosphate dehydrogenase.

© 2013 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/3.0/>).