

Supplementary Materials: Time-Dependent Nerve Growth Factor Signaling Changes in the Rat Retina during Optic Nerve Crush-Induced Degeneration of Retinal Ganglion Cells

Louise A. Mesentier-Louro, Sara De Nicolò, Pamela Rosso, Luigi A. De Vitis, Valerio Castoldi, Letizia Leocani, Rosalia Mendez-Otero, Marcelo F. Santiago, Paola Tirassa, Paolo Rama and Alessandro Lambiase

Table S1. Number of Tuj1- and Brn3a-positive cells in the retina. Table shows the number of cells per square millimeter of retina and SEM. The number of experiments (*n*) is indicated at each point.

	CoEye			7 dac			14 dac		
Central Retina	cells/mm ²	SEM	<i>n</i>	cells/mm ²	SEM	<i>n</i>	cells/mm ²	SEM	<i>n</i>
Brn3a ⁺	1842	39.24	16	876.7	64.86	8	138.3	12.79	8
Tuj1 ⁺	1264	52.31	14	746.8	48.62	8	164.3	11.29	8
Peripheral Retina	cells/mm ²	SEM	<i>n</i>	cells/mm ²	SEM	<i>n</i>	cells/mm ²	SEM	<i>n</i>
Brn3a ⁺	922.8	26.09	16	517.7	42.85	8	115.2	10.55	8
Tuj1 ⁺	1170	47.75	15	949.8	77.81	8	243.7	19.68	8