

Article

Can Selenium and Molybdenum Restrain Cadmium Toxicity to Pollen Grains in *Brassica napus*?

Marwa A. Ismael ^{1,2,3}, Ali Mohamed Elyamine ^{1,2}, Yuan Yuan Zhao ^{1,2}, Mohamed G. Moussa ^{1,2,4}, Muhammad Shoaib Rana ^{1,2}, Javaria Afzal ^{1,2}, Muhammad Imran ^{1,2}, Xiao Hu Zhao ^{1,2} and Cheng Xiao Hu ^{1,2,*}

¹ Key Laboratory of Arable Land Conservation (Middle and Lower Reaches of Yangtze River), Ministry of Agriculture, Huazhong Agricultural University, Wuhan 430070, China; maf02@fayoum.edu.eg (M.A.I.); elyoh@hotmail.fr (A.M.E.); yuanyuanzhao666@gmail.com (Y.Y.Z.); MohamedGomaa_Ali@agr.asu.edu.eg (M.G.M.); muhammadshoaib@webmail.hzau.edu.cn (M.S.R.); juvaria_afzal@outlook.com (J.A.); imrangorayauaf@yahoo.com (M.I.); xhzaowhu@163.com (X.H.Z.); hucx@mail.hzau.edu.cn (C.X.H.)

² Hubei Provincial Engineering Laboratory for New-Type Fertilizers, Huazhong Agricultural University, Wuhan 430070, China

³ Botany Department, Faculty of Science, Fayoum University, Fayoum 63514, Egypt

⁴ Soil and Water Research Department, Nuclear Research Center, Egyptian Atomic Energy Authority, 13759, Abou Zaabl, Egypt.

* Correspondence: hucx@mail.hzau.edu.cn

Table S1. Sequences of primers used for RT-PCR

Target	Primer sequences (5'-3')	Annealing Temperature	Gene IDs
<i>BnActin2.1</i>	F-CTCTTTCACACGCCATCCTCC R-GATTCCAGCAGCTTCCATTCC	57.5 C	
<i>IRT1</i>	F- GCCTGATTCTTTCGAGATGT R- TGCATGCTTGCTAGTGATAG	58 °C	BnaA03g44160D
<i>HMA2</i>	F- CTTTGGCTACGGCTGATATT R- CATTCTCCACCACTTTCCTT	58 °C	BnaC07g50890D
<i>HMA3</i>	F-GGCTGGAGTTTACCCTATTC R-CCAATCTGCAACCGAGAA	58 °C	BnaA01g06890D
<i>HMA4</i>	F-GCAGTCACTGATGAGCTTAG R-TTTCCATCCACCACAATACC	58 °C	BnaC03g48480D
<i>PCS1</i>	F-GACCACCATTGACGACTTT R-GCCATATCTCTCTCAGCATTAT	58 °C	BnaA06g36340D

Note: 'F' was defined as forward primer, and 'R' as reverse primer.