

Rare-earth-doped Calcium Carbonate exposed to X-ray to induce Reactive Oxygen Species for Tumor Treatment

Chun-Chen Yang^a, Wei-Yun Wang^a, Feng-Huei Lin^{b, c, *}, Chun-Han Hou^{d, *}

^a Department of Materials Science and Engineering, National Taiwan University, Taipei 10617, Taiwan

^b Institute of Biomedical Engineering, National Taiwan University, Taipei 10617, Taiwan

^c Institute of Biomedical Engineering and Nanomedicine, National Health Research Institutes, Miaoli 35053, Taiwan

^d Department of Orthopedic Surgery, National Taiwan University, Taipei 10617, Taiwan

Corresponding Author

* Chun-Han Hou: phone, +886-2-23123456#65274; fax, +886-2-23971266; Email, chhou@ntu.edu.tw.

* Feng-Huei Lin: phone, +886-2-27327474; fax, +886-2-27327474; Email, double@ntu.edu.tw.

Table S1. CaCO₃:Ce measured by BET.

CaCO ₃ :Ce	BET Surface Area : 35.8875±0.1818 m ² /g Correlation Coefficient: 0.9999237 Langmuir Surface Area: 48.6989±0.6808 m ² /g
-----------------------	--

	Correlation Coefficient: 0.9996093 BJH Adsorption Average Pore Diameter : 1295.3808 Å BJH Desorption Average Pore Diameter : 1228.4492Å
--	---