

Comparative study of the genetic and biochemical variability of *Polyscias filicifolia* (Araliaceae) regenerants obtained by indirect and direct somatic embryogenesis as a source of triterpenes

Anita A. Śliwińska ¹, Agnieszka Białek ², Renata Orłowska ³, Dariusz Mańkowski ³, Katarzyna Sykłowska-Baranek ^{1*} and Agnieszka Pietrosiuk ¹

¹ Department of Pharmaceutical Biology and Medicinal Plant Biotechnology, Faculty of Pharmacy, Medical University of Warsaw, Banacha 1, 02-097 Warsaw, Poland; asliwinska@wum.edu.pl (AŚ); katarzyna.sykłowska-baranek@wum.edu.pl (KSB); agnieszka.pietrosiuk@wum.edu.pl (AP)

² Department of Biotechnology and Nutrigenomics, Institute of Genetics and Animal Biotechnology, Polish Academy of Sciences, Postępu 36 A, Jastrzębiec, 05-552 Magdalanka, Poland; a.bialek@ighz.pl (AB)

³ Plant Breeding and Acclimatization Institute-National Research Institute, Radzików, 05-870 Błonie, Poland; r.orłowska@ihar.edu.pl (RO); d.mankowski@ihar.edu.pl (DM)

* Correspondence: katarzyna.sykłowska-baranek@wum.edu.pl (KSB)

Table S3. Summary of metAFLP data for three groups of regenerants PSE, DSE, TSE of *Polysia filicifolia*

| Type of variation | % of variation | | |
|----------------------------|----------------|------|------|
| | PSE | DSE | TSE |
| Sequence variation | 1.25 | 1.11 | 0.75 |
| Demethylation | 0.54 | 0.56 | 0.84 |
| <i>De novo</i> methylation | 0.41 | 0.25 | 0.18 |
| Total | 2.20 | 1.92 | 1.77 |