

Antagonizing S1P₃ Receptor with Cell-Penetrating Pepducins in Skeletal Muscle Fibrosis

Angela Corvino ¹, Ida Cerqua ¹, Alessandra Lo Bianco ¹, Giuseppe Caliendo ¹, Ferdinando Fiorino ¹, Francesco Frecentese ¹, Elisa Magli ¹, Elena Morelli ¹, Elisa Perissutti ¹, Vincenzo Santagada ¹, Giuseppe Cirino ¹, Elisabetta Granato ¹, Fiorentina Roviezzo ¹, Elisa Puliti ², Caterina Bernacchioni ², Antonio Lavecchia ¹, Chiara Donati ² and Beatrice Severino ^{1,*†}

¹ Department of Pharmacy, School of Medicine, University of Naples «Federico II», Via D. Montesano, 49, 80131 Napoli, Italy; angela.corvino@unina.it (A.C.); ida.cerqua@unina.it (I.C.); alessandra.lobianco@unina.it (A.L.B.); caliendo@unina.it (G.C.); fefiorin@unina.it (F.F.); frecente@unina.it (F.F.); elisa.magli@unina.it (E.M.); elena.morelli@unina.it (E.M.); perissut@unina.it (E.P.); santagad@unina.it (V.S.); cirino@unina.it (G.C.); elisabetta.granato@unina.it (E.G.); roviezzo@unina.it (F.R.); antonio.lavecchia@unina.it (A.L.)

² Department of Experimental and Clinical Biomedical Sciences “Mario Serio”, University of Florence, Viale GB Morgagni 50, 50134 Firenze, Italy; elisa.puliti@unifi.it (E.P.); caterina.bernacchioni@unifi.it (C.B.); chiara.donati@unifi.it (C.D.)

* Correspondence: bseverin@unina.it; Tel.: +39-081-679-828

† Co-last authors.




Figure S1. (a) Time evolution of the backbone RMSD compared to frame 0 for **KRX-725-II** (blue), compound **V** (green) and compound **VII** (orange) during 1000 ns of MD. (b) Variation of the Rg with time for **KRX-725-II** (blue), **V** (green) and **VII** (orange). (c) Total number of intramolecular H-bonds formed by **KRX-725-II** (blue), **V** (green) and **VII** (orange) during the entire simulation period.