

Table S1

Primers used for qRT-PCR analysis of genes important for hepatic lipid metabolism

<i>Srebp1c</i>	Forward primer	TACAGCGTGGCTGGGAAC
	Reverse primer	CTCCCTGTCTCCGTCAGC
<i>Ppara</i>	Forward primer	CTGAGACCCTCGGGGAAC
	Reverse primer	AAACGTCAGTTCACAGGGAAG
<i>Slc27a2</i>	Forward primer	GCGTGCCTCAACTACAACATT
	Reverse primer	CCTCCTCCACAGCTTCTTGT
<i>Slc27a4</i>	Forward primer	CTTGCCTGAGCTGCACAA
	Reverse primer	GCGGGTCTTTCACAACAGAT
<i>Slc27a5</i>	Forward primer	CTGCGGTACTTGTGTAAACGTCC
	Reverse primer	TCCGAATGGGACCAAAGCGTTG
<i>Acc1</i>	Forward primer	GCGTCGGGTAGATCCAGTT
	Reverse primer	CTCAGTGGGGCTTAGCTCTG
<i>Fas</i>	Forward primer	TACAGCGTGGCTGGGAAC
	Reverse primer	CTCCCTGTCTCCGTCAGC
<i>Acox1</i>	Forward primer	ATCAGGGCACCCTGCTC
	Reverse primer	CCAAGCCTCGAAGATGAGTT
<i>Cpt1a</i>	Forward primer	GACTCCGCTCGCTCATTC
	Reverse primer	TCTGCCATCTTGAGTGGTGA
<i>Mttp</i>	Forward primer	GCCCAACGTA CTCTAATTTATGG
	Reverse primer	TGCTGGCCAACACGTCTA

Primers used for qRT-PCR analysis of genes important for hepatic lipid metabolism