



Review

# Allosteric Interactions between Adenosine A<sub>2A</sub> and Dopamine D<sub>2</sub> Receptors in Heteromeric Complexes: Biochemical and Pharmacological Characteristics, and Opportunities for PET Imaging

Kavya Prasad <sup>1,\*</sup>, Erik F. J. de Vries <sup>1</sup> , Philip H. Elsinga <sup>1</sup>, Rudi A. J. O. Dierckx <sup>1,2</sup> and Aren van Waarde <sup>1,\*</sup>

<sup>1</sup> Department of Nuclear Medicine and Molecular Imaging, University Medical Center Groningen, University of Groningen, Hanzelplein 1, 9713GZ Groningen, The Netherlands; e.f.j.de.vries@umcg.nl (E.F.J.d.V.); p.h.elsinga@umcg.nl (P.H.E.); r.a.dierckx@umcg.nl (R.A.J.O.D.)

<sup>2</sup> Department of Diagnostic Sciences, Ghent University Faculty of Medicine and Health Sciences, C.Heymanslaan 10, 9000 Gent, Belgium

\* Correspondence: k.prasad@umcg.nl (K.P.); a.van.waarde@umcg.nl (A.v.W.); Tel.: +31-50-3613215



**Citation:** Prasad, K.; de Vries, E.F.J.; Elsinga, P.H.; Dierckx, R.A.J.O.; van Waarde, A. Allosteric Interactions between Adenosine A<sub>2A</sub> and Dopamine D<sub>2</sub> Receptors in Heteromeric Complexes: Biochemical and Pharmacological Characteristics, and Opportunities for PET Imaging. *Int. J. Mol. Sci.* **2021**, *22*, 1719. <https://doi.org/10.3390/ijms22041719>

Academic Editor: Xavier Altafaj

Received: 22 January 2021

Accepted: 3 February 2021

Published: 9 February 2021

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.




**Copyright:** © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

**Abstract:** Adenosine and dopamine interact antagonistically in living mammals. These interactions are mediated via adenosine A<sub>2A</sub> and dopamine D<sub>2</sub> receptors (R). Stimulation of A<sub>2A</sub>R inhibits and blockade of A<sub>2A</sub>R enhances D<sub>2</sub>R-mediated locomotor activation and goal-directed behavior in rodents. In striatal membrane preparations, adenosine decreases both the affinity and the signal transduction of D<sub>2</sub>R via its interaction with A<sub>2A</sub>R. Reciprocal A<sub>2A</sub>R/D<sub>2</sub>R interactions occur mainly in striatopallidal GABAergic medium spiny neurons (MSNs) of the indirect pathway that are involved in motor control, and in striatal astrocytes. In the nucleus accumbens, they also take place in MSNs involved in reward-related behavior. A<sub>2A</sub>R and D<sub>2</sub>R co-aggregate, co-internalize, and co-desensitize. They are at very close distance in biomembranes and form heteromers. Antagonistic interactions between adenosine and dopamine are (at least partially) caused by allosteric receptor–receptor interactions within A<sub>2A</sub>R/D<sub>2</sub>R heteromeric complexes. Such interactions may be exploited in novel strategies for the treatment of Parkinson’s disease, schizophrenia, substance abuse, and perhaps also attention deficit-hyperactivity disorder. Little is known about shifting A<sub>2A</sub>R/D<sub>2</sub>R heteromer/homodimer equilibria in the brain. Positron emission tomography with suitable ligands may provide in vivo information about receptor crosstalk in the living organism. Some experimental approaches, and strategies for the design of novel imaging agents (e.g., heterobivalent ligands) are proposed in this review.

**Keywords:** adenosine A<sub>2A</sub> receptor; dopamine D<sub>2</sub> receptor; heteromers; allosteric interaction; receptor–receptor interactions; striatum; GABAergic enkephalinergic neuron

## 1. Introduction

Adenosine, a purine nucleoside, plays several behavioral and physiological roles throughout the central nervous system (CNS). Adenosine is generated in the living brain from adenine nucleotides such as adenosine triphosphate (ATP) and adenosine monophosphate (AMP). A much less important, other source of adenosine is S-adenosylhomocysteine, that originates from S-adenosylmethionine after physiological transmethylation [1]. Increased firing of neurons is associated with increased consumption of ATP, nucleotide dephosphorylation, and increases of intracellular adenosine levels (Figure 1). Since equilibrative nucleoside transporters are present in neuronal membranes, the extracellular levels of adenosine will also increase under such conditions. Thus, extracellular adenosine concentrations fluctuate, depending on neuronal activity.



**Figure 1.** Metabolic pathways involved in the formation and removal of adenosine. 1 = Oxidative phosphorylation (and creatine kinase), 2 = Energy-consuming processes, 3 = Adenylate kinase, 4 = Apyrase, 5 = Adenylate cyclase, 6 = Phosphodiesterase, 7 = 5'-Nucleotidase, 8 = S-adenosyl homocysteine hydrolase, 9 = Adenosine kinase, 10 = Adenosine deaminase, 11 = Purine phosphorylase, 12 = Xanthine oxidase. ATP = adenosine 5'-triphosphate, ADP = adenosine 5'-diphosphate, AMP = adenosine 5'-monophosphate, cAMP = 3',5'-cyclic adenosine monophosphate.

Extracellular adenosine levels in the mammalian brain range from 20 to 250 nM [2–7]. Extracellular adenosine can bind to four subtypes of adenosine receptors, called A<sub>1</sub>, A<sub>2A</sub>, A<sub>2B</sub> and A<sub>3</sub>, which belong to the P1 receptor family. A<sub>1</sub> and A<sub>2A</sub> receptors have a high affinity for adenosine (10–100 nM range), whereas A<sub>2B</sub> and A<sub>3</sub> receptors are only activated when extracellular adenosine reaches very high (micromolar) levels, after tissue damage (e.g., inflammation, hypoxia, ischemia, brain injury). Physiological levels of adenosine will stimulate the A<sub>1</sub> and A<sub>2A</sub> receptors. It is unlikely that adenosine exerts major physiological functions via A<sub>2B</sub> and A<sub>3</sub> receptors in the brain, since physiological levels of adenosine are too low to activate these proteins, and A<sub>2B</sub> and A<sub>3</sub> receptors are mainly expressed in peripheral organs rather than in the CNS [8–13].

A<sub>1</sub> receptors (A<sub>1</sub>R) are coupled to Gi proteins. Stimulation of these receptors by adenosine causes a decrease in cAMP levels through an inhibitory effect on adenylate cyclase. A<sub>2A</sub> receptors (A<sub>2A</sub>R) are coupled to an excitatory Gs protein. Stimulation of A<sub>2A</sub>R results in an increase of cAMP levels and activation of protein kinase A [8–13].

## 2. Antagonistic Interactions between Adenosine and Dopamine

### 2.1. Living Animals

Interactions between adenosine and dopamine in living animals were already observed in 1974. Adenosine antagonists (caffeine and theophyllamine) were then reported to enhance the action of dopamine agonists such as apomorphine, bromocriptine and L-DOPA (stimulation of rotation behavior) in the 6-hydroxydopamine hemiparkinson model of rats [14]. In later studies using reserpinized (i.e., dopamine-depleted) mice, the action of bromocriptine was found to be inhibited by adenosine agonists (L-PIA, NECA)

and this inhibition could be reversed by the adenosine antagonists caffeine, paraxanthine, and theophylline. Since the non-subtype-selective agonist 5'-(N-ethyl)carboxamido-adenosine (NECA) was considerably more potent than the A<sub>1</sub>-selective agonist N6-R-phenylisopropyladenosine (L-PIA), A<sub>2A</sub> rather than A<sub>1</sub> receptors seem to be involved in the inhibition of the locomotor response to dopaminergic stimulation [15,16]. Central administration of the adenosine A<sub>2A</sub>R agonist 2-[p-(2-carboxyethyl)phenethylamino]-5'-N-ethylcarboxamido-adenosine (CGS21680) was shown to induce catalepsy in the rat, and this effect was counteracted by systemic administration of the adenosine antagonist theophylline or the dopamine D<sub>2</sub> agonist 5,6,7,8-Tetrahydro-6-(2-propen-1-yl)-4H-thiazolo[4,5-d]azepin-2-amine dihydrochloride (BHT-920) [17]. The dopamine D<sub>2</sub>R antagonist haloperidol induces catalepsy and Parkinsonian symptoms in rats and mice. Such symptoms can be reversed by treating rats with the non-selective adenosine antagonist caffeine or the selective A<sub>2A</sub>R antagonist SCH58261 [18] and are significantly reduced in A<sub>2A</sub>R knockout mice [19]. Haloperidol-induced motor impairments in monkeys (catalepsy, extrapyramidal syndrome) are counteracted by the A<sub>2A</sub>R antagonists SCH-412348, istradefylline, and caffeine [20].

As A<sub>2A</sub> receptors are known to be located mainly in the striatum, in postsynaptic locations on dendrites and dendritic spines [21,22] and, to a lesser extent (25%), on nerve endings [23,24]. These findings suggest the existence of postsynaptic interactions between adenosine and dopamine receptors, probably the A<sub>2A</sub> and D<sub>2</sub> subtypes. Stimulation of A<sub>2A</sub> receptors results in inhibition, and blockade of A<sub>2A</sub> receptors in enhancement of D<sub>2</sub>-receptor mediated locomotor activation.

Stimulation of A<sub>2A</sub>R in the nucleus accumbens of rats by local infusion of CGS21680 produced behavioral effects similar to those induced by local dopamine depletion (i.e., decreased lever pressing for preferred food and substantially increased consumption of the less preferred but freely accessible chow) [25]. On the other hand, decreases of lever pressing for preferred (high carbohydrate) food caused by the D<sub>2</sub>R antagonist eticlopride could be partially reversed by treating rats with the A<sub>2A</sub>R antagonist MSX-3 [26]. Similar decreases induced by the D<sub>2</sub>R antagonist haloperidol could be reversed by the A<sub>2A</sub>R-subtype-selective antagonist istradefylline or the non-subtype selective AR antagonist caffeine [27]. Thus, antagonistic interactions between A<sub>2A</sub>R and D<sub>2</sub>R occur not only in the dorsal striatum where they control locomotor activity, but also in the nucleus accumbens (ventral striatum) where they affect goal-directed behavior.

## 2.2. Membrane Preparations

Antagonistic interactions between A<sub>2A</sub> and D<sub>2</sub> receptors could also be observed in vitro, in membrane preparations from rat striatum. Administration of the adenosine A<sub>2A</sub> receptor (A<sub>2A</sub>R) agonist CGS21680 resulted in a significant, 40% increase of the K<sub>d</sub> (i.e., a loss of the affinity) of dopamine D<sub>2</sub> receptors to the agonist L-(*–*)-N-[<sup>3</sup>H]propylnorapomorphine without changing the B<sub>max</sub> (i.e., the number of D<sub>2</sub> receptors) [28]. However, the K<sub>d</sub> and B<sub>max</sub> for binding of the dopamine D<sub>2</sub> antagonist [<sup>3</sup>H]raclopride were not affected [28]. The effect of CGS21680 on D<sub>2</sub>R affinity was most pronounced at concentrations similar to the K<sub>d</sub> for binding of CGS21680 to A<sub>2A</sub>R. At very high, saturating doses of CGS21680 (300 nM), the effect of the agonist was reduced, probably because such high doses cause a desensitization of A<sub>2A</sub>R [28]. In striatal membrane preparations of adult (as opposed to young) rats, CGS21680 reduced not only the affinity of D<sub>2</sub> receptors for agonists, but also the fraction of D<sub>2</sub> receptors in the high-affinity state. Thus, A<sub>2A</sub>R stimulation may inhibit the motor responses induced by dopamine receptor agonists by decreasing both the affinity and the signal transduction of D<sub>2</sub> receptors [29,30]. Adenosine appears to regulate the properties of D<sub>2</sub>R via its interaction with A<sub>2A</sub>R. Direct receptor–receptor interactions in striatal membranes were suggested as a potential mechanism involved in this pharmacological crosstalk between A<sub>2A</sub>R and D<sub>2</sub>R [28,31,32].

### 2.3. Intact Cells

Antagonistic interactions between A<sub>2A</sub> and D<sub>2</sub> receptors were also demonstrated in intact cells. In a mouse fibroblast cell line stably transfected with A<sub>2A</sub>R and D<sub>2</sub>R, the D<sub>2</sub>R agonist quinpirole induced a concentration-dependent increase in intracellular (cytosolic) free calcium. This response was completely blocked if cells were pretreated with haloperidol. CGS21680 by itself did not affect intracellular calcium levels (even when it was administered at high dose), but CGS21680 strongly counteracted the response of [Ca<sup>2+</sup>]<sub>i</sub> to quinpirole [33]. Similar observations were made in SH-SY5Y (human neuroblastoma) cells that were transfected with human D<sub>2</sub>R [34]. The effect of CGS21680 was shown to be related to a two- to three-fold decrease of the affinity of the D<sub>2</sub>R in the cells to dopamine receptor agonists [34,35]. A similar three- to four-fold increase of the K<sub>D</sub> of dopamine at high-affinity D<sub>2</sub>R sites after administration of CGS21680 was noted in Chinese hamster ovary (CHO) cells that were co-transfected with A<sub>2A</sub> and D<sub>2</sub> receptors [36]. In such cells, CGS21680 decreased the affinity of D<sub>2</sub> receptors for [<sup>3</sup>H]dopamine but not the number of dopamine binding sites [37]. Since A<sub>2A</sub>R stimulation increases, but D<sub>2</sub>R stimulation decreases, the intracellular formation of cyclic AMP, A<sub>2A</sub>R, and D<sub>2</sub>R may interact not only at the membrane level but also at the second messenger level. The experiments in CHO cells suggested that the latter interaction may be quantitatively the most important [36].

In initial cell experiments, A<sub>2A</sub>R agonists were shown to decrease the affinity of D<sub>2</sub>R for agonists. In later experiments, interactions in the opposite direction were also demonstrated. D<sub>2</sub>R activation by quinpirole resulted in a less rapid and reduced binding of the fluorescent A<sub>2A</sub>R agonist MRS5424 to HEK293 cells, which expressed both A<sub>2A</sub> and D<sub>2</sub> receptors [38]. Similar decreases of A<sub>2A</sub>R agonist binding were observed when the cells were treated with D<sub>2</sub>R agonists in clinical use, such as pramipexole, rotigotine, and apomorphine [39]. On the other hand, chronic D<sub>2</sub>R blockade by haloperidol increased both the affinity and the responsiveness of the A<sub>2A</sub>R to the agonist NECA in CHO cells that expressed both A<sub>2A</sub> and D<sub>2</sub> receptors [40].

In CHO cells transiently transfected with A<sub>2A</sub>R and D<sub>2</sub>R, both the A<sub>2A</sub>R agonist CGS21680 and the AR antagonist caffeine caused a decrease of the affinity of the D<sub>2</sub>R for radioligands, not only the D<sub>2</sub>R agonist [<sup>3</sup>H]quinpirole but also the D<sub>2</sub>R antagonist [<sup>3</sup>H]raclopride. Yet, CGS21680 and caffeine canceled out each other's effect on D<sub>2</sub>R affinity when they were administered together [41]. These apparently paradoxical findings led to a novel hypothesis concerning the structural basis of adenosine–dopamine receptor interactions, which is described in Section 5 of this review.

### 2.4. Brain Slices

Antagonistic interactions between A<sub>2A</sub> and D<sub>2</sub> receptors could also be demonstrated in cryostat sections of rat and human brain. CGS21680 significantly increased the IC<sub>50</sub> values of competition between the D<sub>2/3</sub>R ligand [<sup>125</sup>I]iodosulpiride and dopamine in the striatal region of such preparations [42].

## 3. Regional, Cellular, and Subcellular Distribution of A<sub>2A</sub> and D<sub>2</sub> Receptors

The antagonistic interactions of A<sub>2A</sub> and D<sub>2</sub> receptors that were observed in rat striatal membranes [28–30] suggested that the A<sub>2A</sub> and D<sub>2</sub> receptor genes are co-expressed by some cells in the mammalian brain.


### 3.1. Regional Distribution

Both in the rodent and human brain, A<sub>2A</sub>R mRNA [43–48] and A<sub>2A</sub>R protein [24,49–56] are mainly located in the striatum (caudate-putamen) and nucleus accumbens. In monkeys, A<sub>2A</sub>R immunoreactivity is mainly present in striatum and nucleus accumbens, but can also be detected in the substantia nigra, an area showing very low A<sub>2A</sub>R density in rats. This finding indicates that there may be species differences between rodents and primates concerning the regional distribution of A<sub>2A</sub>R [57].

Caudate, putamen and nucleus accumbens express also high numbers of dopamine D<sub>2</sub>R [58–62]. The distribution of D<sub>2</sub>R in the rodent brain is very similar to that of A<sub>2A</sub>R mRNA, although D<sub>2</sub>R is also present in the substantia nigra and piriform cortex [63].

### 3.2. Cellular Distribution

The majority (more than 95%) of the neurons in the striatum are medium spiny neurons (MSNs; i.e., medium-sized neurons (diameter 12–15 µm in rodents) with large and extensive dendritic trees) [64]. MSNs in the dorsal striatum can be divided in two subtypes [65,66]. Both subtypes use gamma-aminobutyric acid (GABA) as neurotransmitter, but the subtypes have different projection patterns and they express different receptors and neuropeptides. Some MSNs send direct (monosynaptic) projections to the substantia nigra and the globus pallidus internus. Based on this projection pattern, this subtype is said to form part of the “direct pathway” (Figure 2). MSNs of the direct pathway express dopamine D1R and the peptide dynorphin (together with substance P). Other MSNs are indirectly linked to the substantia nigra and the globus pallidus internus, via the globus pallidus externus and the subthalamic nucleus. Because of this distinctive projection pattern, they are said to form part of the “indirect pathway” (Figure 2). MSNs of the indirect pathway express dopamine D<sub>2</sub>R and the peptide enkephalin [63,67,68] (reviewed in [69]).



**Figure 2.** Schematic drawing of the direct and indirect pathways for motor control. Solid and faded lines represent direct and indirect pathways, respectively. Blue lines represent excitatory connections and red lines represent inhibitory connections. Pu = putamen, Gpe = globus pallidus externus, Gpi = globus pallidus internus, STN = subthalamic nucleus, SNC = substantia nigra pars compacta, VA/VL = ventral anterior/ventral lateral thalamic nucleus. Created with BioRender.com.

In early publications, the A<sub>2A</sub>R (at that time still called RDC8) was shown to be present in medium-sized but not in large neurons of the dog and rat striatum [70]. In contrast to A<sub>2A</sub>R, dopamine D<sub>2</sub>R mRNA is present both in medium-sized and large neurons [71]. Later studies employed double *in situ* hybridization [43,44,47,72,73] and double-labeling immunohistochemistry [52] to determine the phenotype of A<sub>2A</sub>R-containing neurons in dorsal and ventral striatum. In the ventral striatum, a population of neurons expresses the gene for the A<sub>2A</sub>R, but not for preproenkephalin. This sub-population is absent in the dorsal striatum. In the dorsal striatum, 95–96% of the A<sub>2A</sub>R mRNA is co-expressed with D<sub>2</sub>R mRNA. Only a few neurons expressing 3–6% of the A<sub>2A</sub>R mRNA co-express dopamine D<sub>1</sub>R or substance P mRNAs. In the ventral striatum, most A<sub>2A</sub>R mRNA (89–92%) co-localizes with preproenkephalin A mRNA, and the vast majority (93–95%) with D<sub>2</sub>R mRNA [74]. Adenosine A<sub>2A</sub> receptors were shown to co-localize with enkephalin and dopamine D<sub>2</sub>R, but not with dopamine D<sub>1</sub>R, substance P or somatostatin. These data were interpreted as evidence for a preferential expression of A<sub>2A</sub>R in striatopallidal GABAergic MSNs of the indirect pathway, cells which also express D<sub>2</sub>R [44,72,75]. Microdialysis experiments in intact freely moving rats supported this hypothesis. In these experiments, adenosine and dopamine agonists and antagonists were infused in the striatum, either alone or in combination, and the effect on the release of GABA was measured in the ipsilateral globus pallidus [76].

MSNs from the indirect pathway are the main, but not the only, cells in the striatum that co-express A<sub>2A</sub> and D<sub>2</sub> receptors. Striatal astrocytes also express both proteins [77–80] and receptor–receptor interactions between A<sub>2A</sub>R and D<sub>2</sub>R have been demonstrated in glia. Administration of the D<sub>2</sub>R agonist quinpirole to rat striatal astrocytes inhibits the 4-aminopyridine-provoked release of glutamate. The A<sub>2A</sub>R agonist CGS21680 alone did not affect glutamate release but reduced the D<sub>2</sub>R-mediated inhibiting effect of quinpirole [81]. A third class of cells in the striatum which express both A<sub>2A</sub>R and D<sub>2</sub>R are cholinergic interneurons [82].

### 3.3. Subcellular Location

In bright field photomicrographs of coronal sections of rat striatum, A<sub>2A</sub>R protein was detected on the cell bodies of GABA/enkephalin striatopallidal neurons [73]. Using immuno-electron microscopy, A<sub>2A</sub>Rs were mainly detected on dendrites, to a lesser extent on axon terminals, soma and astrocytic processes [23,24,52]. Subcellular fractionation experiments using the radioligand [<sup>3</sup>H]SCH58261 suggested that A<sub>2A</sub>R in the striatum of the rat are not enriched in synaptosomes [22]. In dendrites and soma, A<sub>2A</sub>R were shown to be present not only on the plasmalemma, but also throughout the cytoplasm and around intracellular membranous structures [23]. The predominantly postsynaptic location of A<sub>2A</sub>Rs (on dendrites and dendritic spines) was interpreted as evidence for an important function of these receptors in modulating the excitatory glutamatergic input to the striatum [24].

D<sub>2</sub> receptor immunoreactivity was detected by immunocytochemistry and electron micrography in rat basal ganglia. Subcellular experiments using fusion protein antibodies depicted predominant localization of D<sub>2</sub> in spiny dendrites and spine heads within the neutrophil of the striatum. The receptors were also located in submembranous sites of dendritic shafts and dendritic spines [83].

## 4. A<sub>2A</sub>R and D<sub>2</sub>R Co-Aggregate, Co-Internalize and Co-Desensitize

Interactions between A<sub>2A</sub> and D<sub>2</sub> receptors were found to affect not only the signaling but also the intracellular trafficking of the two proteins. The human neuroblastoma cell line SH-SY5Y constitutively expresses A<sub>2A</sub> receptors. In a groundbreaking article [84], SH-SY5Y cells were transfected with D<sub>2</sub> receptors, and incubated with fluorescein-conjugated anti-A<sub>2A</sub>R (green fluorescence) and rhodamine-conjugated anti-D<sub>2</sub>R antibodies (red fluorescence). Receptor trafficking in the cells could then be monitored with confocal laser microscopy. In untreated cells, A<sub>2A</sub>R and D<sub>2</sub>R were shown to be generally at close distance

(<100 nm) but rather uniformly distributed in the plasma membrane. When the cells were treated with either CGS21680 (100 nM) or quinpirole (10  $\mu$ M) for 3 h, the distribution of the receptors in the plasma membrane became less uniform and significant co-aggregates were formed (yellow hotspots). When the same doses of the A<sub>2A</sub>R and D<sub>2</sub>R agonist were administered together for 3 h, the total intensity of the fluorescence signals was decreased, suggesting that co-aggregation of the A<sub>2A</sub>R and D<sub>2</sub>R was followed by co-internalization. This effect was dose-dependent, both the co-aggregation and the signal loss being stronger after treatment with 200 nM CGS21680 plus 50  $\mu$ M quinpirole than with 100 nM CGS21680 plus 10  $\mu$ M quinpirole. In cells lacking D<sub>2</sub>R, quinpirole did not cause any aggregation or internalization of the A<sub>2A</sub>R. Prolonged (3 h) administration of either 1  $\mu$ M of CGS21680 or 1  $\mu$ M of quinpirole to cells expressing both A<sub>2A</sub>R and D<sub>2</sub>R resulted in desensitization of their A<sub>2A</sub> receptors (decrease of the cAMP response to A<sub>2A</sub>R stimulation), but desensitization of the D<sub>2</sub>R occurred only when both agonists were simultaneously administered [84]. When A<sub>2A</sub>R in the cells were immunoprecipitated with A<sub>2A</sub>R antibodies, Western blots indicated that the D<sub>2</sub>R was co-precipitated and that three glycosylated forms of the D<sub>2</sub>R were present in the precipitate [84]. Thus, A<sub>2A</sub>R and D<sub>2</sub>R were shown to co-aggregate, co-internalize, co-desensitize, and co-precipitate in the presence of D<sub>2</sub>R and A<sub>2A</sub>R agonists.

Computer-assisted analysis of dual-channel fluorescence laser microscopy images indicated co-localization, co-aggregation and co-internalization of A<sub>2A</sub>R and D<sub>2</sub>R also in Chinese hamster ovary (CHO) cells [85,86]. In the CHO cell experiments, the effect of receptor stimulation was examined at different time intervals (3, 15 and 24 h) after administration of quinpirole. Co-aggregation of A<sub>2A</sub>R and D<sub>2</sub>R was observed after 3 h, and the co-aggregates internalized after 15 h. A return to the plasma membrane was detected after 24 h. In contrast to treatment with quinpirole, treatment of CHO cells with the D<sub>2</sub>R antagonist raclopride did not decrease but increased the fluorescence signal of both A<sub>2A</sub>R and D<sub>2</sub>R, indicating that a D<sub>2</sub>R antagonist reduced the internalization of the two receptors [86].

Similar microscopy techniques suggested that A<sub>2A</sub> and D<sub>2</sub> receptors form a macrocomplex with caveolin-1 that internalizes when cells are treated with an A<sub>2A</sub> and a D<sub>2</sub> agonist. Thus, caveolin-1 may play a role in the process of co-internalization [87]. Later experiments using bioluminescence resonance energy transfer (BRET) indicated that A<sub>2A</sub> and D<sub>2</sub> receptors also form a macrocomplex with  $\beta$ -arrestin2, A<sub>2A</sub>R agonists promoting (and A<sub>2A</sub>R antagonists reducing) the D<sub>2</sub>R agonist-induced recruitment of  $\beta$ -arrestin2 by the D<sub>2</sub>R protomer and subsequent co-internalization [88,89].

The D<sub>2</sub>R agonist 3-(3,4-dimethylphenyl)-1-(2-piperidin-1-yl)ethyl)-piperidine was shown to reduce the affinity and functional responsiveness of A<sub>2A</sub>R to agonists. In addition, this D<sub>2</sub>R agonist induced co-internalization of the A<sub>2A</sub>R and D<sub>2</sub>R proteins [90].

## 5. A<sub>2A</sub>R and D<sub>2</sub>R Are at Very Close Distance in Biomembranes and Form Heteromers

At the end of the twentieth and beginning of the twenty-first century, several biophysical techniques, such as atomic force microscopy (AFM), bimolecular fluorescence complementation (BiFC), fluorescence resonance energy transfer (FRET), bioluminescence resonance energy transfer (BRET), in situ proximity ligation assay (PLA), and AlphaScreen technology, were developed that allow the detection of spatial proximity of protein molecules, and such techniques have also been applied to A<sub>2A</sub> and D<sub>2</sub> receptors [85,91–100]. The results of these techniques and the observed co-aggregation, co-internalization and co-immuno-precipitation of A<sub>2A</sub>R and D<sub>2</sub>R indicate that both receptors are at very close distance in biological membranes (<10 nm) and form heteromers. Molecular biology experiments have provided insight in the mechanisms and atomic interactions that are involved in heteromer formation.

Using BRET technology, Japanese authors demonstrated that A<sub>2A</sub>R form homomers and also heteromers with D<sub>2</sub>R in living HEK293T cells. A<sub>2A</sub> and D<sub>2</sub> receptors were fused to either an energy donor (Renilla luciferase) or an energy acceptor (modified green fluorescent protein) without affecting the ligand binding affinity, subcellular distribution or

co-immunoprecipitation of the two receptor proteins [101]. BRET and FRET techniques were also applied to quantify A<sub>2A</sub>R/D<sub>2</sub>R heteromers in receptor co-transfected cells, including cells that were transfected with modified D<sub>2</sub> receptors: Chimeric proteins in which part of the D<sub>2</sub> receptor protein was replaced by the corresponding part of the D<sub>1</sub> receptor protein. Such experiments, and molecular modeling studies, suggested that heteromerization between A<sub>2A</sub>R and D<sub>2</sub>R depends on interaction of the third intracellular loop of the D<sub>2</sub>R with the C-terminal tail of the A<sub>2A</sub>R [102,103]. Transmembrane domains of the D<sub>2</sub>R, particularly the fifth transmembrane domain, also appeared to play a role [37]. A comprehensive molecular model of the A<sub>2A</sub>R/D<sub>2</sub>R heteromer was developed [104].

Triplet homologies in A<sub>2A</sub>R and D<sub>2</sub>R (e.g., alanine-alanine-arginine) have been proposed to guide the heteromer partners and to clasp them together [105]. “Pull-down” assays are in vitro methods to identify and determine physical interactions between two proteins. Using such techniques and mass spectrometry, a strong electrostatic interaction was demonstrated between negatively charged motifs (aspartic/phosphorylated serine residues) in the C-terminal tail of the A<sub>2A</sub>R and a positively charged (arginine-rich) epitope in the third intracellular loop of the D<sub>2</sub>R [106]. This electrostatic interaction was shown to possess an amazing stability, comparable to the stability of a covalent bond [107]. The importance of the serine residue in the C-terminal tail of the A<sub>2A</sub>R for A<sub>2A</sub>R-D<sub>2</sub>R receptor–receptor interaction was proven by mutation studies. A point mutation (change of serine 374 to alanine) reduced the formation of A<sub>2A</sub>/D<sub>2</sub> heteromers and the allosteric modulation of D<sub>2</sub>R by A<sub>2A</sub>R agonists and antagonists [108]. Additional mutation of two aspartate residues (401–402 to alanine) in the C-terminal tail of the A<sub>2A</sub>R reduced the heteromer formation even further and completely abolished the allosteric modulation of D<sub>2</sub>R by A<sub>2A</sub> ligands [109]. The importance of transmembrane domains of the D<sub>2</sub>R for heteromer formation was proven by administering synthetic peptides corresponding to the structure of the fourth and fifth transmembrane domain of the D<sub>2</sub>R. Such peptides reduced the ability of A<sub>2A</sub>R and D<sub>2</sub>R to form heteromers [109]. BRET techniques also demonstrated that calmodulin (CaM) interacts with the C-terminal tail of the A<sub>2A</sub>R and provided evidence for the formation of CaM-A<sub>2A</sub>R-D<sub>2</sub>R oligomeric complexes [110].

Japanese investigators created a single-polypeptide chain A<sub>2A</sub>R/D<sub>2</sub>R heteromer by fusing the C-terminus of the A<sub>2A</sub>R to the N-terminus of the D<sub>2</sub>R via a type II transmembrane protein. The resulting synthetic heterodimer showed similar specific binding of A<sub>2A</sub>R and D<sub>2</sub>R ligands and functional coupling to G-proteins as the original wild-type receptors [111].

A very interesting study used BiFC to demonstrate the presence of receptor oligomers in CAD cells, a differentiated neuronal cell model. Prolonged treatment of the cells with the D<sub>2</sub>R agonist quinpirole led to internalization of D<sub>2</sub>R/D<sub>2</sub>R oligomers and A<sub>2A</sub>R/D<sub>2</sub>R heteromers and decreased the relative number of A<sub>2A</sub>R/D<sub>2</sub>R heteromers compared to A<sub>2A</sub>R/A<sub>2A</sub>R oligomers. This effect of quinpirole was reversed by D<sub>2</sub>R antagonists (spiperone, sulpiride), and prolonged treatment of the cells with either a D<sub>2</sub>R antagonist or the A<sub>2A</sub>R agonist MECA resulted in a significant increase of the relative number of A<sub>2A</sub>R/D<sub>2</sub>R heteromers compared to A<sub>2A</sub>R/A<sub>2A</sub>R oligomers. Changes of the heteromer:oligomer ratio were not equivalent to the changes of total A<sub>2A</sub>R and D<sub>2</sub>R numbers in the cells. Thus, drug treatment appeared to modulate G-protein-coupled receptor oligomerization [112].

Investigators from Taiwan demonstrated that both A<sub>2A</sub>R and D<sub>2</sub>R are substrates for sialyltransferases (e.g., St8sia3) in the mouse striatum. If sialylation is reduced (as in St8sia3 knockout mice), a larger fraction of both receptors moves to lipid rafts and a greater number of D<sub>2</sub>R form heteromers with A<sub>2A</sub>R. Thus, sialylation may be a mechanism counteracting heteromer formation and shifting the homomer/heteromer equilibrium in the living brain [113]. Treatment of mice with an A<sub>2A</sub>R antagonist (SCH58261) causes a dose-dependent increase of locomotor activity. This response is much lower in St8sia3-knockout animals than in wild-type mice [113]. On the other hand, treatment of mice with a D<sub>2</sub>R antagonist (L741626) results in a dose-dependent reduction of their locomotor activity, and St8sia3-knockout animals are more sensitive to this effect of a D<sub>2</sub>R antagonist than their wild-type counterparts [113]. Alterations of the A<sub>2A</sub>R/D<sub>2</sub>R homomer/heteromer

equilibrium in the striatum thus appear to be associated with altered responses of the animals to adenosine and dopamine receptor blockade.

D<sub>2</sub>R-agonists can inhibit the 4-aminopyridine-provoked glutamate release in rat striatal astrocytes. Modulation of this inhibition with CGS21680 was shown to depend on the formation of A<sub>2A</sub>R/D<sub>2</sub>R heteromers, whereas the synthetic peptide VLRRRKRVN abolished the effect of CGS21680 [81,114]. VLRRRKRVN binds to the region of the D<sub>2</sub>R that is involved in electrostatic interaction with the A<sub>2A</sub>R and thus blocks the formation of A<sub>2A</sub>R-D<sub>2</sub>R heteromers [106].

Using fluorescent PLA and time-resolved FRET, A<sub>2A</sub>R/D<sub>2</sub>R complexes were detected in the striatum of rodents [94,115–117], monkeys [118], and humans [119]. Such complexes could also be demonstrated and quantified in postmortem brain tissue from patients with Parkinson's disease and healthy control subjects, using AlphaScreen technology [97].

A<sub>2A</sub>R/D<sub>2</sub>R heteromers are now considered to be receptor heterotetramers, consisting of an A<sub>2A</sub>R homodimer and a D<sub>2</sub>R homodimer, each coupled to its own G-protein (G<sub>s</sub> and G<sub>i</sub>, respectively). Adenylate cyclase subtype AC5 also forms part of this multi-protein complex [41,120–125]. The heterotetramer model can explain the apparently paradoxical effects of A<sub>2A</sub>R agonists and antagonists on D<sub>2</sub>R ligand binding in CHO cells that were described in Section 2.3 of this review. Occupancy of the A<sub>2A</sub>R homodimer by either an agonist or an antagonist (at high dose) causes a conformational change in the heterotetramer, resulting in decreased function of the D<sub>2</sub>R protomer in the complex. However, when one of the two adenosine binding sites in the A<sub>2A</sub>R homodimer is occupied by an agonist and the other is simultaneously occupied by an antagonist, the conformational change does not occur [41].

## 6. Pharmacological Consequences of A<sub>2A</sub>/D<sub>2</sub> Heteromer Formation

Receptors can form heteromers if certain basic criteria are met. These include: (a) The individual receptors that can form a heteromeric complex (protomers) must co-localize (i.e., be present in the same membrane domains, at very close distance from each other) and physically interact; (b) formed receptor complexes must exhibit distinct properties which differ from those of the individual, isolated protomers; and (c) chemical compounds that bind selectively to the heteromers should alter the properties or functions of the heteromers [126].

A<sub>2A</sub>R and D<sub>2</sub>R meet all these criteria. Biophysical and molecular biology techniques have demonstrated that these receptors co-localize and physically interact, both in cells and in mammalian tissues (see above, Section 5). Synthetic peptides that interact with the receptor domains involved in heteromer formation affect the electrostatic interactions between the protomers and alter the response of cells to certain drugs (Section 5). In addition, within A<sub>2A</sub>R/D<sub>2</sub>R heterotetramers, various receptor–receptor interactions are possible [125]:

(i) “Canonical interaction”. The agonist-activated G<sub>i</sub>-coupled receptor in the complex (i.e., the D<sub>2</sub>R) will inhibit the activation of adenylate cyclase AC5 by the G<sub>s</sub>-coupled receptor (i.e., the A<sub>2A</sub>R) [36]. The Ras GTPase domain of the subunits of the G<sub>s</sub> and G<sub>i</sub> proteins will interact with the C2 and C1 catalytic domains of adenylate cyclase AC5. The receptor partners in the complex can modulate each other's downstream signaling cascade [36].

(ii) “Allosteric interaction”. Allostery is defined as communication between distant sites in a protein (or protein complex) in which energy associated with ligand binding or conformational change at one site is transferred to other, remote sites of the protein (or protein complex) resulting in changes of the kinetic or conformational properties of these sites. When a ligand binds to one of the receptors in an A<sub>2A</sub>R-D<sub>2</sub>R complex, the conformation of the complex (quaternary structure of the heterotetramer) is altered, resulting in different binding and signaling properties of the other receptor proteins in the complex [41,121,127–129]. When an A<sub>2A</sub>R ligand (either an agonist or an antagonist) binds to the A<sub>2A</sub>R homodimer in the complex, the affinity and signaling efficacy of D<sub>2</sub>R agonists is decreased. On the other hand, when a D<sub>2</sub>R agonist binds to the D<sub>2</sub>R heteromer in the complex, the binding of A<sub>2A</sub>R agonists is suppressed. Such allosteric effects between

A<sub>2A</sub>R and D<sub>2</sub>R have been demonstrated in isolated biomembranes, intact cells, brain slices, and living animals (see above, Section 2).

(iii) “Formation of new modulatory sites”. When different receptor proteins associate to form a heteromer, novel binding sites may be created that are not present in the isolated receptors. Ligands specific to the receptor complex as such may exist [130] and, if discovered, may be used to specifically modulate the complex when it is present [131] (see also Section 11.3).

(iv) “Higher order interaction”. A<sub>2A</sub>R/D<sub>2</sub>R heterotetramers may become part of higher order heteromers, so-called “receptor mosaics” [132]. Such interactions may, for example, involve the metabotropic glutamate receptor 5 (mGluR5) [133,134] or the sigma-1 receptor [117,135–137]. The presence or absence of such additional partners in a higher-order heteromer changes the strength of A<sub>2A</sub>R-D<sub>2</sub>R allosteric interactions and alters the response of the A<sub>2A</sub> and D<sub>2</sub> protomers to adenosine or dopamine. Since an unknown (and variable) number of additional proteins may bind to A<sub>2A</sub>R and D<sub>2</sub>R, the term “heteroreceptor complexes” is used in recent literature rather than A<sub>2A</sub>/D<sub>2</sub>R heterotetramers [138].

Thus, A<sub>2A</sub>R/D<sub>2</sub>R heterotetramers have a distinct pharmacology and distinct functions which differ from those of the individual constituent receptors [139].

## 7. A<sub>2A</sub>/D<sub>2</sub> Interactions and Parkinson’s Disease

Upper motor neurons in the motor regions of the cortex initiate movements, such as continuous postural control, body locomotion, orientation towards sensory stimuli, and orofacial behavior. The activity of lower motor neurons in the spinal cord is coordinated by the upper motor neurons. These lower motor neurons directly or indirectly innervate skeletal muscle fibers [140].

In movement control, there is also a close cooperation of regions in the cortex with the basal ganglia [65,141] (Figure 2). Neurons that belong to the basal ganglia regulate the activity of the upper motor neurons although they do not directly project to them. The major nuclei that comprise the basal ganglia are: The striatum, the globus pallidus (GP), the substantia nigra (SN), and the subthalamic nucleus (STN) [142] (Figure 2). In the rodent brain, the striatum is a single nucleus whereas in primates, it is divided into caudate nucleus and putamen [143]. The basal ganglia receive input from areas of the cerebral cortex and their output is directed towards the thalamus, from where there is a transient excitation back to the motor regions in the cortex (Figure 2). MSNs in the striatum are known to be involved in movement control.

Activation of GABAergic MSNs of the “direct pathway” results in inhibition of the globus pallidus internus (GPi), for GABA is an inhibitory neurotransmitter. Since the GPi is connected to the thalamus via another GABAergic projection, inhibition of the GPi causes disinhibition of the thalamus. Because the thalamus contains excitatory neurons that project to the cortex, activation of the direct pathway results in facilitation of motor activity [140] (Figure 2).

In the “indirect pathway”, GABAergic MSNs project from the striatum to the globus pallidus externus (GPe). A second GABAergic projection runs from the GPe to the subthalamic nucleus (STN) and an excitatory glutamatergic projection connects the STN to the GPi. Activation of the indirect pathway therefore results in disinhibition of the STN and activation of the GPi. This activation of the GPi causes inhibition of the thalamus and reduced activity of the excitatory neurons that run from the thalamus to the cortex. Thus, activation of the indirect pathway results in suppression of motor activity. Although this description of the indirect pathway is probably a gross over-simplification [144], the concept is still widely used as a basis for research and therapy.

Normal movements require a delicate, coordinated balance of activity in the direct and indirect pathways [65]. The healthy brain contains dopaminergic neurons in the substantia nigra pars compacta that project to the striatum. Dopamine from these neurons stimulates the MSNs from the direct pathway via D1R and inhibits the MSNs from the indirect pathway via D2R. Both actions of dopamine facilitate motor activity. Loss of dopaminergic

neurons from the brain, as occurs in Parkinson's disease, will result in a decreased activity of the direct pathway, an increased activity of the indirect pathway and impaired motor control, particularly hypokinesia.

Since loss of dopamine results in overactivity of the indirect pathway, A<sub>2A</sub>R antagonists have been proposed as therapeutic drugs for the treatment of Parkinson's disease [75,145–149]. Such drugs may restore the disturbed balance between the indirect and direct pathways and may increase the effect of endogenous dopamine, L-DOPA and specific D<sub>2</sub> agonists, at least in the early stages of Parkinson's disease [150,151]. A<sub>2A</sub>R antagonists may bind to the A<sub>2A</sub>R protomer in A<sub>2A</sub>R-D<sub>2</sub>R heteromeric complexes and increase the affinity of the D<sub>2</sub>R protomer for dopamine, its coupling to the G-protein and its signaling. In accordance with this hypothesis, perfusion measurements with MRI and pulsed arterial spin labeling have proven that the A<sub>2A</sub>R antagonist tozadenant inhibits (i.e., suppresses the overactivity of) the indirect pathway in the brain of Parkinson's patients [152].

A<sub>2A</sub>R antagonists have been shown to be beneficial in various animal models of PD (e.g., D<sub>2</sub>R knockout mice [153], 6-OHDA-lesioned rats [154,155] and mice [156], rats with pharmacological D<sub>2</sub>R blockade [157], MPTP-treated marmosets [158], and MPTP-treated monkeys [159]). Since locomotor abnormalities in D<sub>2</sub>R knockout mice were rescued by the blockade of A<sub>2A</sub>R, not all actions of A<sub>2A</sub>R are related to the formation of A<sub>2A</sub>R-D<sub>2</sub>R heteromers. Apparently, striatal neuronal activity can also be regulated by A<sub>2A</sub>R via a dopamine D<sub>2</sub>R-independent pathway [152].

Many clinical studies have been performed to explore the effect of adenosine antagonists in Parkinson patients. These studies involved the non-subtype selective adenosine antagonists theophylline [160–162] and caffeine [163], and the A<sub>2A</sub>R-antagonists istradefylline [164–174] and tozadenant [175]. In a single study, theophylline was reported to have no significant effect, probably because group sizes were too small to reach adequate statistical power [162], but in two other studies, the drug caused mild improvement of the objective and subjective symptoms of disability and did not worsen dyskinesia [160,161]. Caffeine temporarily improved freezing of gait in Parkinson's patients with symptoms of total immobility, but not in subjects who suffered from episodes of trembling with incapacity to any further movement [163]. Istradefylline as monotherapy was reported to not improve motor symptoms in early PD [169], but as adjunct therapy was shown to potentiate and prolong the action of L-DOPA. In the presence of istradefylline, lower doses of L-DOPA could be given to the patients and the severity of dyskinesia and resting tremor were reduced [164]. Several studies reported a reduction in "off" time (i.e., the time intervals in which disease symptoms return) when patients were given istradefylline [165–168,170,171] or tozadenant [175] in combination with L-DOPA, and this beneficial effect was not associated with any increase of dyskinesia [168]. Other symptoms of Parkinson's disease, such as daytime sleepiness [172], gait disturbance, freezing of gait, and postural instability [174], were also improved by istradefylline. As a consequence of these positive findings, istradefylline is now a registered drug for treatment of Parkinson's disease, both in Japan [176] and in the U.S. [177].

## 8. A<sub>2A</sub>R-D<sub>2</sub>R Interactions and Schizophrenia

Schizophrenia is thought to be associated with an overactivity of dopamine neurons in the ventral tegmental area of the brain, resulting in increased D<sub>2</sub>R signaling in the nucleus accumbens [178]. As explained above (Sections 2.1 and 3.1), A<sub>2A</sub>R and D<sub>2</sub>R are present not only in the dorsal striatum, but also in the nucleus accumbens. Powerful antagonistic interactions between both receptors occur in this area of the brain and could be detected both in receptor binding studies and in microdialysis experiments. Administration of CGS21680 resulted in a reduced efficacy of dopamine to displace [<sup>125</sup>I]iodosulpiride from D<sub>2</sub>R in the nucleus accumbens. Infusion of the A<sub>2A</sub>R agonist CGS21680 in the nucleus accumbens had the same effect as infusion of the D<sub>2</sub>R antagonist raclopride (i.e., increasing the extracellular levels of GABA in the ipsilateral ventral globus pallidus),

and the stimulation of GABA release by an A<sub>2A</sub>R agonist and a D<sub>2</sub>R antagonist were found to be synergistic [179].

According to several hypotheses, altered levels of extracellular adenosine and adenosine receptors are involved in the pathophysiology of schizophrenia [180–182]. In accordance with such hypotheses, A<sub>2A</sub>R were found to be upregulated in the striatum [183,184] and hippocampus [185] of chronic schizophrenics (although this upregulation could also be a consequence of the antipsychotic treatment that the patients received). A Chinese study reported significant associations between single nucleotide polymorphisms of the A<sub>2A</sub>R gene and schizophrenia in the northern Chinese Han population [186].

Since D<sub>2</sub>R of the ventral striatopallidal neurons are implied in the antipsychotic effects of neuroleptics [187], A<sub>2A</sub>R agonists, either alone or in combination with D<sub>2</sub>R antagonists, have been proposed as potential anti-schizophrenic drugs [179]. The ventral striatopallidal GABA pathway is considered as an anti-reward pathway which is overactivated in schizophrenia due to increased activation of its D<sub>2</sub>R [188]. The antagonistic A<sub>2A</sub>R-D<sub>2</sub>R interactions in the nucleus accumbens, which presumably occur within receptor heteromers, could be exploited to reduce the activity of the D<sub>2</sub>R protomer in the heteroreceptor complex [151,189]. In support of this idea, CGS21680 was shown to act as an atypical antipsychotic drug in rodent models of schizophrenia (phencyclidine, amphetamine) [190] and also in monkeys [191].

Some findings in humans have suggested that stimulation of A<sub>2A</sub>R may be beneficial in the treatment of psychosis. Dipyridamole, a nucleoside transport inhibitor that increases the extracellular levels of adenosine, has been tested as an add-on therapy in the treatment of schizophrenics. Combined treatment with haloperidol and dipyridamole (16 patients) was found to be significantly better than treatment with haloperidol and placebo (14 patients) in reducing positive and general psychopathology symptoms as well as PANSS scores [192]. Administration of allopurinol, a drug which blocks the degradation of purines and increases the levels of adenosine and inosine in the brain, resulted in clinical improvement in two poorly responsive schizophrenic patients [193].

Chronic treatment of rodents with clozapine, an atypical antipsychotic which is more effective than classical antipsychotics in some patients, was found to increase the activity of the enzyme ecto-5'-nucleotidase in the striatum, whereas chronic treatment with haloperidol did not have this effect [194]. These preclinical data suggest that clozapine treatment, in contrast to treatment with typical antipsychotics, is associated with increases of the levels of extracellular adenosine in the brain and with stimulation of A<sub>2A</sub>R.

## 9. A<sub>2A</sub>R-D<sub>2</sub>R Interactions and Treatment of Drug Addiction

According to a common hypothesis of reward-related behavior, the nucleus accumbens exerts tonic inhibitory effects on downstream structures in the brain. When MSNs in the nucleus accumbens are inhibited (e.g., by stimulation of dopamine D<sub>2</sub>R), these downstream structures are excited and an endogenous brake on reward-related behavior is released [195]. Addictive drugs are believed to be rewarding and reinforcing due to their effects on the dopamine reward pathway. They enhance dopamine release as is, for example, the case with nicotine, or they inhibit the reuptake of dopamine as does cocaine, or they act themselves as agonists at D<sub>2</sub>R [196].

Physiologically-relevant rewarding stimuli cause a release of dopamine in the shell of the nucleus accumbens, and this response is subject to habituation when the stimuli are repeatedly administered. Thus, the amount of dopamine that is released by a rewarding non-drug stimulus decreases as a result of repeated exposure to that stimulus. However, the dopamine response in the nucleus accumbens to addictive drugs is not prone to habituation but rather to sensitization, meaning that the amount of dopamine that is released by the drug increases as a result of repeated drug exposure.

Animal experiments in which rats with electrodes implanted in the medial forebrain bundle were trained to rotate a wheel in order to receive a rewarding electrical current

have indicated that A<sub>2A</sub>R agonists elevate current reward thresholds (i.e., inhibit central reward processes) [197]. This observation suggests that A<sub>2A</sub>R modulate reward.

Studies in animal models of cocaine addiction have indicated that stimulation or blockade of A<sub>2A</sub>R has a significant impact on cocaine use. Administration of the non-subtype selective adenosine receptor antagonist caffeine to rats facilitates cocaine self-administration [198,199], whereas an A<sub>2A</sub>R agonist, like CGS21680 or NECA, suppresses the tendency of animals to take cocaine [200,201]. Adenosine receptor agonists appear to suppress cocaine intake by an interaction with A<sub>2A</sub>R in the nucleus accumbens, since microinjections of CGS21680 in the nucleus accumbens, but not in the prefrontal cortex, dose-dependently decrease cocaine self-administration [202]. Microinjections of a synthetic TM5 peptide (which interacts with the fifth transmembrane domain of the A<sub>2A</sub>R and disrupts A<sub>2A</sub>/D<sub>2</sub> heterotetramers), completely counteracted the inhibitory effect of CGS21680 on cocaine intake [203]. In contrast to this striking impact of a TM5 peptide, microinjections of a TM2 peptide (which disrupts A<sub>2A</sub>/A<sub>2A</sub> homodimers but not A<sub>2A</sub>/D<sub>2</sub> heterotetramers) did not counteract the effect of CGS21680 on cocaine self-administration [204]. These results suggest that the beneficial actions of CGS21680 in animal models of cocaine abuse are mediated by the triggering of an allosteric inhibition of D<sub>2</sub> protomer signaling in A<sub>2A</sub>R-D<sub>2</sub>R heteromeric complexes.

The development of cocaine sensitization is enhanced when rats are treated with the A<sub>2A</sub>R antagonist MSX-3 but is reduced when they are treated with the A<sub>2A</sub>R agonist CGS21680 or the D<sub>2</sub>R antagonist raclopride [205]. Administration of CGS21680 (0.25 to 0.5 mg/kg) to rats decreases the acquisition and expression of conditioned place preference induced by cocaine [206] or amphetamines [207].

In the treatment of substance abuse, relapse or drug-seeking behavior after a period of abstinence is a very serious problem. Thus, the finding that CGS21680 dose-dependently inhibits cocaine-induced reinstatement in rats after a period of drug abstinence of at least one week [201,208] is of great interest. On the other hand, A<sub>2A</sub>R antagonists (MSX-3, istradefylline, SCH58261, CGS15943), when administered systemically or by microinjections in the nucleus accumbens, promote cocaine-seeking behavior [202,209–211]. The impact of A<sub>2A</sub>R antagonists appears to be dependent on the question whether postsynaptic or presynaptic A<sub>2A</sub>R are blocked. Istradefylline is a postsynaptic A<sub>2A</sub>R antagonist, whereas SCH442416 blocks mainly presynaptic A<sub>2A</sub>R [212]. Postsynaptic blocking was found to enhance whereas presynaptic blocking reduced reinstatement of cocaine seeking [211,213]. The different antagonist affinities of pre- and postsynaptic A<sub>2A</sub>R may be due to the fact that presynaptic A<sub>2A</sub>R form heteromers with adenosine A<sub>1</sub>R, whereas postsynaptic A<sub>2A</sub>R interact with dopamine D<sub>2</sub>R.

Prolonged cocaine self-administration in rats is associated with a significant upregulation of A<sub>2A</sub>R in the nucleus accumbens [214,215]. After seven days of cocaine withdrawal, A<sub>2A</sub>R numbers in this area of the brain return to normal. This upregulation has been interpreted as a compensatory mechanism to counteract cocaine-induced increases in D<sub>2</sub>R signaling [214]. Mice that were prenatally exposed to cocaine showed an upregulation of D<sub>2</sub>R function and a downregulation of adenosine transporter function, consistent with increased levels of extracellular adenosine and more stimulation of A<sub>2A</sub>R [216]. Thus, cocaine exposure both prenatally and in later life, has direct effects on the dopamine and modulatory adenosine systems.

Cocaine is known to also increase the density of sigma-1R in the nucleus accumbens [217] and to cause trafficking of intracellular sigma-1R to the plasma membrane, where they can interact with D<sub>2</sub>R [135,218]. In fact, cocaine self-administration has been reported to increase the number of A<sub>2A</sub>R-D<sub>2</sub>R and D<sub>2</sub>R-sigma-1R heteromers in the nucleus accumbens shell [117]. These data can also be interpreted as the formation of A<sub>2A</sub>R-D<sub>2</sub>R-sigma-1R heteromeric complexes in response to cocaine, the addition of the sigma-1R to the complex resulting in increased strength of antagonistic A<sub>2A</sub>R-D<sub>2</sub>R interactions [136,137,219,220].

BRET experiments in HEK-293T cells that were co-transfected with A<sub>2A</sub>R and D<sub>2</sub>R demonstrated that cocaine induces a concentration-dependent transient decrease of D<sub>2</sub>R

homodimers and A<sub>2A</sub>R/D<sub>2</sub>R heteromers, but not of A<sub>2A</sub>R homodimers, via a specific interaction with the D<sub>2</sub>R. In co-transfected CHO cells, cocaine was found to cause an increase of the affinity of D<sub>2</sub>R for dopamine and increased coupling of D<sub>2</sub>R to G-proteins by changing the conformation of the receptor protein [221].

Based on such findings (and many others, which are extensively reviewed in [196]), it has been postulated that stimulation of A<sub>2A</sub>R could be a possible strategy to treat drug addiction [201,222–224]. A<sub>2A</sub>R antagonists that preferentially block presynaptic A<sub>2A</sub>R may also offer therapeutic benefits.

## 10. A<sub>2A</sub>R-D<sub>2</sub>R Interactions and Attention Deficit Hyperactivity Disorder

Attention-deficit hyperactivity disorder (ADHD) is a disorder of human behavior that involves dysfunctions of sustained attention, behavioral hyperactivity and impulsivity. ADHD seems to be characterized by reduced functioning of the dopaminergic reward pathway [225,226]. Oral methylphenidate, an inhibitor of noradrenaline and dopamine reuptake, is often prescribed as a therapeutic drug to treat ADHD.

A study that was published in 2000 reported that apart from several genes of the noradrenergic system, polymorphisms of the A<sub>2A</sub>R gene are significantly associated with human ADHD [227]. A later Swedish study confirmed that the A<sub>2A</sub>R gene may indeed be involved in ADHD traits [228]. In rodent models of ADHD, A<sub>2A</sub>R were found to be upregulated in various brain regions [229,230] and adenosine A<sub>2A</sub>R antagonists were shown to have beneficial effects, such as improvement of short-term object-recognition ability, attention and memory function [230,231] and improved development of frontal cortical neurons [232].

A large study involving 1239 human subjects reported an association between the rs2298383 TT genotype of the A<sub>2A</sub>R and anxiety disorders in ADHD. No association with the D<sub>2</sub>R genotype was detected, but a significant, positive gene-gene interaction effect between A<sub>2A</sub>R and D<sub>2</sub>R on the presence of anxiety disorders was noted [233]. This synergistic effect between the A<sub>2A</sub>R and D<sub>2</sub>R genes suggests that A<sub>2A</sub>R-D<sub>2</sub>R heteromers could be explored as a possible target in the treatment of ADHD.

## 11. PET Imaging of Adenosine–Dopamine Interactions

Positron emission tomography (PET) is a minimally invasive imaging technique that allows quantitative assessment of the interaction of radioactive ligands with receptors, enzymes, or transporters in the living brain. Since PET makes it possible to study such interactions repeatedly in experimental animals and humans, this imaging modality may be employed to acquire information about adenosine–dopamine interactions in the healthy human brain, their alterations in disease, and the impact of treatment. Radioligands for adenosine A<sub>2A</sub> and dopamine D<sub>2</sub> receptors are currently available (see Tables 1 and 2, and [234–238] for an overview). However, until now the number of PET studies aiming to demonstrate A<sub>2A</sub>/D<sub>2</sub> interactions have been very limited.

Based on findings acquired with other techniques and reported in the literature, three classes of PET studies concerning adenosine–dopamine interactions appear possible:

**Table 1.** Overview of ligands for positron emission tomography (PET) imaging of A<sub>2A</sub> receptors.

Ligand (Alphabetic Order)	Animal Study		Human Study	Comments
	Animal Model	Reference		
[ <sup>11</sup> C]CSC	Rodent	[239]		
DMPX analogues	Rodent	[240]		
[ <sup>18</sup> F]FDA-PP <sub>1</sub> [ <sup>18</sup> F]FDA-PP <sub>2</sub>				Agonists, no in vivo data [241]
[ <sup>18</sup> F]FESCH (=[ <sup>18</sup> F]MRS5425)	Rodent	[242–245]		
[ <sup>11</sup> C]Istradefylline (=[ <sup>11</sup> C]KW6002)	Rodent	[246,247]	[247]	Extrastriatal off-target binding
[ <sup>11</sup> C]KF17837	Rodent Monkey	[248–250] [251]		High non-specific binding
[ <sup>11</sup> C]TMSX (=[ <sup>11</sup> C]KF18446)	Rodent	[252–257]	[257–270]	
[ <sup>11</sup> C]KF21213	Rodent	[271]		
[ <sup>18</sup> F]MNI-444	Monkey	[272,273]	[274]	
[ <sup>11</sup> C]Preladenant	Rodent Monkey	[275–278] [279]	[280,281]	
[ <sup>11</sup> C]SCH442416	Rodent Monkey	[282,283] [282,284]	[285,286]	

CSC = 8-(3-Chlorostyryl)caffeine, DMPX = 3,7-Dimethyl-1-propargylxanthine, TMSX = [7-methyl-<sup>11</sup>C]-(E)-8-(3,4,5-trimethoxystyryl)-1,3,7-trimethylxanthine. Other compounds are numbered by the producing institutions or pharmaceutical companies.

**Table 2.** Overview of ligands for positron emission tomography (PET) imaging of dopamine D<sub>2/3</sub> receptors.

Ligand (Alphabetic Order)	Rodent, Pig or Cat Study	Monkey or Baboon Study	Human Study	Comments
[ <sup>18</sup> F]Benperidol		[287,288]		
[ <sup>18</sup> F]DMFP	[289–291]		[292,293]	Longer half-life than [ <sup>11</sup> C]raclopride
N-Ethyl-[ <sup>11</sup> C]-et裾lopride		[294]		
[ <sup>11</sup> C]Fallypride		[295]	[296]	
[ <sup>18</sup> F]Fallypride	[291,297–299]	[297,300–304]	[305–311]	High-affinity, visualizes also extrastriatal D <sub>2</sub> R, numerous studies *
[ <sup>18</sup> F]FCP		[312]		
[ <sup>18</sup> F]FEBF	[313]			
[ <sup>18</sup> F]FESP	[314,315]	[314–316]	[314,317]	
[ <sup>11</sup> C]FLB457	[318]	[319,320]	[296,321–328]	High-affinity, visualizes also extrastriatal D <sub>2</sub> R, numerous studies *
[ <sup>11</sup> C]FLB524		[329]	[329]	
5-[ <sup>18</sup> F]FPE		[330]		

**Table 2.** *Cont.*

Ligand (Alpha-Betic Order)	Rodent, Pig or Cat Study	Monkey or Baboon Study	Human Study	Comments
[ <sup>18</sup> F]FPSP	[315,331]	[315,331]	[331]	
[ <sup>18</sup> F]Haloperidol	[332,333]	[334]		Binds also to sigmaR
[ <sup>18</sup> F]MABN	[335]	[336]		
[ <sup>18</sup> F]MBP	[335]	[312,336]		Binds also to rho1
Methyl-[ <sup>11</sup> C]-etclopride		[294]		
[ <sup>11</sup> C]MNPA	[337]	[338,339]	[340]	Agonist ligand
[ <sup>11</sup> C]Nemonapride	[341–343]			Binds also to sigmaR
[ <sup>11</sup> C]NMSP	[341,344,345]		[346–349]	
[ <sup>18</sup> F]NMSP	[350]	[336,350]	[351]	
[ <sup>11</sup> C]NPA	[352,353]	[352,354,355]	[356–358]	Agonist ligand
[ <sup>11</sup> C]PPHT	[359,360]	[359]		Agonist ligand
[ <sup>11</sup> C]Raclopride	[361,362]	[363–367]	[368–379]	Moderate affinity, visualizes mainly striatal D <sub>2</sub> R, numerous studies *
[ <sup>18</sup> F]Spiperone		[287]		
[ <sup>11</sup> C]SV-III-130		[380]		Partial agonist ligand
[ <sup>11</sup> C]ZYY339	[359,360]	[359]		Agonist ligand

\* Only a small selection of the available publications is cited for this radioligand. DMFP = Desmethoxyfallypride, FCP = Fluoroclebopride, FEBF = Fluorethyl-2,3-dihydrobenzofuran, FESP = Fluoroethyl-spiperone, FPE = Fluoropropyl-epidepride, FPSP = Fluoropropyl-spiperone, MABN = 2,3-dimethoxy-N-[9-(4-fluorobenzyl)-9-azabicyclo[3.3.1]nonan-3beta-yl]benzamide, MBP = 2,3-dimethoxy-N-[1-(4-fluorobenzyl)piperidin4yl]benzamide, MNPA = Methoxy-N-n-propylnorapomorphine, NMSP = N-methyl-spiperone, NPA = N-n-propylnorapomorphine, PPHT = (+/-)-2-(N-phenethyl-N-propyl)amino-5-hydroxytetralin. Other compounds are numbered by the producing institutions or pharmaceutical companies.

### 11.1. Pharmacological Challenge Studies

In these studies, subjects are scanned twice with a radioligand for adenosine A<sub>2A</sub>R or dopamine D<sub>2</sub>R, first at baseline (or after administration of a placebo) and then at follow-up, after a pharmacological challenge with a drug that binds to the other receptor system (a dopaminergic drug in the case of A<sub>2A</sub>R imaging, and a purinergic drug in the case of D<sub>2</sub>R imaging). Three investigations that used PET imaging have shown that this experimental set-up allows the detection of adenosine–dopamine interactions in the brain of living mammals.

In the first study [242], the radiotracer [<sup>18</sup>F]MRS5425, an analogue of the A<sub>2A</sub>R antagonist SCH442416, was used to image A<sub>2A</sub>AR in the brain of rats that had been unilaterally lesioned with 6-hydroxydopamine. In this animal model of Parkinson’s disease, the authors observed an increased binding of the tracer in the ipsilateral (lesioned) striatum with respect to the contralateral (healthy) striatum. The increase of [<sup>18</sup>F]MRS5425 in the lesioned hemisphere suggests that loss of dopaminergic neurons can cause upregulation of postsynaptic D<sub>2</sub> and A<sub>2A</sub> receptors, and binding of the PET ligand [<sup>18</sup>F]MRS5425 may be used as a biomarker to monitor Parkinson’s disease. Some animals were subsequently treated with the dopamine D<sub>2</sub>R agonist, quinpirole. A significant (15–20%) decrease of the striatal uptake of [<sup>18</sup>F]MRS5425 was observed after acute administration of quinpirole. The decreased binding of the A<sub>2A</sub>R ligand after a dopaminergic challenge indicates that interactions between D<sub>2</sub>R and A<sub>2A</sub>R can be monitored in living animals with PET [242].

In the second study [381], healthy human subjects with low levels of daily caffeine intake received oral caffeine (300 mg) and the impact of this challenge on the dopaminergic system was assessed by measuring changes of the binding of [<sup>11</sup>C]raclopride to D<sub>2</sub>R in

the brain. A small but significant increase in the binding potential of [<sup>11</sup>C]raclopride was detected in the putamen and ventral striatum (5 to 6%), but not in the caudate nucleus. The rise in the ventral striatum was associated with an increase of alertness caused by caffeine [381]. In an earlier study, which involved administration of 200 mg of oral caffeine to eight human subjects with higher levels of daily caffeine intake, a trend towards increased [<sup>11</sup>C]raclopride binding in the ventral striatum was also noted, but this did not reach statistical significance [382].

In the third study (which was performed in our own institution), anesthetized healthy rats received either the A<sub>2A</sub>R agonist CGS21680 (1 mg/kg, i.p.), the A<sub>2A</sub>R antagonist istradefylline (1 mg/kg, i.p.) or vehicle (saline) and the impact of these challenges on the dopaminergic system was assessed by PET imaging, using full kinetic modeling of the cerebral uptake of the radioligand [<sup>11</sup>C]raclopride. Significant decreases of [<sup>11</sup>C]raclopride binding potential were detected, which were strong (>50%) after intraperitoneal administration of CGS21680 and moderate (30%) after administration of istradefylline [383].

However, these studies also highlighted the complexity of interactions in the living brain and difficulties in pinpointing the exact mechanisms underlying the observed changes. Altered binding potentials in PET imaging may indicate: (i) An altered size of the total receptor population (i.e., altered expression of the receptor gene). (ii) An altered affinity of existing receptors for the radioligand (which may be due to allosteric receptor-receptor interactions within heteromeric complexes). Both A<sub>2A</sub>R agonists (like CGS21680) and A<sub>2A</sub>R antagonists (like istradefylline) can allosterically decrease the affinity of the D<sub>2</sub>R protomer for agonists and antagonists [41,129]. (iii) Increases or decreases of the fraction of internalized receptors (since, in most cases only receptors on the cell surface will bind the radioligand). The adenosine A<sub>2A</sub>R agonist CGS21680 promotes the recruitment of β-arrestin2 to the D<sub>2</sub>R protomers in an A<sub>2A</sub>/D<sub>2</sub> heteromer complex and causes subsequent co-internalization of A<sub>2A</sub> and D<sub>2</sub> receptors [84,88], a process in which caveolin-1 is involved [87]. (iv) Increases or decreases of the extracellular concentration of the endogenous neurotransmitter or neuromodulator (which competes with the radioligand for binding to a limited number of receptor sites). Selective adenosine A<sub>2A</sub>R antagonists may increase the release of dopamine [384] and may also inhibit the enzyme monoamine oxidase B and thus raise the levels of extracellular dopamine [385]. The first mechanism (altered gene expression) is unlikely as an explanation for the observed changes of [<sup>11</sup>C]raclopride binding potential, since the PET studies employed an acute drug challenge and measured radioligand binding shortly after the challenge. The increased binding potential of [<sup>11</sup>C]raclopride that was noticed in the ventral striatum after administration of caffeine cannot reflect a decrease of extracellular dopamine, since increased alertness was noticed under these conditions. Increased alertness is normally related to augmented release of dopamine in the striatum, whereas reductions of extracellular dopamine are accompanied by increased tiredness and sleepiness [381]. Thus, the increase of [<sup>11</sup>C] raclopride binding after caffeine intake may reflect an altered affinity of D<sub>2</sub>R for the radioligand or a reduced internalization of D<sub>2</sub>R in the presence of caffeine.

The PET studies mentioned above [242,381–383] indicate that adenosine–dopamine receptor interactions can be visualized and quantified in the brain of living mammals, but various mechanisms or a combination of mechanisms may be involved and may cause the observed changes.

Other PET studies have indicated that antagonistic effects between adenosine A<sub>2A</sub> and dopamine D<sub>2</sub> receptors at the MSNs of the striatum occur at physiological levels of receptor occupancy in the living brain. The D<sub>2</sub>R antagonist haloperidol is widely used as an antipsychotic, but can induce extrapyramidal symptoms (i.e., movement disorders, such as catalepsy (rigidity, muscle stiffness, fixed posture)). In non-human primates, the duration of the cataleptic posture induced by haloperidol (0.03 mg/kg, i.m.) was reduced when animals were treated with the A<sub>2A</sub>R antagonist ASP5854 (0.1 mg/kg, oral). A PET study with the A<sub>2A</sub>R ligand [<sup>11</sup>C]SCH442416 showed that the anti-cataleptic effect of ASP5854 (0.1 mg/kg, oral) was reached at an A<sub>2A</sub>R occupancy of 85% [284].

In Parkinson's patients treated with dopaminergic medication (and rodents with 6-OHDA induced hemiparkinsonism), A<sub>2A</sub>R were found to be upregulated if dyskinesia was present, but not when dyskinesia was absent [56,262,278,285]. This finding indicates that adenosine–dopamine interactions are clinically relevant and A<sub>2A</sub>R antagonists may be applied as therapeutic drugs [386].


### 11.2. Studies with Bivalent Radioligands

As discussed above, PET imaging with a suitable radioligand for adenosine A<sub>2A</sub> or dopamine D<sub>2</sub> receptors may be used to gain information on adenosine–dopamine interactions. Changes of radioligand binding to one protomer after a pharmacological challenge to the other protomer can be monitored with PET. The magnitude of these changes may be proportional to the relative abundance of A<sub>2A</sub>/D<sub>2</sub>R heterotetramers and/or the strength of the A<sub>2A</sub>R–D<sub>2</sub>R interaction, which could be altered by disease or after a successful treatment.

Another approach to visualize and quantify A<sub>2A</sub>R/D<sub>2</sub>R heterotetramers is the use of radiolabeled bivalent ligands for PET imaging, so-called “bivalent probes”. In early attempts to target receptor homodimers, the orthosteric sites of two homodimer partners were bridged by a ligand consisting of two identical pharmacophores connected by a short linker. A similar approach could be tried to target heteromers. The two receptor partners in a heteromer may be bridged by a ligand consisting of two different pharmacophores (appropriately designed for each individual heteromer partner) connected via a short spacer. In this way, the ligand can bind simultaneously to two GPCR receptors if these receptors are closely together (i.e., within a receptor heteromer). A successful bivalent ligand will bind more avidly (with 10–100 fold greater affinity) to the appropriate receptor heteromer than to the isolated receptor monomers or homodimers.

Some experience with bivalent ligands has been acquired by pharmacochemists. In the past, virtually all therapeutic drugs were designed to target a single protein. The discovery of heteromeric receptors has led to a new interest in the development of mixed action drugs for combination therapies, or drugs which selectively bind to receptor heteromers [138,387]. A heterobivalent ligand combining D<sub>2</sub>R agonism with A<sub>2A</sub>R antagonism could be an effective antiparkinsonian drug and might also be radiolabeled for PET imaging of A<sub>2A</sub>R/D<sub>2</sub>R heteromers [388]. The potential of such mixed-actions drugs has been demonstrated in the opioid system, where successful bivalent analgesics combining μ-agonism with δ-antagonism have been developed [389,390].

The synthesis of A<sub>2A</sub> antagonist-D<sub>2</sub> agonist heterobivalent drugs was first reported in 2009 [391]. The spacer in these drugs was based on trifunctional amino acids that were combined with PEG-polyamide unit repeats. Various bivalent ligands were constituted by connecting the A<sub>2A</sub>R antagonist 8-(p-carboxymethoxy)phenyl-1,3 dipropylxanthine (XCC) and the D<sub>2</sub>R agonist (+/-)-2-(N-phenethyl-N-propyl)amino-5-hydroxytetralin (PPHT-NH<sub>2</sub>) via a Lys-Lys-[PEG/polyamide]<sub>n</sub>-Lys-Glu (n = 0–7) linker (Figure 3). In competition radioligand binding experiments using striatal membranes, it was shown that these bivalent ligands could displace specific A<sub>2A</sub>R and D<sub>2</sub>R radioligand binding. The bivalent compounds displaced the specific monovalent ligands [<sup>3</sup>H]-ZM241385 and [<sup>3</sup>H]-YM09151-2 only when both A<sub>2A</sub> and D<sub>2</sub> receptors were expressed in cells. Such displacement could also be observed in striatal tissue, indicating the presence of A<sub>2A</sub>/D<sub>2</sub> receptor heteromers. This suggests that heterobivalent ligands could potentially serve as PET probes for A<sub>2A</sub>/D<sub>2</sub> receptor heteromers in native tissues and as pharmacological tools to investigate the properties of A<sub>2A</sub>R-D<sub>2</sub>R heterotetramers. They could also pave the way for the design of heteromer-selective drugs for the treatment of Parkinson's disease.




**Figure 3.** Chemical structures of the A<sub>2A</sub> antagonist XCC (**A**) and the D<sub>2</sub> agonist PPHT-NH<sub>2</sub> (**B**). By attaching a linker to the atomic positions indicated by the arrowheads, a bivalent ligand for A<sub>2A</sub>AR/D<sub>2</sub>R heteromers can be created [392].

The distance between the ligand pharmacophores should correspond to the distance of the two binding sites in the receptor heteromer. Thus, the pharmacophore units should be connected with a linker of the appropriate length. Docking experiments predicted that a linker of 26 atoms allows two pharmacophore moieties to bind to the A<sub>2A</sub>/D<sub>2</sub> heteromer complex. The points of attachment of the linker to the pharmacophore units are another crucial aspect of bivalent ligand design. During the development of a bivalent ligand with A<sub>2A</sub>R antagonist and D<sub>2</sub>R agonist action, it was found that the [COOH] position of the adenosine antagonist XCC and the N-terminal position of dopamine D<sub>2</sub>R agonist were most suitable for the attachment of a linker [391]. Another difficulty in the design of bivalent ligands is the fact that the attachment of a linker and a spacer generally results in a reduction in affinity of each pharmacophore for its target. Even if the lead compounds of bivalent ligands have target affinities in the low nanomolar range, the fused ligand with its spacer may have a too low affinity for successful PET imaging [392]. Lead compounds with very high affinities may be required for the design of a successful bivalent probe for PET imaging.


Bivalent ligands also face challenges concerning CNS uptake, metabolism, and excretion. CNS drugs and radioligands for CNS imaging must cross the blood–brain barrier (BBB) in order to be effective. The ability of compounds to penetrate the BBB by passive diffusion is guided by a number of molecular properties: Topological polar surface area (tPSA) should be <90 Å, hydrogen bond donors (HBDs) < 3, cLog P = 2–5, and the molecular weight <450 Da [393]. Although some properties of bivalent and heterobivalent drugs may not satisfy these criteria, they may still cross the BBB. The serotonin receptor agonist sumatriptan has a molecular weight of 721 Da (i.e., greater than 450 Da), yet it crosses the BBB and is a successful therapeutic drug [392]. Thus, although the design of a bivalent A<sub>2A</sub>AR/D<sub>2</sub>R ligand remains a serious challenge, the problems may be overcome.

Another approach to development of bivalent A<sub>2A</sub>R-D<sub>2</sub>R ligands has been described in the literature [394]. This approach resulted in a new compound, DP-L-A2AANT, that was prepared by amide conjugation of dopamine (DP) to an A<sub>2A</sub> antagonist (A2AANT) via a succinic spacer (L) (Figure 4). The spacer was bound to the amine group of A2AANT. The fusion compound showed a high A<sub>2A</sub>R affinity ( $K_i$  2.07 ± 0.23 nM in rat striatum). Although the drug did not exhibit a high affinity towards D<sub>2</sub>R and cannot be considered as a suitable candidate for labeling with a positron emitter for PET imaging, it could be a lead compound for the development of antiparkinsonian drugs and PET tracers. Administration of DP-L-A2AANT led to a release of L-A2AANT and dopamine that could be detected in heparinized human whole blood after one and two hours, and DP-L could be detected after 8 h. The bivalent drug may allow prolonged delivery of small amounts of dopamine which are associated with little neuronal toxicity and limited side effects compared to conventional dopaminergic treatments, especially since succinic acid is known to have low toxicity in humans [394].



**Figure 4.** Conjugation of dopamine with the A<sub>2A</sub> antagonist 7-amino-5-(aminomethyl)-cyclohexyl 2-methyl-amino-2-(2-furyl)-1,2,4-triazolo[1,5-*a*]-1,3,5-triazine trifluoroacetate via a succinate spacer, to obtain the prodrug DP-L-A<sub>2A</sub>ANT, a bivalent ligand [394].

The potent A<sub>2A</sub>R antagonist ZM 241385 and the D<sub>2</sub>R agonist ropinirole are considered to have favorable properties for the design of bivalent therapeutic drugs. These classic ligands were used in 2015 to synthesize a novel series of compounds [395] (Figure 5). A cyclic linker between the A<sub>2A</sub> and D<sub>2</sub> pharmacophores could be used to increase the structural rigidity of a bivalent ligand. Generally, triazine-linker based members of the family showed a 4-fold decrease in A<sub>2A</sub> inhibitory potency compared to the parent A<sub>2A</sub>R antagonist and maintained their functional potency towards the D<sub>2</sub>R. Non-cyclic dual acting ligands showed a 28- to 54-fold reduction in their A<sub>2A</sub>R inhibitory potency. Many of the developed compounds passed preliminary BBB permeability tests. However, the *in vivo* brain uptake kinetics of these dual acting ligands should still be determined [395].



**Figure 5.** Conjugation of the adenosine A<sub>2A</sub>R antagonist ZM241385 (**A**) and the dopamine D<sub>2</sub> agonist ropinirole (**B**) to form a heterobivalent ligand (**C**). Dual action drugs can be prepared by using cyclic (**D**) or non-cyclic (**E**) spacers, and the latter may contain an ionizable tertiary amine [395].

In conclusion, although the development of bivalent radioligands for PET imaging of A<sub>2A</sub>R/D<sub>2</sub>R heterotetramers will be a major challenge, the design of such compounds may prove to be possible.

### 11.3. Studies with Radiolabeled Heteromer-Specific Allosteric Modulators

Experiments in which the impact of homocysteine on A<sub>2A</sub>R-D<sub>2</sub>R heteromers was examined have suggested that heteromer-specific allosteric modulators may exist [131,220,396,397]. In CHO cells that express both A<sub>2A</sub>R and D<sub>2</sub>R, homocysteine reduces the internalization of A<sub>2A</sub>R-D<sub>2</sub>R complexes after stimulation of the D<sub>2</sub>R [396]. Homocysteine was shown to form a non-covalent complex with an arginine-rich epitope involved in heteromer formation but did not disrupt or prevent the formation of A<sub>2A</sub>R-D<sub>2</sub>R heteromers in co-transfected HEK cells [396]. In striatal astrocytes, homocysteine reduces the D<sub>2</sub>R-mediated inhibition of glutamate release but does not affect the A<sub>2A</sub>R-mediated antagonism of this D<sub>2</sub>R effect [397]. These data have been interpreted as evidence that homocysteine binds to A<sub>2A</sub>R-D<sub>2</sub>R heteromers and modulates the allosteric energy transmission between A<sub>2A</sub>R and D<sub>2</sub>R in a heteromer complex [220].

Labeling homocysteine with a positron emitter is definitely not a viable strategy to develop a heteromer-specific radioligand for PET imaging, but if other substances can be identified which bind to specific pockets in the A<sub>2A</sub>R-D<sub>2</sub>R interface within a heteromer without disrupting or preventing heteromer formation, such substances could be used as lead compounds to develop heteromer-specific imaging agents or therapeutic drugs.

### 11.4. Other Opportunities for PET Imaging

PET may not only be used to visualize and quantify A<sub>2A</sub>R-D<sub>2</sub>R heterotetramers or to determine the strength of receptor–receptor interactions within heteromeric complexes, but also to assess the physiological consequences of pharmacological targeting of A<sub>2A</sub>R-D<sub>2</sub>R heteromers. Regional changes of cerebral glucose metabolism after administration of a heterobivalent drug may be measured with the PET tracer [<sup>18</sup>F]fluorodeoxyglucose (FDG), and regional changes of cerebral perfusion can be measured with flow tracers and PET or single photon emission computed tomography (SPECT), or with functional magnetic resonance imaging (fMRI). In animal models of Parkinson’s disease, A<sub>2A</sub>R antagonists have been found to not only improve motor function, but also to be neuroprotective [398–403]. The neuroprotective actions of such drugs, drug combinations or heterobivalent drugs may be assessed with PET imaging (e.g., by visualizing dopaminergic nerve endings with dopamine transporter ligands such as [<sup>11</sup>C]PE2I or [<sup>18</sup>F]-FE-PE2I, or by visualizing neuroinflammation with radiolabeled TSPO ligands, in longitudinal studies. An exciting final possibility is to compare changes of the binding of a radioligand for presynaptic A<sub>2A</sub> receptors, such as [<sup>11</sup>C]SCH442416, with changes of the binding of a radioligand for postsynaptic A<sub>2A</sub> receptors in disease models (although [<sup>11</sup>C]KW6002 = istradefylline is probably not ideal for this purpose).

## 12. Conclusions

Understanding the complexities of A<sub>2A</sub>/D<sub>2</sub> interactions is important in unravelling basal ganglia physiology. Initial observation of antagonistic interactions between adenosine and dopamine in membrane preparations, intact cells and living animals was followed by proof of direct interactions between A<sub>2A</sub> and D<sub>2</sub> receptors supplied by various biophysical techniques and demonstration of the molecular mechanisms involved in these protein–protein interactions. Such biochemical and biophysical findings facilitated further studies of basal ganglia disorders.

Several neurodegenerative and neuropsychiatric disorders are associated with a dysregulation of corticostriatal and nigrostriatal afferents that leads to aberrant neurotransmission. Membrane interactions between adenosine A<sub>2A</sub>R and dopamine D<sub>2</sub>R are an important aspect of striatal function and appear to be altered in Parkinson’s disease, schizophrenia, substance abuse, and ADHD. A<sub>2A</sub>R and D<sub>2</sub>R have been proposed as therapeutic targets.

Decreasing A<sub>2A</sub> signaling by selective A<sub>2A</sub> antagonists may result in a recovery of GPe activity in Parkinson's disease, thereby reinstating the thalamocortical motor stimulatory activity. A<sub>2A</sub> agonists, either alone or in combination with D<sub>2</sub>R antagonists, have been proposed for the treatment of schizophrenia. Such combination treatments reduce overactivity of the D<sub>2</sub>R protomer in the A<sub>2A</sub>/D<sub>2</sub> receptor complex. Drugs that are known to increase the levels of extracellular adenosine (such as nucleoside transport inhibitors, inhibitors of purine degradation and antipsychotics increasing the activity of the enzyme 5'-nucleotidase) may be used as a replacement for A<sub>2A</sub> agonists in A<sub>2A</sub>R-D<sub>2</sub>R based combination therapies. Stimulation of A<sub>2A</sub>R has also been postulated as a possible strategy to treat substance abuse, in particular addiction to cocaine. A<sub>2A</sub>R antagonists, on the other hand, could have a beneficial effect in combination therapies for ADHD. Thus, heteromers of A<sub>2A</sub>R and D<sub>2</sub>R are potential targets for the treatment of several human disorders.

PET imaging may provide significant in vivo information that leads to greater understanding of the role of A<sub>2A</sub>R/D<sub>2</sub>R heteromers in the physiology of the healthy and diseased brain. PET studies with radioligands for A<sub>2A</sub>R or D<sub>2</sub>R before and after a pharmacological challenge to the other protomer in the A<sub>2A</sub>R-D<sub>2</sub>R complex may be used to assess the strength of A<sub>2A</sub>/D<sub>2</sub> receptor interactions. Bivalent ligands that bind simultaneously to A<sub>2A</sub> and D<sub>2</sub> receptors if the receptor proteins are at close proximity may be used as molecular probes to assess the regional abundance of A<sub>2A</sub>R-D<sub>2</sub>R heteromers. The physiological consequences of pharmacological targeting of A<sub>2A</sub>R-D<sub>2</sub>R heteromers in patients can be assessed by PET imaging with tracers visualizing cerebral energy metabolism, cerebral perfusion, neuroinflammation, or dopamine transporter expression.

**Author Contributions:** A.v.W., K.P. and E.F.J.d.V. were involved in the drafting of the manuscript. P.H.E. and R.A.J.O.D. read the initial drafts and provided helpful suggestions for improvement. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research received no external funding.

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** Not applicable.

**Conflicts of Interest:** The authors declare no conflict of interest.

## Abbreviations

A2AR	Adenosine A2A receptor
ADHD	Attention deficit hyperactivity disorder
AlphaScreen	Amplified luminescent proximity homogeneous assay screen (based on luminescent oxygen channeling immunoassay)
BBB	Blood-brain barrier
BHT-920	5,6,7,8-Tetrahydro-6-(2-propen-1-yl)-4H-thiazolo[4,5-d]azepin-2-amine dihydrochloride
BiFC	Bimolecular fluorescence complementation
BRET	Bioluminescence resonance energy transfer
CGS21680	2-[p-(2-carboxyethyl)phenethylamino]-5'-N-ethylcarboxamido-adenosine
CNS	Central nervous system
CSC	8-(3-Chlorostyryl)caffeine
D2R	Dopamine D2 receptor
DMFP	Desmethoxyfallypride
DMPX	3,7-Dimethyl-1-propargylxanthine
FCP	Fluoroclebopride
FEBF	Fluoreethyl-2,3-dihydrobenzofuran
FESP	Fluoroethyl-spiperone
FPE	Fluoropropyl-epidepride

FPSP	Fluoropropyl-spiperone
FRET	Fluorescence resonance energy transfer
MABN	2,3-dimethoxy-N-[9-(4-fluorobenzyl)-9-azabicyclo[3.3.1]nonan-3beta-yl]benzamide
MBP	2,3-dimethoxy-N-[1-(4-fluorobenzyl)piperidin4yl]benzamide
MNPA	Methoxy-N-n-propylnorapomorphine
MSN	Medium spiny neuron
MECA	5'-(N-methyl)carboxamido-adenosine
NECA	5'-(N-ethyl)carboxamido-adenosine
NMSP	N-methyl-spiperone
NPA	N-n-propylnorapomorphine
PANNS	Positive and Negative Syndrome Scale
PIA	N6-R-phenylisopropyladenosine
PPHT	(+/-)-2-(N-phenethyl-N-propyl)amino-5-hydroxytetralin
TMSX	[7-methyl- <sup>11</sup> C]-{(E)-8-(3,4,5-trimethoxystyryl)-1,3,7-trimethylxanthine}
XCC	8-(p-carboxymethoxy)phenyl-1,3 dipropylxanthine

## References

- Latini, S.; Corsi, C.; Pedata, F.; Pepeu, G. The Source of Brain Adenosine Outflow during Ischemia and Electrical Stimulation. *Neurochem. Int.* **1996**, *28*, 113–118. [[CrossRef](#)]
- Ballarín, M.; Fredholm, B.B.; Ambrosio, S.; Mahy, N. Extracellular Levels of Adenosine and Its Metabolites in the Striatum of Awake Rats: Inhibition of Uptake and Metabolism. *Acta Physiol. Scand.* **1991**, *142*, 97–103. [[CrossRef](#)] [[PubMed](#)]
- Pazzaglia, M.; Corsi, C.; Fratti, S.; Pedata, F.; Pepeu, G. Regulation of Extracellular Adenosine Levels in the Striatum of Aging Rats. *Brain Res.* **1995**, *684*, 103–106. [[CrossRef](#)]
- Murillo-Rodriguez, E.; Liu, M.; Blanco-Centurion, C.; Shiromani, P.J. Effects of Hypocretin (Orexin) Neuronal Loss on Sleep and Extracellular Adenosine Levels in the Rat Basal Forebrain. *Eur. J. Neurosci.* **2008**, *28*, 1191–1198. [[CrossRef](#)] [[PubMed](#)]
- Nelson, A.M.; Battersby, A.S.; Baghdoyan, H.A.; Lydic, R. Opioid-Induced Decreases in Rat Brain Adenosine Levels Are Reversed by Inhibiting Adenosine Deaminase. *Anesthesiology* **2009**, *111*, 1327–1333. [[CrossRef](#)] [[PubMed](#)]
- Sharma, R.; Engemann, S.C.; Sahota, P.; Thakkar, M.M. Effects of Ethanol on Extracellular Levels of Adenosine in the Basal Forebrain: An in Vivo Microdialysis Study in Freely Behaving Rats. *Alcohol. Clin. Exp. Res.* **2010**, *34*, 813–818. [[CrossRef](#)] [[PubMed](#)]
- Miranda, M.F.; Hamani, C.; De Almeida, A.C.; Amorim, B.O.; Macedo, C.E.; Fernandes, M.J.; Nobrega, J.N.; Aarão, M.C.; Madureira, A.P.; Rodrigues, A.M.; et al. Role of Adenosine in the Antiepileptic Effects of Deep Brain Stimulation. *Front. Cell. Neurosci.* **2014**, *8*, 312. [[CrossRef](#)]
- Collis, M.G.; Hourani, S.M. Adenosine Receptor Subtypes. *Trends Pharmacol. Sci.* **1993**, *14*, 360–366. [[CrossRef](#)]
- Fredholm, B.B.; Abbracchio, M.P.; Burnstock, G.; Daly, J.W.; Harden, T.K.; Jacobson, K.A.; Leff, P.; Williams, M. Nomenclature and Classification of Purinoceptors. *Pharmacol. Rev.* **1994**, *46*, 143–156.
- Palmer, T.M.; Stiles, G.L. Adenosine Receptors. *Neuropharmacology* **1995**, *34*, 683–694. [[CrossRef](#)]
- Dunwiddie, T.V.; Masino, S.A. The Role and Regulation of Adenosine in the Central Nervous System. *Annu. Rev. Neurosci.* **2001**, *24*, 31–55. [[CrossRef](#)] [[PubMed](#)]
- Fredholm, B.B.; IJzerman, A.P.; Jacobson, K.A.; Klotz, K.N.; Linden, J. International Union of Pharmacology. XXV. Nomenclature and Classification of Adenosine Receptors. *Pharmacol. Rev.* **2001**, *53*, 527–552. [[PubMed](#)]
- Fredholm, B.B.; IJzerman, A.P.; Jacobson, K.A.; Linden, J.; Müller, C.E. International Union of Basic and Clinical Pharmacology. LXXXI. Nomenclature and Classification of Adenosine Receptors—an Update. *Pharmacol. Rev.* **2011**, *63*, 1–34. [[CrossRef](#)] [[PubMed](#)]
- Fuxe, K.; Ungerstedt, U. Action of Caffeine and Theophyllamine on Supersensitive Dopamine Receptors: Considerable Enhancement of Receptor Response to Treatment with DOPA and Dopamine Receptor Agonists. *Med. Biol.* **1974**, *52*, 48–54. [[PubMed](#)]
- Ferré, S.; Herrera-Marschitz, M.; Grabowska-Andén, M.; Ungerstedt, U.; Casas, M.; Andén, N.E. Postsynaptic Dopamine/Adenosine Interaction: I. Adenosine Analogues Inhibit Dopamine D2-Mediated Behaviour in Short-Term Reserpinized Mice. *Eur. J. Pharmacol.* **1991**, *192*, 25–30. [[CrossRef](#)]
- Ferré, S.; Herrera-Marschitz, M.; Grabowska-Andén, M.; Casas, M.; Ungerstedt, U.; Andén, N.E. Postsynaptic Dopamine/Adenosine Interaction: II. Postsynaptic Dopamine Agonism and Adenosine Antagonism of Methylxanthines in Short-Term Reserpinized Mice. *Eur. J. Pharmacol.* **1991**, *192*, 31–37. [[CrossRef](#)]
- Ferré, S.; Rubio, A.; Fuxe, K. Stimulation of Adenosine A2 Receptors Induces Catalepsy. *Neurosci. Lett.* **1991**, *130*, 162–164. [[CrossRef](#)]
- Trevitt, J.; Vallance, C.; Harris, A.; Goode, T. Adenosine Antagonists Reverse the Cataleptic Effects of Haloperidol: Implications for the Treatment of Parkinson’s Disease. *Pharmacol. Biochem. Behav.* **2009**, *92*, 521–527. [[CrossRef](#)]
- El Yacoubi, M.; Ledent, C.; Parmentier, M.; Costentin, J.; Vaugeois, J.M. Adenosine A2A Receptor Knockout Mice Are Partially Protected Against Drug-Induced Catalepsy. *Neuroreport* **2001**, *12*, 983–986. [[CrossRef](#)]

20. Varty, G.B.; Hodgson, R.A.; Pond, A.J.; Grzelak, M.E.; Parker, E.M.; Hunter, J.C. The Effects of Adenosine A2A Receptor Antagonists on Haloperidol-Induced Movement Disorders in Primates. *Psychopharmacology* **2008**, *200*, 393–401. [CrossRef]
21. Alexander, S.P.; Reddington, M. The Cellular Localization of Adenosine Receptors in Rat Neostriatum. *Neuroscience* **1989**, *28*, 645–651. [CrossRef]
22. Rebola, N.; Canas, P.M.; Oliveira, C.R.; Cunha, R.A. Different Synaptic and Subsynaptic Localization of Adenosine A2A Receptors in the Hippocampus and Striatum of the Rat. *Neuroscience* **2005**, *132*, 893–903. [CrossRef]
23. Hettinger, B.D.; Lee, A.; Linden, J.; Rosin, D.L. Ultrastructural Localization of Adenosine A2A Receptors Suggests Multiple Cellular Sites for Modulation of GABAergic Neurons in Rat Striatum. *J. Comp. Neurol.* **2001**, *431*, 331–346. [CrossRef]
24. Rosin, D.L.; Hettinger, B.D.; Lee, A.; Linden, J. Anatomy of Adenosine A2A Receptors in Brain: Morphological Substrates for Integration of Striatal Function. *Neurology* **2003**, *61*, S12–S18. [CrossRef] [PubMed]
25. Font, L.; Mingote, S.; Farrar, A.M.; Pereira, M.; Worden, L.; Stopper, C.; Port, R.G.; Salamone, J.D. Intra-Accumbens Injections of the Adenosine A2A Agonist CGS 21680 Affect Effort-Related Choice Behavior in Rats. *Psychopharmacology* **2008**, *199*, 515–526. [CrossRef]
26. Worden, L.T.; Shahriari, M.; Farrar, A.M.; Sink, K.S.; Hockemeyer, J.; Müller, C.E.; Salamone, J.D. The Adenosine A2A Antagonist MSX-3 Reverses the Effort-Related Effects of Dopamine Blockade: Differential Interaction With D1 and D2 Family Antagonists. *Psychopharmacology* **2009**, *203*, 489–499. [CrossRef] [PubMed]
27. Salamone, J.D.; Farrar, A.M.; Font, L.; Patel, V.; Schlar, D.E.; Nunes, E.J.; Collins, L.E.; Sager, T.N. Differential Actions of Adenosine A1 and A2A Antagonists on the Effort-Related Effects of Dopamine D2 Antagonism. *Behav. Brain Res.* **2009**, *201*, 216–222. [CrossRef] [PubMed]
28. Ferré, S.; von Euler, G.; Johansson, B.; Fredholm, B.B.; Fuxe, K. Stimulation of High-Affinity Adenosine A2 Receptors Decreases the Affinity of Dopamine D2 Receptors in Rat Striatal Membranes. *Proc. Natl. Acad. Sci. USA* **1991**, *88*, 7238–7241. [CrossRef]
29. Ferré, S.; Fuxe, K. Dopamine Denervation Leads to an Increase in the Intramembrane Interaction between Adenosine A2 and Dopamine D2 Receptors in the Neostriatum. *Brain Res.* **1992**, *594*, 124–130. [CrossRef]
30. Ferré, S.; Snaprud, P.; Fuxe, K. Opposing Actions of an Adenosine A2 Receptor Agonist and a GTP Analogue on the Regulation of Dopamine D2 Receptors in Rat Neostratal Membranes. *Eur. J. Pharmacol.* **1993**, *244*, 311–315. [CrossRef]
31. Agnati, L.F.; Fuxe, K.; Benfenati, F.; von Euler, G.; Fredholm, B. Intramembrane Receptor-Receptor Interactions: Integration of Signal Transduction Pathways in the Nervous System. *Neurochem. Int.* **1993**, *22*, 213–222. [CrossRef]
32. Fuxe, K.; Ferré, S.; Zoli, M.; Agnati, L.F. Integrated Events in Central Dopamine Transmission As Analyzed at Multiple Levels. Evidence for Intramembrane Adenosine A2A/Dopamine D2 and Adenosine A1/Dopamine D1 Receptor Interactions in the Basal Ganglia. *Brain Res. Rev.* **1998**, *26*, 258–273. [CrossRef]
33. Yang, S.N.; Dasgupta, S.; Lledo, P.M.; Vincent, J.D.; Fuxe, K. Reduction of Dopamine D2 Receptor Transduction by Activation of Adenosine A2a Receptors in Stably A2a/D2 (Long-Form) Receptor Co-Transfected Mouse Fibroblast Cell Lines: Studies on Intracellular Calcium Levels. *Neuroscience* **1995**, *68*, 729–736. [CrossRef]
34. Salim, H.; Ferré, S.; Dalal, A.; Peterfreund, R.A.; Fuxe, K.; Vincent, J.D.; Lledo, P.M. Activation of Adenosine A1 and A2A Receptors Modulates Dopamine D2 Receptor-Induced Responses in Stably Transfected Human Neuroblastoma Cells. *J. Neurochem.* **2000**, *74*, 432–439. [CrossRef] [PubMed]
35. Dasgupta, S.; Ferré, S.; Kull, B.; Hedlund, P.B.; Finnman, U.B.; Ahlberg, S.; Arenas, E.; Fredholm, B.B.; Fuxe, K. Adenosine A2A Receptors Modulate the Binding Characteristics of Dopamine D2 Receptors in Stably Cotransfected Fibroblast Cells. *Eur. J. Pharmacol.* **1996**, *316*, 325–331. [CrossRef]
36. Kull, B.; Ferré, S.; Arslan, G.; Svenssonsson, P.; Fux, K.; Owman, C.; Fredholm, B.B. Reciprocal Interactions Between Adenosine A2A and Dopamine D2 Receptors in Chinese Hamster Ovary Cells Co-Transfected With the Two Receptors. *Biochem. Pharmacol.* **1999**, *58*, 1035–1045. [CrossRef]
37. Torvinen, M.; Kozell, L.B.; Neve, K.A.; Agnati, L.F.; Fuxe, K. Biochemical Identification of the Dopamine D2 Receptor Domains Interacting With the Adenosine A2A Receptor. *J. Mol. Neurosci.* **2004**, *24*, 173–180. [CrossRef]
38. Fernández-Dueñas, V.; Gómez-Soler, M.; Jacobson, K.A.; Kumar, S.T.; Fux, K.; Borroto-Escuela, D.O.; Ciruela, F. Molecular Determinants of A2AR-D2R Allostery: Role of the Intracellular Loop 3 of the D2R. *J. Neurochem.* **2012**, *123*, 373–384. [CrossRef]
39. Fernández-Dueñas, V.; Gómez-Soler, M.; Morató, X.; Núñez, F.; Das, A.; Kumar, T.S.; Jaumà, S.; Jacobson, K.A.; Ciruela, F. Dopamine D(2) Receptor-Mediated Modulation of Adenosine A(2A) Receptor Agonist Binding Within the A(2A)R/D(2)R Oligomer Framework. *Neurochem. Int.* **2013**, *63*, 42–46. [CrossRef] [PubMed]
40. Trincavelli, M.L.; Cuboni, S.; Catena, D.M.; Maggio, R.; Klotz, K.N.; Novi, F.; Panighini, A.; Daniele, S.; Martini, C. Receptor Crosstalk: Haloperidol Treatment Enhances A(2A) Adenosine Receptor Functioning in a Transfected Cell Model. *Purinergic Signal.* **2010**, *6*, 373–381. [CrossRef]
41. Bonaventura, J.; Navarro, G.; Casadó-Anguera, V.; Azdad, K.; Rea, W.; Moreno, E.; Brugarolas, M.; Mallol, J.; Canela, E.I.; Lluís, C.; et al. Allosteric Interactions Between Agonists and Antagonists Within the Adenosine A2A Receptor-Dopamine D2 Receptor Heterotetramer. *Proc. Natl. Acad. Sci. USA* **2015**, *112*, E3609–E3618. [CrossRef]
42. Díaz-Cabiale, Z.; Hurd, Y.; Guidolin, D.; Finnman, U.B.; Zoli, M.; Agnati, L.F.; Vanderhaeghen, J.J.; Fux, K.; Ferré, S. Adenosine A2A Agonist CGS 21680 Decreases the Affinity of Dopamine D2 Receptors for Dopamine in Human Striatum. *Neuroreport* **2001**, *12*, 1831–1834. [CrossRef]

43. Schiffmann, S.N.; Libert, F.; Vassart, G.; Vanderhaeghen, J.J. Distribution of Adenosine A2 Receptor mRNA in the Human Brain. *Neurosci. Lett.* **1991**, *130*, 177–181. [[CrossRef](#)]
44. Fink, J.S.; Weaver, D.R.; Rivkees, S.A.; Peterfreund, R.A.; Pollack, A.E.; Adler, E.M.; Reppert, S.M. Molecular Cloning of the Rat A2 Adenosine Receptor: Selective Co-Expression With D2 Dopamine Receptors in Rat Striatum. *Mol. Brain Res.* **1992**, *14*, 186–195. [[CrossRef](#)]
45. Peterfreund, R.A.; MacCollin, M.; Gusella, J.; Fink, J.S. Characterization and Expression of the Human A2a Adenosine Receptor Gene. *J. Neurochem.* **1996**, *66*, 362–368. [[CrossRef](#)] [[PubMed](#)]
46. Dixon, A.K.; Gubitz, A.K.; Sirinathsinghji, D.J.; Richardson, P.J.; Freeman, T.C. Tissue Distribution of Adenosine Receptor MRNAs in the Rat. *Br. J. Pharmacol.* **1996**, *118*, 1461–1468. [[CrossRef](#)] [[PubMed](#)]
47. Svenningsson, P.; Le, M.C.; Aubert, I.; Burbaud, P.; Fredholm, B.B.; Bloch, B. Cellular Distribution of Adenosine A2A Receptor mRNA in the Primate Striatum. *J. Comp. Neurol.* **1998**, *399*, 229–240. [[CrossRef](#)]
48. Kaelin-Lang, A.; Liniger, P.; Probst, A.; Lauterburg, T.; Burgunder, J.M. Adenosine A2A Receptor Gene Expression in the Normal Striatum and After 6-OH-Dopamine Lesion. *J. Neural. Transm.* **2000**, *107*, 851–859. [[CrossRef](#)] [[PubMed](#)]
49. Martinez-Mir, M.I.; Probst, A.; Palacios, J.M. Adenosine A2 Receptors: Selective Localization in the Human Basal Ganglia and Alterations with Disease. *Neuroscience* **1991**, *42*, 697–706. [[CrossRef](#)]
50. Nonaka, H.; Mori, A.; Ichimura, M.; Shindou, T.; Yanagawa, K.; Shimada, J.; Kase, H. Binding of [<sup>3</sup>H]KF17837S, a Selective Adenosine A2 Receptor Antagonist, to Rat Brain Membranes. *Mol. Pharmacol.* **1994**, *46*, 817–822. [[PubMed](#)]
51. Fredholm, B.B.; Lindström, K.; Dionisotti, S.; Ongini, E. [<sup>3</sup>H]SCH 58261, a Selective Adenosine A2A Receptor Antagonist, Is a Useful Ligand in Autoradiographic Studies. *J. Neurochem.* **1998**, *70*, 1210–1216. [[CrossRef](#)] [[PubMed](#)]
52. Rosin, D.L.; Robeva, A.; Woodard, R.L.; Guyenet, P.G.; Linden, J. Immunohistochemical Localization of Adenosine A2A Receptors in the Rat Central Nervous System. *J. Comp. Neurol.* **1998**, *401*, 163–186. [[CrossRef](#)]
53. Kull, B.; Svenningsson, P.; Hall, H.; Fredholm, B.B. GTP Differentially Affects Antagonist Radioligand Binding to Adenosine A(1) and A(2A) Receptors in Human Brain. *Neuropharmacology* **2000**, *39*, 2374–2380. [[CrossRef](#)]
54. Alexander, S.P.; Millns, P.J. [(<sup>3</sup>H)]ZM241385—an Antagonist Radioligand for Adenosine A(2A) Receptors in Rat Brain. *Eur. J. Pharmacol.* **2001**, *411*, 205–210. [[CrossRef](#)]
55. DeMet, E.M.; Chicz-DeMet, A. Localization of Adenosine A2A-Receptors in Rat Brain With [<sup>3</sup>H]ZM-241385. *Naunyn Schmiedebergs Arch. Pharmacol.* **2002**, *366*, 478–481. [[CrossRef](#)] [[PubMed](#)]
56. Calon, F.; Dridi, M.; Hornykiewicz, O.; Bédard, P.J.; Rajput, A.H.; Di, P.T. Increased Adenosine A2A Receptors in the Brain of Parkinson’s Disease Patients With Dyskinesias. *Brain* **2004**, *127*, 1075–1084. [[CrossRef](#)] [[PubMed](#)]
57. Bogenpohl, J.W.; Ritter, S.L.; Hall, R.A.; Smith, Y. Adenosine A2A Receptor in the Monkey Basal Ganglia: Ultrastructural Localization and Colocalization with the Metabotropic Glutamate Receptor 5 in the Striatum. *J. Comp. Neurol.* **2012**, *520*, 570–589. [[CrossRef](#)] [[PubMed](#)]
58. Boyson, S.J.; McGonigle, P.; Molinoff, P.B. Quantitative Autoradiographic Localization of the D1 and D2 Subtypes of Dopamine Receptors in Rat Brain. *J. Neurosci.* **1986**, *6*, 3177–3188. [[CrossRef](#)] [[PubMed](#)]
59. De Keyser, J.; Claeys, A.; De Backer, J.P.; Ebinger, G.; Roels, F.; Vauquelin, G. Autoradiographic Localization of D1 and D2 Dopamine Receptors in the Human Brain. *Neurosci. Lett.* **1988**, *91*, 142–147. [[CrossRef](#)]
60. Meador-Woodruff, J.H.; Mansour, A.; Bunzow, J.R.; Van Tol, H.H.; Watson, S.J., Jr.; Civelli, O. Distribution of D2 Dopamine Receptor mRNA in Rat Brain. *Proc. Natl. Acad. Sci. USA* **1989**, *86*, 7625–7628. [[CrossRef](#)]
61. Camps, M.; Cortés, R.; Gueye, B.; Probst, A.; Palacios, J.M. Dopamine Receptors in Human Brain: Autoradiographic Distribution of D2 Sites. *Neuroscience* **1989**, *28*, 275–290. [[CrossRef](#)]
62. Landwehrmeyer, B.; Mengod, G.; Palacios, J.M. Differential Visualization of Dopamine D2 and D3 Receptor Sites in Rat Brain. A Comparative Study Using In Situ Hybridization Histochemistry and Ligand Binding Autoradiography. *Eur. J. Neurosci.* **1993**, *5*, 145–153. [[CrossRef](#)] [[PubMed](#)]
63. Le Moine, C.; Normand, E.; Guitteny, A.F.; Fouque, B.; Teoule, R.; Bloch, B. Dopamine Receptor Gene Expression by Enkephalin Neurons in Rat Forebrain. *Proc. Natl. Acad. Sci. USA* **1990**, *87*, 230–234. [[CrossRef](#)]
64. Kemp, J.M.; Powell, T.P. The Structure of the Caudate Nucleus of the Cat: Light and Electron Microscopy. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **1971**, *262*, 383–401.
65. Alexander, G.E.; Crutcher, M.D. Functional Architecture of Basal Ganglia Circuits: Neural Substrates of Parallel Processing. *Trends Neurosci.* **1990**, *13*, 266–271. [[CrossRef](#)]
66. Gerfen, C.R. The Neostratal Mosaic: Multiple Levels of Compartmental Organization. *Trends Neurosci.* **1992**, *15*, 133–139. [[CrossRef](#)]
67. Kawaguchi, Y.; Wilson, C.J.; Emson, P.C. Projection Subtypes of Rat Neostratal Matrix Cells Revealed by Intracellular Injection of Biocytin. *J. Neurosci.* **1990**, *10*, 3421–3438. [[CrossRef](#)]
68. Gerfen, C.R.; Engber, T.M.; Mahan, L.C.; Susel, Z.; Chase, T.N.; Monsma, F.J., Jr.; Sibley, D.R. D1 and D2 Dopamine Receptor-Regulated Gene Expression of Striatonigral and Striatopallidal Neurons. *Science* **1990**, *250*, 1429–1432. [[CrossRef](#)] [[PubMed](#)]
69. Gerfen, C.R.; Surmeier, D.J. Modulation of Striatal Projection Systems by Dopamine. *Annu. Rev. Neurosci.* **2011**, *34*, 441–466. [[CrossRef](#)] [[PubMed](#)]

70. Schiffmann, S.N.; Libert, F.; Vassart, G.; Dumont, J.E.; Vanderhaeghen, J.J. A Cloned G Protein-Coupled Protein With a Distribution Restricted to Striatal Medium-Sized Neurons. Possible Relationship with D1 Dopamine Receptor. *Brain Res.* **1990**, *519*, 333–337. [[CrossRef](#)]
71. Weiner, D.M.; Levey, A.I.; Brann, M.R. Expression of Muscarinic Acetylcholine and Dopamine Receptor MRNAs in Rat Basal Ganglia. *Proc. Natl. Acad. Sci. USA* **1990**, *87*, 7050–7054. [[CrossRef](#)] [[PubMed](#)]
72. Schiffmann, S.N.; Jacobs, O.; Vanderhaeghen, J.J. Striatal Restricted Adenosine A2 Receptor (RDC8) Is Expressed by Enkephalin but Not by Substance P Neurons: An in Situ Hybridization Histochemistry Study. *J. Neurochem.* **1991**, *57*, 1062–1067. [[CrossRef](#)] [[PubMed](#)]
73. Augood, S.J.; Emson, P.C. Adenosine A2a Receptor mRNA Is Expressed by Enkephalin Cells but Not by Somatostatin Cells in Rat Striatum: A Co-Expression Study. *Mol. Brain Res.* **1994**, *22*, 204–210. [[CrossRef](#)]
74. Svenssonsson, P.; Le, M.C.; Kull, B.; Sunahara, R.; Bloch, B.; Fredholm, B.B. Cellular Expression of Adenosine A2A Receptor Messenger RNA in the Rat Central Nervous System With Special Reference to Dopamine Innervated Areas. *Neuroscience* **1997**, *80*, 1171–1185. [[CrossRef](#)]
75. Ferré, S.; Fuxe, K.; von Euler, G.; Johansson, B.; Fredholm, B.B. Adenosine-Dopamine Interactions in the Brain. *Neuroscience* **1992**, *51*, 501–512. [[CrossRef](#)]
76. Ferré, S.; O'Connor, W.T.; Fuxe, K.; Ungerstedt, U. The Striopallidal Neuron: A Main Locus for Adenosine-Dopamine Interactions in the Brain. *J. Neurosci.* **1993**, *13*, 5402–5406. [[CrossRef](#)]
77. Miyazaki, I.; Asanuma, M.; Diaz-Corrales, F.J.; Miyoshi, K.; Ogawa, N. Direct Evidence for Expression of Dopamine Receptors in Astrocytes from Basal Ganglia. *Brain Res.* **2004**, *1029*, 120–123. [[CrossRef](#)] [[PubMed](#)]
78. Daré, E.; Schulte, G.; Karovic, O.; Hammarberg, C.; Fredholm, B.B. Modulation of Glial Cell Functions by Adenosine Receptors. *Physiol. Behav.* **2007**, *92*, 15–20. [[CrossRef](#)]
79. Matos, M.; Augusto, E.; Santos-Rodrigues, A.D.; Schwarzschild, M.A.; Chen, J.F.; Cunha, R.A.; Agostinho, P. Adenosine A2A Receptors Modulate Glutamate Uptake in Cultured Astrocytes and Gliosomes. *Glia* **2012**, *60*, 702–716. [[CrossRef](#)]
80. Matos, M.; Augusto, E.; Agostinho, P.; Cunha, R.A.; Chen, J.F. Antagonistic Interaction Between Adenosine A2A Receptors and Na<sup>+</sup>/K<sup>+</sup>-ATPase-A2 Controlling Glutamate Uptake in Astrocytes. *J. Neurosci.* **2013**, *33*, 18492–18502. [[CrossRef](#)]
81. Cervetto, C.; Venturini, A.; Passalacqua, M.; Guidolin, D.; Genedani, S.; Fuxe, K.; Borroto-Escuela, D.O.; Cortelli, P.; Woods, A.; Maura, G.; et al. A2A-D2 Receptor-Receptor Interaction Modulates Gliotransmitter Release From Striatal Astrocyte Processes. *J. Neurochem.* **2017**, *140*, 268–279. [[CrossRef](#)]
82. Tozzi, A.; de Iure, A.; Di Filippo, M.; Tantucci, M.; Costa, C.; Borsini, F.; Ghiglieri, V.; Giampà, C.; Fusco, F.R.; Picconi, B.; et al. The Distinct Role of Medium Spiny Neurons and Cholinergic Interneurons in the D2/A2A Receptor Interaction in the Striatum: Implications for Parkinson’s Disease. *J. Neurosci.* **2011**, *31*, 1850–1862. [[CrossRef](#)]
83. Levey, A.I.; Hersch, S.M.; Rye, D.B.; Sunahara, R.K.; Niznik, H.B.; Kitt, C.A.; Price, D.L.; Maggio, R.; Brann, M.R.; Ciliax, B.J. Localization of D1 and D2 Dopamine Receptors in Brain With Subtype-Specific Antibodies. *Proc. Natl. Acad. Sci. USA* **1993**, *90*, 8861–8865. [[CrossRef](#)] [[PubMed](#)]
84. Hillion, J.; Canals, M.; Torvinen, M.; Casado, V.; Scott, R.; Terasmaa, A.; Hansson, A.; Watson, S.; Olah, M.E.; Mallol, J.; et al. Coaggregation, Cointernalization, and Codesensitization of Adenosine A2A Receptors and Dopamine D2 Receptors. *J. Biol. Chem.* **2002**, *277*, 18091–18097. [[CrossRef](#)] [[PubMed](#)]
85. Agnati, L.F.; Fuxe, K.; Torvinen, M.; Genedani, S.; Franco, R.; Watson, S.; Nussdorfer, G.G.; Leo, G.; Guidolin, D. New Methods to Evaluate Colocalization of Fluorophores in Immunocytochemical Preparations As Exemplified by a Study on A2A and D2 Receptors in Chinese Hamster Ovary Cells. *J. Histochem. Cytochem.* **2005**, *53*, 941–953. [[CrossRef](#)] [[PubMed](#)]
86. Torvinen, M.; Torri, C.; Tombesi, A.; Marcellino, D.; Watson, S.; Lluis, C.; Franco, R.; Fuxe, K.; Agnati, L.F. Trafficking of Adenosine A2A and Dopamine D2 Receptors. *J. Mol. Neurosci.* **2005**, *25*, 191–200. [[CrossRef](#)]
87. Genedani, S.; Guidolin, D.; Leo, G.; Filaferro, M.; Torvinen, M.; Woods, A.S.; Fuxe, K.; Ferré, S.; Agnati, L.F. Computer-Assisted Image Analysis of Caveolin-1 Involvement in the Internalization Process of Adenosine A2A-Dopamine D2 Receptor Heterodimers. *J. Mol. Neurosci.* **2005**, *26*, 177–184. [[CrossRef](#)]
88. Borroto-Escuela, D.O.; Romero-Fernandez, W.; Tarakanov, A.O.; Ciruela, F.; Agnati, L.F.; Fuxe, K. On the Existence of a Possible A2A-D2-β-Arrestin2 Complex: A2A Agonist Modulation of D2 Agonist-Induced β-Arrestin2 Recruitment. *J. Mol. Biol.* **2011**, *406*, 687–699. [[CrossRef](#)]
89. Huang, L.; Wu, D.D.; Zhang, L.; Feng, L.Y. Modulation of A2a Receptor Antagonist on D2 Receptor Internalization and ERK Phosphorylation. *Acta Pharmacol. Sin.* **2013**, *34*, 1292–1300. [[CrossRef](#)] [[PubMed](#)]
90. Trincavelli, M.L.; Daniele, S.; Orlandini, E.; Navarro, G.; Casadó, V.; Giacomelli, C.; Nencetti, S.; Nuti, E.; Macchia, M.; Huebner, H.; et al. A New D2 Dopamine Receptor Agonist Allosterically Modulates A(2A) Adenosine Receptor Signalling by Interacting With the A(2A)/D2 Receptor Heteromer. *Cell Signal.* **2012**, *24*, 951–960. [[CrossRef](#)] [[PubMed](#)]
91. Navarro, G.; Carrión, P.; Gandía, J.; Ciruela, F.; Casadó, V.; Cortés, A.; Mallol, J.; Canela, E.I.; Lluis, C.; Franco, R. Detection of Heteromers Formed by Cannabinoid CB1, Dopamine D2, and Adenosine A2A G-Protein-Coupled Receptors by Combining Bimolecular Fluorescence Complementation and Bioluminescence Energy Transfer. *Sci. World J.* **2008**, *8*, 1088–1097. [[CrossRef](#)]
92. Agnati, L.F.; Guidolin, D.; Leo, G.; Carone, C.; Genedani, S.; Fuxe, K. Receptor-Receptor Interactions: A Novel Concept in Brain Integration. *Prog. Neurobiol.* **2010**, *90*, 157–175. [[CrossRef](#)] [[PubMed](#)]

93. Cavig, M.; Lluís, C.; Moreno, E.; Bakesová, J.; Canela, E.I.; Navarro, G. Production of Functional Recombinant G-Protein Coupled Receptors for Heteromerization Studies. *J. Neurosci. Methods* **2011**, *199*, 258–264. [[CrossRef](#)]
94. Trifilieff, P.; Rives, M.L.; Urizar, E.; Piskorowski, R.A.; Vishwasrao, H.D.; Castrillon, J.; Schmauss, C.; Slättman, M.; Gullberg, M.; Javitch, J.A. Detection of Antigen Interactions Ex Vivo by Proximity Ligation Assay: Endogenous Dopamine D2-Adenosine A2A Receptor Complexes in the Striatum. *Biotechniques* **2011**, *51*, 111–118. [[CrossRef](#)] [[PubMed](#)]
95. Fernández-Dueñas, V.; Llorente, J.; Gandía, J.; Borroto-Escuela, D.O.; Agnati, L.F.; Tasca, C.I.; Fuxé, K.; Ciruela, F. Fluorescence Resonance Energy Transfer-Based Technologies in the Study of Protein-Protein Interactions at the Cell Surface. *Methods* **2012**, *57*, 467–472. [[CrossRef](#)] [[PubMed](#)]
96. Borroto-Escuela, D.O.; Romero-Fernandez, W.; Garriga, P.; Ciruela, F.; Narvaez, M.; Tarakanov, A.O.; Palkovits, M.; Agnati, L.F.; Fuxé, K. G Protein-Coupled Receptor Heterodimerization in the Brain. *Methods Enzymol.* **2013**, *521*, 281–294. [[PubMed](#)]
97. Fernández-Dueñas, V.; Gómez-Soler, M.; Valle-León, M.; Watanabe, M.; Ferrer, I.; Ciruela, F. Revealing Adenosine A(2A)-Dopamine D(2) Receptor Heteromers in Parkinson’s Disease Post-Mortem Brain Through a New AlphaScreen-Based Assay. *Int. J. Mol. Sci.* **2019**, *20*, 3600. [[CrossRef](#)] [[PubMed](#)]
98. López-Cano, M.; Fernández-Dueñas, V.; Ciruela, F. Proximity Ligation Assay Image Analysis Protocol: Addressing Receptor-Receptor Interactions. *Methods Mol. Biol.* **2019**, *2040*, 41–50. [[PubMed](#)]
99. Söderberg, O.; Gullberg, M.; Jarvius, M.; Ridderstråle, K.; Leuchowius, K.J.; Jarvius, J.; Wester, K.; Hydbring, P.; Bahram, F.; Larsson, L.G.; et al. Direct Observation of Individual Endogenous Protein Complexes in Situ by Proximity Ligation. *Nat. Methods* **2006**, *3*, 995–1000. [[CrossRef](#)]
100. Gandia, J.; Galino, J.; Amaral, O.B.; Soriano, A.; Lluís, C.; Franco, R.; Ciruela, F. Detection of Higher-Order G Protein-Coupled Receptor Oligomers by a Combined BRET-BiFC Technique. *FEBS Lett.* **2008**, *582*, 2979–2984. [[CrossRef](#)]
101. Kamiya, T.; Saitoh, O.; Yoshioka, K.; Nakata, H. Oligomerization of Adenosine A2A and Dopamine D2 Receptors in Living Cells. *Biochem. Biophys. Res. Commun.* **2003**, *306*, 544–549. [[CrossRef](#)]
102. Canals, M.; Marcellino, D.; Fanelli, F.; Ciruela, F.; de Benedetti, P.; Goldberg, S.R.; Neve, K.; Fuxé, K.; Agnati, L.F.; Woods, A.S.; et al. Adenosine A2A-Dopamine D2 Receptor-Receptor Heteromerization: Qualitative and Quantitative Assessment by Fluorescence and Bioluminescence Energy Transfer. *J. Biol. Chem.* **2003**, *278*, 46741–46749. [[CrossRef](#)] [[PubMed](#)]
103. Prakash, A.; Luthra, P.M. Insilico Study of the A(2A)R-D(2)R Kinetics and Interfacial Contact Surface for Heteromerization. *Amino Acids* **2012**, *43*, 1451–1464. [[CrossRef](#)] [[PubMed](#)]
104. Borroto-Escuela, D.O.; Rodriguez, D.; Romero-Fernandez, W.; Kapla, J.; Jaiteh, M.; Ranganathan, A.; Lazarova, T.; Fuxé, K.; Carlsson, J. Mapping the Interface of a GPCR Dimer: A Structural Model of the A(2A) Adenosine and D(2) Dopamine Receptor Heteromer. *Front. Pharmacol.* **2018**, *9*, 829. [[CrossRef](#)]
105. Tarakanov, A.O.; Fuxé, K.G. Triplet Puzzle: Homologies of Receptor Heteromers. *J. Mol. Neurosci.* **2010**, *41*, 294–303. [[CrossRef](#)]
106. Ciruela, F.; Burgueño, J.; Casadó, V.; Canals, M.; Marcellino, D.; Goldberg, S.R.; Bader, M.; Fuxé, K.; Agnati, L.F.; Lluis, C.; et al. Combining Mass Spectrometry and Pull-Down Techniques for the Study of Receptor Heteromerization. Direct Epitope-Epitope Electrostatic Interactions Between Adenosine A2A and Dopamine D2 Receptors. *Anal. Chem.* **2004**, *76*, 5354–5363. [[CrossRef](#)] [[PubMed](#)]
107. Woods, A.S.; Ferré, S. Amazing Stability of the Arginine-Phosphate Electrostatic Interaction. *J. Proteome Res.* **2005**, *4*, 1397–1402. [[CrossRef](#)] [[PubMed](#)]
108. Borroto-Escuela, D.O.; Marcellino, D.; Narvaez, M.; Flajolet, M.; Heintz, N.; Agnati, L.; Ciruela, F.; Fuxé, K. A Serine Point Mutation in the Adenosine A2AR C-Terminal Tail Reduces Receptor Heteromerization and Allosteric Modulation of the Dopamine D2R. *Biochem. Biophys. Res. Commun.* **2010**, *394*, 222–227. [[CrossRef](#)]
109. Borroto-Escuela, D.O.; Romero-Fernandez, W.; Tarakanov, A.O.; Gómez-Soler, M.; Corrales, F.; Marcellino, D.; Narvaez, M.; Frankowska, M.; Flajolet, M.; Heintz, N.; et al. Characterization of the A2AR-D2R Interface: Focus on the Role of the C-Terminal Tail and the Transmembrane Helices. *Biochem. Biophys. Res. Commun.* **2010**, *402*, 801–807. [[CrossRef](#)] [[PubMed](#)]
110. Navarro, G.; Aymerich, M.S.; Marcellino, D.; Cortés, A.; Casadó, V.; Mallol, J.; Canela, E.I.; Agnati, L.; Woods, A.S.; Fuxé, K.; et al. Interactions Between Calmodulin, Adenosine A2A, and Dopamine D2 Receptors. *J. Biol. Chem.* **2009**, *284*, 28058–28068. [[CrossRef](#)] [[PubMed](#)]
111. Kamiya, T.; Saitoh, O.; Nakata, H. Functional Expression of Single-Chain Heterodimeric G-Protein-Coupled Receptors for Adenosine and Dopamine. *Cell Struct. Funct.* **2005**, *29*, 139–145. [[CrossRef](#)]
112. Vidi, P.A.; Chemel, B.R.; Hu, C.D.; Watts, V.J. Ligand-Dependent Oligomerization of Dopamine D(2) and Adenosine A(2A) Receptors in Living Neuronal Cells. *Mol. Pharmacol.* **2008**, *74*, 544–551. [[CrossRef](#)] [[PubMed](#)]
113. Lin, C.Y.; Lai, H.L.; Chen, H.M.; Siew, J.J.; Hsiao, C.T.; Chang, H.C.; Liao, K.S.; Tsai, S.C.; Wu, C.Y.; Kitajima, K.; et al. Functional Roles of ST8SIA3-Mediated Sialylation of Striatal Dopamine D(2) and Adenosine A(2A) Receptors. *Transl. Psychiatry* **2019**, *9*, 209. [[CrossRef](#)]
114. Pelassa, S.; Guidolin, D.; Venturini, A.; Averna, M.; Frumento, G.; Campanini, L.; Bernardi, R.; Cortelli, P.; Buonaura, G.C.; Maura, G.; et al. A2A-D2 Heteromers on Striatal Astrocytes: Biochemical and Biophysical Evidence. *Int. J. Mol. Sci.* **2019**, *20*, 2457. [[CrossRef](#)]
115. Fernández-Dueñas, V.; Taura, J.J.; Cottet, M.; Gómez-Soler, M.; López-Cano, M.; Ledent, C.; Watanabe, M.; Trinquet, E.; Pin, J.P.; Luján, R.; et al. Untangling Dopamine-Adenosine Receptor-Receptor Assembly in Experimental Parkinsonism in Rats. *Dis. Models Mech.* **2015**, *8*, 57–63. [[CrossRef](#)]

116. He, Y.; Li, Y.; Chen, M.; Pu, Z.; Zhang, F.; Chen, L.; Ruan, Y.; Pan, X.; He, C.; Chen, X.; et al. Habit Formation After Random Interval Training Is Associated With Increased Adenosine A(2A) Receptor and Dopamine D(2) Receptor Heterodimers in the Striatum. *Front. Mol. Neurosci.* **2016**, *9*, 151. [[CrossRef](#)]
117. Borroto-Escuela, D.O.; Narváez, M.; Wydra, K.; Pintsuk, J.; Pinton, L.; Jimenez-Beristain, A.; Di Palma, M.; Jastrzebska, J.; Filip, M.; Fuxe, K. Cocaine Self-Administration Specifically Increases A2AR-D2R and D2R-Sigma1R Heteroreceptor Complexes in the Rat Nucleus Accumbens Shell. Relevance for Cocaine Use Disorder. *Pharmacol. Biochem. Behav.* **2017**, *155*, 24–31. [[CrossRef](#)] [[PubMed](#)]
118. Bonaventura, J.; Rico, A.J.; Moreno, E.; Sierra, S.; Sánchez, M.; Luquin, N.; Farré, D.; Müller, C.E.; Martínez-Pinilla, E.; Cortés, A.; et al. L-DOPA-Treatment in Primates Disrupts the Expression of A(2A) Adenosine-CB(1) Cannabinoid-D(2) Dopamine Receptor Heteromers in the Caudate Nucleus. *Neuropharmacology* **2014**, *79*, 90–100. [[CrossRef](#)] [[PubMed](#)]
119. Zhu, Y.; Mészáros, J.; Walle, R.; Fan, R.; Sun, Z.; Dwork, A.J.; Trifilieff, P.; Javitch, J.A. Detecting G Protein-Coupled Receptor Complexes in Postmortem Human Brain With Proximity Ligation Assay and a Bayesian Classifier. *Biotechniques* **2020**, *68*, 122–129. [[CrossRef](#)] [[PubMed](#)]
120. Ferré, S.; Baler, R.; Bouvier, M.; Caron, M.G.; Devi, L.A.; Durroux, T.; Fuxe, K.; George, S.R.; Javitch, J.A.; Lohse, M.J.; et al. Building a New Conceptual Framework for Receptor Heteromers. *Nat. Chem. Biol.* **2009**, *5*, 131–134. [[CrossRef](#)] [[PubMed](#)]
121. Ferré, S.; Casadó, V.; Devi, L.A.; Filizola, M.; Jockers, R.; Lohse, M.J.; Milligan, G.; Pin, J.P.; Guitart, X. G Protein-Coupled Receptor Oligomerization Revisited: Functional and Pharmacological Perspectives. *Pharmacol. Rev.* **2014**, *66*, 413–434. [[CrossRef](#)] [[PubMed](#)]
122. Ferré, S. The GPCR Heterotetramer: Challenging Classical Pharmacology. *Trends Pharmacol. Sci.* **2015**, *36*, 145–152. [[CrossRef](#)] [[PubMed](#)]
123. Casadó-Anguera, V.; Bonaventura, J.; Moreno, E.; Navarro, G.; Cortés, A.; Ferré, S.; Casadó, V. Evidence for the Heterotetrameric Structure of the Adenosine A2A-Dopamine D2 Receptor Complex. *Biochem. Soc. Trans.* **2016**, *44*, 595–600. [[CrossRef](#)] [[PubMed](#)]
124. Navarro, G.; Cordini, A.; Brugarolas, M.; Moreno, E.; Aguinaga, D.; Pérez-Benito, L.; Ferre, S.; Cortés, A.; Casadó, V.; Mallol, J.; et al. Cross-Communication Between G(i) and G(s) in a G-Protein-Coupled Receptor Heterotetramer Guided by a Receptor C-Terminal Domain. *BMC Biol.* **2018**, *16*, 24. [[CrossRef](#)]
125. Ferré, S.; Ciruela, F. Functional and Neuroprotective Role of Striatal Adenosine A(2A) Receptor Heterotetramers. *J. Caffeine Adenosine Res.* **2019**, *9*, 89–97. [[CrossRef](#)]
126. Gomes, I.; Ayoub, M.A.; Fujita, W.; Jaeger, W.C.; Pfleger, K.D.; Devi, L.A. G Protein-Coupled Receptor Heteromers. *Annu. Rev. Pharmacol. Toxicol.* **2016**, *56*, 403–425. [[CrossRef](#)]
127. Fuxe, K.; Borroto-Escuela, D.O.; Marcellino, D.; Romero-Fernandez, W.; Frankowska, M.; Guidolin, D.; Filip, M.; Ferraro, L.; Woods, A.S.; Tarakanov, A.; et al. GPCR Heteromers and Their Allosteric Receptor-Receptor Interactions. *Curr. Med. Chem.* **2012**, *19*, 356–363. [[CrossRef](#)]
128. Fuxe, K.; Marcellino, D.; Leo, G.; Agnati, L.F. Molecular Integration via Allosteric Interactions in Receptor Heteromers. A Working Hypothesis. *Curr. Opin. Pharmacol.* **2010**, *10*, 14–22. [[CrossRef](#)]
129. Ferré, S.; Bonaventura, J.; Tomasi, D.; Navarro, G.; Moreno, E.; Cortés, A.; Lluís, C.; Casadó, V.; Volkow, N.D. Allosteric Mechanisms Within the Adenosine A2A-Dopamine D2 Receptor Heterotetramer. *Neuropharmacology* **2016**, *104*, 154–160. [[CrossRef](#)]
130. Fuxe, K.; Marcellino, D.; Borroto-Escuela, D.O.; Frankowska, M.; Ferraro, L.; Guidolin, D.; Ciruela, F.; Agnati, L.F. The Changing World of G Protein-Coupled Receptors: From Monomers to Dimers and Receptor Mosaics with Allosteric Receptor-Receptor Interactions. *J. Recept. Signal Transduct. Res.* **2010**, *30*, 272–283. [[CrossRef](#)]
131. Agnati, L.F.; Leo, G.; Genedani, S.; Andreoli, N.; Marcellino, D.; Woods, A.; Piron, L.; Guidolin, D.; Fuxe, K. Structural Plasticity in G-Protein Coupled Receptors As Demonstrated by the Allosteric Actions of Homocysteine and Computer-Assisted Analysis of Disordered Domains. *Brain Res. Rev.* **2008**, *58*, 459–474. [[CrossRef](#)]
132. Agnati, L.F.; Guidolin, D.; Vilardaga, J.P.; Ciruela, F.; Fuxe, K. On the Expanding Terminology in the GPCR Field: The Meaning of Receptor Mosaics and Receptor Heteromers. *J. Recept. Signal Transduct. Res.* **2010**, *30*, 287–303. [[CrossRef](#)] [[PubMed](#)]
133. Popoli, P.; Pèzzolla, A.; Torvinen, M.; Reggio, R.; Pintor, A.; Scarchilli, L.; Fuxe, K.; Ferré, S. The Selective mGlu(5) Receptor Agonist CHPG Inhibits Quinpirole-Induced Turning in 6-Hydroxydopamine-Lesioned Rats and Modulates the Binding Characteristics of Dopamine D(2) Receptors in the Rat Striatum: Interactions With Adenosine A(2a) Receptors. *Neuropsychopharmacology* **2001**, *25*, 505–513. [[CrossRef](#)]
134. Cabello, N.; Gandía, J.; Bertarelli, D.C.; Watanabe, M.; Lluís, C.; Franco, R.; Ferré, S.; Luján, R.; Ciruela, F. Metabotropic Glutamate Type 5, Dopamine D2 and Adenosine A2a Receptors Form Higher-Order Oligomers in Living Cells. *J. Neurochem.* **2009**, *109*, 1497–1507. [[CrossRef](#)] [[PubMed](#)]
135. Navarro, G.; Moreno, E.; Bonaventura, J.; Brugarolas, M.; Farré, D.; Aguinaga, D.; Mallol, J.; Cortés, A.; Casadó, V.; Lluís, C.; et al. Cocaine Inhibits Dopamine D2 Receptor Signaling Via Sigma-1-D2 Receptor Heteromers. *PLoS ONE* **2013**, *8*, e61245. [[CrossRef](#)]
136. Borroto-Escuela, D.O.; Wydra, K.; Filip, M.; Fuxe, K. A2AR-D2R Heteroreceptor Complexes in Cocaine Reward and Addiction. *Trends Pharmacol. Sci.* **2018**, *39*, 1008–1020. [[CrossRef](#)]
137. Romero-Fernandez, W.; Zhou, Z.; Beggiato, S.; Wydra, K.; Filip, M.; Tanganeli, S.; Borroto-Escuela, D.O.; Ferraro, L.; Fuxe, K. Acute Cocaine Treatment Enhances the Antagonistic Allosteric Adenosine A2A-Dopamine D2 Receptor-Receptor Interactions in Rat Dorsal Striatum Without Increasing Significantly Extracellular Dopamine Levels. *Pharmacol. Rep.* **2020**, *72*, 332–339. [[CrossRef](#)]

138. Fuxe, K.; Borroto-Escuela, D.O. Heteroreceptor Complexes and Their Allosteric Receptor-Receptor Interactions as a Novel Biological Principle for Integration of Communication in the CNS: Targets for Drug Development. *Neuropharmacology* **2016**, *41*, 380–382. [[CrossRef](#)]
139. Ferré, S.; Ciruela, F.; Woods, A.S.; Lluis, C.; Franco, R. Functional Relevance of Neurotransmitter Receptor Heteromers in the Central Nervous System. *Trends Neurosci.* **2007**, *30*, 440–446. [[CrossRef](#)]
140. Purves, D.; Augustine, G.J.; Fitzpatrick, D.; Katz, L.C.; LaMantia, A.S.; McNamara, J.O.; Williams, S.M. *Neuroscience*; Sinauer Associates: Sunderland, MA, USA, 2001.
141. DeLong, M.R. Primate Models of Movement Disorders of Basal Ganglia Origin. *Trends Neurosci.* **1990**, *13*, 281–285. [[CrossRef](#)]
142. Tepper, J.M.; Abercrombie, E.D.; Bolam, J.P. Basal Ganglia Macrocircuits. *Prog. Brain Res.* **2007**, *160*, 3–7.
143. Lanciego, J.L.; Luquin, N.; Obeso, J.A. Functional Neuroanatomy of the Basal Ganglia. *Cold Spring Harb. Perspect. Med.* **2012**, *2*, a009621. [[CrossRef](#)]
144. Parent, A.; Hazrati, L.N. Functional Anatomy of the Basal Ganglia. II. The Place of Subthalamic Nucleus and External Pallidum in Basal Ganglia Circuitry. *Brain Res. Rev.* **1995**, *20*, 128–154. [[CrossRef](#)]
145. Richardson, P.J.; Kase, H.; Jenner, P.G. Adenosine A2A Receptor Antagonists as New Agents for the Treatment of Parkinson’s Disease. *Trends Pharmacol. Sci.* **1997**, *18*, 338–344. [[CrossRef](#)]
146. Müller, C.E.; Ferré, S. Blocking Striatal Adenosine A2A Receptors: A New Strategy for Basal Ganglia Disorders. *Recent Pat. CNS Drug Discov.* **2007**, *2*, 1–21. [[CrossRef](#)] [[PubMed](#)]
147. Armentero, M.T.; Pinna, A.; Ferré, S.; Lanciego, J.L.; Müller, C.E.; Franco, R. Past, Present and Future of A(2A) Adenosine Receptor Antagonists in the Therapy of Parkinson’s Disease. *Pharmacol. Ther.* **2011**, *132*, 280–299. [[CrossRef](#)] [[PubMed](#)]
148. Szabó, N.; Kincses, Z.T.; Vécsei, L. Novel Therapy in Parkinson’s Disease: Adenosine A(2A) Receptor Antagonists. *Expert Opin. Drug Metab. Toxicol.* **2011**, *7*, 441–455. [[CrossRef](#)] [[PubMed](#)]
149. Preti, D.; Baraldi, P.G.; Moorman, A.R.; Borea, P.A.; Varani, K. History and Perspectives of A2A Adenosine Receptor Antagonists as Potential Therapeutic Agents. *Med. Res. Rev.* **2015**, *35*, 790–848. [[CrossRef](#)] [[PubMed](#)]
150. Mori, A.; Shindou, T. Modulation of GABAergic Transmission in the Striatopallidal System by Adenosine A2A Receptors: A Potential Mechanism for the Antiparkinsonian Effects of A2A Antagonists. *Neurology* **2003**, *61*, S44–S48. [[CrossRef](#)] [[PubMed](#)]
151. Fuxe, K.; Ferré, S.; Genedani, S.; Franco, R.; Agnati, L.F. Adenosine Receptor-Dopamine Receptor Interactions in the Basal Ganglia and Their Relevance for Brain Function. *Physiol. Behav.* **2007**, *92*, 210–217. [[CrossRef](#)]
152. Black, K.J.; Koller, J.M.; Campbell, M.C.; Gusnard, D.A.; Bandak, S.I. Quantification of Indirect Pathway Inhibition by the Adenosine A2a Antagonist SYN115 in Parkinson Disease. *J. Neurosci.* **2010**, *30*, 16284–16292. [[CrossRef](#)] [[PubMed](#)]
153. Aoyama, S.; Kase, H.; Borrelli, E. Rescue of Locomotor Impairment in Dopamine D2 Receptor-Deficient Mice by an Adenosine A2A Receptor Antagonist. *J. Neurosci.* **2000**, *20*, 5848–5852. [[CrossRef](#)] [[PubMed](#)]
154. Strömborg, I.; Popoli, P.; Müller, C.E.; Ferré, S.; Fuxe, K. Electrophysiological and Behavioural Evidence for an Antagonistic Modulatory Role of Adenosine A2A Receptors in Dopamine D2 Receptor Regulation in the Rat Dopamine-Denervated Striatum. *Eur. J. Neurosci.* **2000**, *12*, 4033–4037. [[CrossRef](#)] [[PubMed](#)]
155. Bové, J.; Marin, C.; Bonastre, M.; Tolosa, E. Adenosine A2A Antagonism Reverses Levodopa-Induced Motor Alterations in Hemiparkinsonian Rats. *Synapse* **2002**, *46*, 251–257. [[CrossRef](#)] [[PubMed](#)]
156. Matsuya, T.; Takuma, K.; Sato, K.; Asai, M.; Murakami, Y.; Miyoshi, S.; Noda, A.; Nagai, T.; Mizoguchi, H.; Nishimura, S.; et al. Synergistic Effects of Adenosine A2A Antagonist and L-DOPA on Rotational Behaviors in 6-Hydroxydopamine-Induced Hemiparkinsonian Mouse Model. *J. Pharmacol. Sci.* **2007**, *103*, 329–332. [[CrossRef](#)] [[PubMed](#)]
157. Hauber, W.; Neuscheler, P.; Nagel, J.; Müller, C.E. Catalepsy Induced by a Blockade of Dopamine D1 or D2 Receptors Was Reversed by a Concomitant Blockade of Adenosine A(2A) Receptors in the Caudate-Putamen of Rats. *Eur. J. Neurosci.* **2001**, *14*, 1287–1293. [[CrossRef](#)] [[PubMed](#)]
158. Kanda, T.; Tashiro, T.; Kuwana, Y.; Jenner, P. Adenosine A2A Receptors Modify Motor Function in MPTP-Treated Common Marmosets. *Neuroreport* **1998**, *9*, 2857–2860. [[CrossRef](#)] [[PubMed](#)]
159. Bibbiani, F.; Oh, J.D.; Petzer, J.P.; Castagnoli, N., Jr.; Chen, J.F.; Schwarzschild, M.A.; Chase, T.N. A2A Antagonist Prevents Dopamine Agonist-Induced Motor Complications in Animal Models of Parkinson’s Disease. *Exp. Neurol.* **2003**, *184*, 285–294. [[CrossRef](#)]
160. Mally, J.; Stone, T.W. The Effect of Theophylline on Parkinsonian Symptoms. *J Pharm Pharmacol* **1994**, *46*, 515–517. [[CrossRef](#)]
161. Kostic, V.S.; Svetel, M.; Sternic, N.; Dragasevic, N.; Przedborski, S. Theophylline Increases “on” Time in Advanced Parkinsonian Patients. *Neurology* **1999**, *52*, 1916. [[CrossRef](#)]
162. Kulisevsky, J.; Barbanoj, M.; Gironell, A.; Antonijs, R.; Casas, M.; Pascual-Sedano, B. A Double-Blind Crossover, Placebo-Controlled Study of the Adenosine A2A Antagonist Theophylline in Parkinson’s Disease. *Clin. Neuropharmacol.* **2002**, *25*, 25–31. [[CrossRef](#)] [[PubMed](#)]
163. Kitagawa, M.; Houzen, H.; Tashiro, K. Effects of Caffeine on the Freezing of Gait in Parkinson’s Disease. *Mov. Disord.* **2007**, *22*, 710–712. [[CrossRef](#)] [[PubMed](#)]
164. Bara-Jimenez, W.; Sherzai, A.; Dimitrova, T.; Favit, A.; Bibbiani, F.; Gillespie, M.; Morris, M.J.; Mouradian, M.M.; Chase, T.N. Adenosine A(2A) Receptor Antagonist Treatment of Parkinson’s Disease. *Neurology* **2003**, *61*, 293–296. [[CrossRef](#)] [[PubMed](#)]
165. Hauser, R.A.; Hubble, J.P.; Truong, D.D. Randomized Trial of the Adenosine A(2A) Receptor Antagonist Istradefylline in Advanced PD. *Neurology* **2003**, *61*, 297–303. [[CrossRef](#)] [[PubMed](#)]

166. Stacy, M.A. The US-005/US-006 Clinical Investigator Group. Istradefylline (KW-6002) As Adjunctive Therapy in Patients with Advanced Parkinson's Disease: A Positive Safety Profile with Supporting Efficacy. *Mov. Disord.* **2004**, *19*, S215–S216.
167. Hauser, R.A.; Shulman, L.M.; Trugman, J.M.; Roberts, J.W.; Mori, A.; Ballerini, R.; Sussman, N.M. Study of Istradefylline in Patients With Parkinson's Disease on Levodopa With Motor Fluctuations. *Mov. Disord.* **2008**, *23*, 2177–2185. [CrossRef]
168. LeWitt, P.A.; Guttman, M.; Tetrud, J.W.; Tuite, P.J.; Mori, A.; Chaikin, P.; Sussman, N.M. Adenosine A2A Receptor Antagonist Istradefylline (KW-6002) Reduces "Off" Time in Parkinson's Disease: A Double-Blind, Randomized, Multicenter Clinical Trial (6002-US-005). *Ann. Neurol.* **2008**, *63*, 295–302. [CrossRef]
169. Fernandez, H.H.; Greeley, D.R.; Zweig, R.M.; Wojcieszek, J.; Mori, A.; Sussman, N.M. Istradefylline As Monotherapy for Parkinson Disease: Results of the 6002-US-051 Trial. *Parkinsonism Relat. Disord.* **2010**, *16*, 16–20. [CrossRef]
170. Mizuno, Y.; Kondo, T. Adenosine A2A Receptor Antagonist Istradefylline Reduces Daily off Time in Parkinson's Disease. *Mov. Disord.* **2013**, *28*, 1138–1141. [CrossRef]
171. Kondo, T.; Mizuno, Y. A Long-Term Study of Istradefylline Safety and Efficacy in Patients with Parkinson Disease. *Clin. Neuropharmacol.* **2015**, *38*, 41–46. [CrossRef]
172. Suzuki, K.; Miyamoto, M.; Miyamoto, T.; Uchiyama, T.; Watanabe, Y.; Suzuki, S.; Kadokawa, T.; Fujita, H.; Matsubara, T.; Sakuramoto, H.; et al. Istradefylline Improves Daytime Sleepiness in Patients With Parkinson's Disease: An Open-Label, 3-Month Study. *J. Neurol. Sci.* **2017**, *380*, 230–233. [CrossRef] [PubMed]
173. Yabe, I.; Kitagawa, M.; Takahashi, I.; Matsushima, M.; Sasaki, H. The Efficacy of Istradefylline for Treating Mild Wearing-Off in Parkinson Disease. *Clin. Neuropharmacol.* **2017**, *40*, 261–263. [CrossRef] [PubMed]
174. Iijima, M.; Orimo, S.; Terashi, H.; Suzuki, M.; Hayashi, A.; Shimura, H.; Mitoma, H.; Kitagawa, K.; Okuma, Y. Efficacy of Istradefylline for Gait Disorders With Freezing of Gait in Parkinson's Disease: A Single-Arm, Open-Label, Prospective, Multicenter Study. *Expert Opin. Pharmacother.* **2019**, *20*, 1405–1411. [CrossRef] [PubMed]
175. Hauser, R.A.; Olanow, C.W.; Kieburtz, K.D.; Pourcher, E.; Docu-Axelerad, A.; Lew, M.; Kozyolkina, O.; Neale, A.; Resburg, C.; Meya, U.; et al. Tozadenant (SYN115) in Patients With Parkinson's Disease Who Have Motor Fluctuations on Levodopa: A Phase 2b, Double-Blind, Randomised Trial. *Lancet Neurol.* **2014**, *13*, 767–776. [CrossRef]
176. Dungo, R.; Deeks, E.D. Istradefylline: First Global Approval. *Drugs* **2013**, *73*, 875–882. [CrossRef]
177. Chen, J.F.; Cunha, R.A. The Belated US FDA Approval of the Adenosine A(2A) Receptor Antagonist Istradefylline for Treatment of Parkinson's Disease. *Purinergic Signal.* **2020**, *16*, 167–174. [CrossRef]
178. Seeman, P. Targeting the Dopamine D2 Receptor in Schizophrenia. *Expert Opin. Ther. Targets* **2006**, *10*, 515–531. [CrossRef] [PubMed]
179. Ferré, S.; O'Connor, W.T.; Snaprud, P.; Ungerstedt, U.; Fuxe, K. Antagonistic Interaction Between Adenosine A2A Receptors and Dopamine D2 Receptors in the Ventral Striopallidal System. Implications for the Treatment of Schizophrenia. *Neuroscience* **1994**, *63*, 765–773. [CrossRef]
180. Lara, D.R.; Dall'Igna, O.P.; Ghisolfi, E.S.; Brunstein, M.G. Involvement of Adenosine in the Neurobiology of Schizophrenia and Its Therapeutic Implications. *Prog. Neuropsychopharmacol. Biol. Psychiatry* **2006**, *30*, 617–629. [CrossRef] [PubMed]
181. Lara, D.R.; Souza, D.O. Schizophrenia: A Purinergic Hypothesis. *Med. Hypotheses* **2000**, *54*, 157–166. [CrossRef]
182. Boison, D.; Singer, P.; Shen, H.Y.; Feldon, J.; Yee, B.K. Adenosine Hypothesis of Schizophrenia—Opportunities for Pharmacotherapy. *Neuropharmacology* **2012**, *62*, 1527–1543. [CrossRef] [PubMed]
183. Kurumaji, A.; Toru, M. An Increase in [<sup>3</sup>H] CGS21680 Binding in the Striatum of Postmortem Brains of Chronic Schizophrenics. *Brain Res.* **1998**, *808*, 320–323. [CrossRef]
184. Deckert, J.; Brenner, M.; Durany, N.; Zöchling, R.; Paulus, W.; Ransmayr, G.; Tatschner, T.; Danielczyk, W.; Jellinger, K.; Riederer, P. Up-Regulation of Striatal Adenosine A(2A) Receptors in Schizophrenia. *Neuroreport* **2003**, *14*, 313–316. [CrossRef] [PubMed]
185. Hwang, Y.; Kim, J.; Shin, J.Y.; Kim, J.I.; Seo, J.S.; Webster, M.J.; Lee, D.; Kim, S. Gene Expression Profiling by mRNA Sequencing Reveals Increased Expression of Immune/Inflammation-Related Genes in the Hippocampus of Individuals With Schizophrenia. *Transl. Psychiatry* **2013**, *3*, e321. [CrossRef] [PubMed]
186. Miao, J.; Liu, L.; Yan, C.; Zhu, X.; Fan, M.; Yu, P.; Ji, K.; Huang, Y.; Wang, Y.; Zhu, G. Association Between ADORA2A Gene Polymorphisms and Schizophrenia in the North Chinese Han Population. *Neuropsychiatr. Dis. Treat.* **2019**, *15*, 2451–2458. [CrossRef]
187. Ferré, S. Adenosine-Dopamine Interactions in the Ventral Striatum. Implications for the Treatment of Schizophrenia. *Psychopharmacology* **1997**, *133*, 107–120.
188. Borroto-Escuela, D.O.; Ferraro, L.; Narvaez, M.; Tanganelli, S.; Beggiato, S.; Liu, F.; Rivera, A.; Fuxe, K. Multiple Adenosine-Dopamine (A2A-D2 Like) Heteroreceptor Complexes in the Brain and Their Role in Schizophrenia. *Cells* **2020**, *9*, 1077. [CrossRef] [PubMed]
189. Fuxe, K.; Borroto-Escuela, D.O.; Tarakanov, A.O.; Romero-Fernandez, W.; Ferraro, L.; Tanganelli, S.; Perez-Alea, M.; Di, P.M.; Agnati, L.F. Dopamine D2 Heteroreceptor Complexes and Their Receptor-Receptor Interactions in Ventral Striatum: Novel Targets for Antipsychotic Drugs. *Prog. Brain Res.* **2014**, *211*, 113–139. [PubMed]
190. Rimondini, R.; Ferré, S.; Ogren, S.O.; Fuxe, K. Adenosine A2A Agonists: A Potential New Type of Atypical Antipsychotic. *Neuropsychopharmacology* **1997**, *17*, 82–91. [CrossRef]
191. Andersen, M.B.; Fuxe, K.; Werge, T.; Gerlach, J. The Adenosine A2A Receptor Agonist CGS 21680 Exhibits Antipsychotic-Like Activity in Cebus Apella Monkeys. *Behav. Pharmacol.* **2002**, *13*, 639–644. [CrossRef]

192. Akhondzadeh, S.; Shasavand, E.; Jamilian, H.; Shabestari, O.; Kamalipour, A. Dipyridamole in the Treatment of Schizophrenia: Adenosine-Dopamine Receptor Interactions. *J. Clin. Pharm. Ther.* **2000**, *25*, 131–137. [CrossRef] [PubMed]
193. Lara, D.R.; Brunstein, M.G.; Ghisolfi, E.S.; Lobato, M.I.; Belmonte-de-Abreu, P.; Souza, D.O. Allopurinol Augmentation for Poorly Responsive Schizophrenia. *Int. Clin. Psychopharmacol.* **2001**, *16*, 235–237. [CrossRef] [PubMed]
194. Lara, D.R.; Vianna, M.R.; de Paris, F.; Quevedo, J.; Oses, J.P.; Battastini, A.M.; Sarkis, J.J.; Souza, D.O. Chronic Treatment With Clozapine, but Not Haloperidol, Increases Striatal Ecto-5'-Nucleotidase Activity in Rats. *Neuropsychobiology* **2001**, *44*, 99–102. [CrossRef] [PubMed]
195. Durieux, P.F.; Bearzatto, B.; Guiducci, S.; Buch, T.; Waisman, A.; Zoli, M.; Schiffmann, S.N.; de Kerchove, d.A. D2R Striatopallidal Neurons Inhibit Both Locomotor and Drug Reward Processes. *Nat. Neurosci.* **2009**, *12*, 393–395. [CrossRef] [PubMed]
196. Wydra, K.; Gawlinski, D.; Gawlinska, K.; Frankowska, M.; Borroto-Escuela, D.O.; Fuxé, K.; Filip, M. Adenosine A(2A)Receptors in Substance Use Disorders: A Focus on Cocaine. *Cells* **2020**, *9*, 1372. [CrossRef] [PubMed]
197. Baldo, B.A.; Koob, G.F.; Markou, A. Role of Adenosine A2 Receptors in Brain Stimulation Reward under Baseline Conditions and During Cocaine Withdrawal in Rats. *J. Neurosci.* **1999**, *19*, 11017–11026. [CrossRef]
198. Horger, B.A.; Wellman, P.J.; Morien, A.; Davies, B.T.; Schenk, S. Caffeine Exposure Sensitizes Rats to the Reinforcing Effects of Cocaine. *Neuroreport* **1991**, *2*, 53–56. [CrossRef]
199. Ferré, S. Mechanisms of the Psychostimulant Effects of Caffeine: Implications for Substance Use Disorders. *Psychopharmacology* **2016**, *233*, 1963–1979. [CrossRef]
200. Knapp, C.M.; Foye, M.M.; Cottam, N.; Ciraulo, D.A.; Kornetsky, C. Adenosine Agonists CGS 21680 and NECA Inhibit the Initiation of Cocaine Self-Administration. *Pharmacol. Biochem. Behav.* **2001**, *68*, 797–803. [CrossRef]
201. Wydra, K.; Suder, A.; Borroto-Escuela, D.O.; Filip, M.; Fuxé, K. On the Role of A2A and D2 Receptors in Control of Cocaine and Food-Seeking Behaviors in Rats. *Psychopharmacology* **2015**, *232*, 1767–1778. [CrossRef] [PubMed]
202. Wydra, K.; Suder, A.; Frankowska, M.; Borroto Escuela, D.O.; Fuxé, K.; Filip, M. Effects of Intra-Accumbal or Intra-Prefrontal Cortex Microinjections of Adenosine 2A Receptor Ligands on Responses to Cocaine Reward and Seeking in Rats. *Psychopharmacology* **2018**, *235*, 3509–3523. [CrossRef] [PubMed]
203. Borroto-Escuela, D.O.; Wydra, K.; Li, X.; Rodriguez, D.; Carlsson, J.; Jastrzebska, J.; Filip, M.; Fuxé, K. Disruption of A2AR-D2R Heteroreceptor Complexes After A2AR Transmembrane 5 Peptide Administration Enhances Cocaine Self-Administration in Rats. *Mol. Neurobiol.* **2018**, *55*, 7038–7048. [CrossRef] [PubMed]
204. Borroto-Escuela, D.O.; Wydra, K.; Romero-Fernandez, W.; Zhou, Z.; Frankowska, M.; Filip, M.; Fuxé, K. A2AR Transmembrane 2 Peptide Administration Disrupts the A2AR-A2AR Homoreceptor but Not the A2AR-D2R Heteroreceptor Complex: Lack of Actions on Rodent Cocaine Self-Administration. *Int. J. Mol. Sci.* **2019**, *20*, 6100. [CrossRef] [PubMed]
205. Filip, M.; Frankowska, M.; Zaniewska, M.; Przegalinski, E.; Muller, C.E.; Agnati, L.; Franco, R.; Roberts, D.C.; Fuxé, K. Involvement of Adenosine A2A and Dopamine Receptors in the Locomotor and Sensitizing Effects of Cocaine. *Brain Res.* **2006**, *1077*, 67–80. [CrossRef] [PubMed]
206. Poleszak, E.; Malec, D. Adenosine Receptor Ligands and Cocaine in Conditioned Place Preference (CPP) Test in Rats. *Pol. J. Pharmacol.* **2002**, *54*, 119–126. [PubMed]
207. Poleszak, E.; Malec, D. Effects of Adenosine Receptor Agonists and Antagonists in Amphetamine-Induced Conditioned Place Preference Test in Rats. *Pol. J. Pharmacol.* **2003**, *55*, 319–326.
208. Bachtell, R.K.; Self, D.W. Effects of Adenosine A2A Receptor Stimulation on Cocaine-Seeking Behavior in Rats. *Psychopharmacology* **2009**, *206*, 469–478. [CrossRef] [PubMed]
209. Weerts, E.M.; Griffiths, R.R. The Adenosine Receptor Antagonist CGS15943 Reinstates Cocaine-Seeking Behavior and Maintains Self-Administration in Baboons. *Psychopharmacology* **2003**, *168*, 155–163. [CrossRef] [PubMed]
210. O'Neill, C.E.; LeTendre, M.L.; Bachtell, R.K. Adenosine A2A Receptors in the Nucleus Accumbens Bi-Directionally Alter Cocaine Seeking in Rats. *Neuropsychopharmacology* **2012**, *37*, 1245–1256. [CrossRef]
211. Haynes, N.S.; O'Neill, C.E.; Hobson, B.D.; Bachtell, R.K. Effects of Adenosine A(2A) Receptor Antagonists on Cocaine-Induced Locomotion and Cocaine Seeking. *Psychopharmacology* **2019**, *236*, 699–708. [CrossRef]
212. Orru, M.; Bakesová, J.; Brugarolas, M.; Quiroz, C.; Beaumont, V.; Goldberg, S.R.; Lluís, C.; Cortés, A.; Franco, R.; Casadó, V.; et al. Striatal Pre- and Postsynaptic Profile of Adenosine A(2A) Receptor Antagonists. *PLoS ONE* **2011**, *6*, e16088. [CrossRef] [PubMed]
213. O'Neill, C.E.; Hobson, B.D.; Levis, S.C.; Bachtell, R.K. Persistent Reduction of Cocaine Seeking by Pharmacological Manipulation of Adenosine A1 and A 2A Receptors During Extinction Training in Rats. *Psychopharmacology* **2014**, *231*, 3179–3188. [CrossRef]
214. Marcellino, D.; Roberts, D.C.; Navarro, G.; Filip, M.; Agnati, L.; Lluís, C.; Franco, R.; Fuxé, K. Increase in A2A Receptors in the Nucleus Accumbens After Extended Cocaine Self-Administration and Its Disappearance After Cocaine Withdrawal. *Brain Res.* **2007**, *1143*, 208–220. [CrossRef] [PubMed]
215. Frankowska, M.; Marcellino, D.; Adamczyk, P.; Filip, M.; Fuxé, K. Effects of Cocaine Self-Administration and Extinction on D2 -Like and A2A Receptor Recognition and D2 -Like/Gi Protein Coupling in Rat Striatum. *Addict. Biol.* **2013**, *18*, 455–466. [CrossRef] [PubMed]
216. Kubrusly, R.C.; Bhide, P.G. Cocaine Exposure Modulates Dopamine and Adenosine Signaling in the Fetal Brain. *Neuropharmacology* **2010**, *58*, 436–443. [CrossRef] [PubMed]

217. Romieu, P.; Meunier, J.; Garcia, D.; Zozime, N.; Martin-Fardon, R.; Bowen, W.D.; Maurice, T. The Sigma1 (Sigma1) Receptor Activation Is a Key Step for the Reactivation of Cocaine Conditioned Place Preference by Drug Priming. *Psychopharmacology* **2004**, *175*, 154–162. [CrossRef] [PubMed]
218. Kourrich, S.; Su, T.P.; Fujimoto, M.; Bonci, A. The Sigma-1 Receptor: Roles in Neuronal Plasticity and Disease. *Trends Neurosci.* **2012**, *35*, 762–771. [CrossRef] [PubMed]
219. Pintsuk, J.; Borroto-Escuela, D.O.; Pomierny, B.; Wydra, K.; Zaniewska, M.; Filip, M.; Fuxe, K. Cocaine Self-Administration Differentially Affects Allosteric A2A-D2 Receptor-Receptor Interactions in the Striatum. Relevance for Cocaine Use Disorder. *Pharmacol. Biochem. Behav.* **2016**, *144*, 85–91. [CrossRef] [PubMed]
220. Guidolin, D.; Marcoli, M.; Tortorella, C.; Maura, G.; Agnati, L.F. Adenosine A(2A)-Dopamine D(2) Receptor-Receptor Interaction in Neurons and Astrocytes: Evidence and Perspectives. *Prog. Mol. Biol. Transl. Sci.* **2020**, *169*, 247–277. [PubMed]
221. Marcellino, D.; Navarro, G.; Sahlholm, K.; Nilsson, J.; Agnati, L.F.; Canela, E.I.; Lluís, C.; Århem, P.; Franco, R.; Fuxé, K. Cocaine Produces D2R-Mediated Conformational Changes in the Adenosine A(2A)R-Dopamine D2R Heteromer. *Biochem. Biophys. Res. Commun.* **2010**, *394*, 988–992. [CrossRef]
222. Filip, M.; Zaniewska, M.; Frankowska, M.; Wydra, K.; Fuxé, K. The Importance of the Adenosine A(2A) Receptor-Dopamine D(2) Receptor Interaction in Drug Addiction. *Curr. Med. Chem.* **2012**, *19*, 317–355. [CrossRef]
223. Borroto-Escuela, D.O.; Wydra, K.; Pintsuk, J.; Narvaez, M.; Corrales, F.; Zaniewska, M.; Agnati, L.F.; Franco, R.; Tanganelli, S.; Ferraro, L.; et al. Understanding the Functional Plasticity in Neural Networks of the Basal Ganglia in Cocaine Use Disorder: A Role for Allosteric Receptor-Receptor Interactions in A2A-D2 Heteroreceptor Complexes. *Neural Plast.* **2016**, *2016*, 4827268. [CrossRef] [PubMed]
224. Ballesteros-Yáñez, I.; Castillo, C.A.; Merighi, S.; Gessi, S. The Role of Adenosine Receptors in Psychostimulant Addiction. *Front. Pharmacol.* **2017**, *8*, 985. [CrossRef] [PubMed]
225. Volkow, N.D.; Wang, G.J.; Kollins, S.H.; Wigal, T.L.; Newcorn, J.H.; Telang, F.; Fowler, J.S.; Zhu, W.; Logan, J.; Ma, Y.; et al. Evaluating Dopamine Reward Pathway in ADHD: Clinical Implications. *JAMA* **2009**, *302*, 1084–1091. [CrossRef] [PubMed]
226. Volkow, N.D.; Wang, G.J.; Newcorn, J.H.; Kollins, S.H.; Wigal, T.L.; Telang, F.; Fowler, J.S.; Goldstein, R.Z.; Klein, N.; Logan, J.; et al. Motivation Deficit in ADHD Is Associated With Dysfunction of the Dopamine Reward Pathway. *Mol. Psychiatry* **2011**, *16*, 1147–1154. [CrossRef]
227. Comings, D.E.; Gade-Andavolu, R.; Gonzalez, N.; Wu, S.; Muhleman, D.; Blake, H.; Chiu, F.; Wang, E.; Farwell, K.; Darakjy, S.; et al. Multivariate Analysis of Associations of 42 Genes in ADHD, ODD and Conduct Disorder. *Clin. Genet.* **2000**, *58*, 31–40. [CrossRef] [PubMed]
228. Molero, Y.; Gumpert, C.; Serlachius, E.; Lichtenstein, P.; Walum, H.; Johansson, D.; Anckarsäter, H.; Westberg, L.; Eriksson, E.; Halldner, L. A Study of the Possible Association between Adenosine A2A Receptor Gene Polymorphisms and Attention-Deficit Hyperactivity Disorder Traits. *Genes Brain Behav.* **2013**, *12*, 305–310. [CrossRef]
229. Masuo, Y.; Ishido, M.; Morita, M.; Sawa, H.; Nagashima, K.; Niki, E. Behavioural Characteristics and Gene Expression in the Hyperactive Wiggling (Wig) Rat. *Eur. J. Neurosci.* **2007**, *25*, 3659–3666. [CrossRef]
230. Pandolfo, P.; Machado, N.J.; Köfalvi, A.; Takahashi, R.N.; Cunha, R.A. Caffeine Regulates Frontocorticostratial Dopamine Transporter Density and Improves Attention and Cognitive Deficits in an Animal Model of Attention Deficit Hyperactivity Disorder. *Eur. Neuropsychopharmacol.* **2013**, *23*, 317–328. [CrossRef] [PubMed]
231. Pires, V.A.; Pamplona, F.A.; Pandolfo, P.; Fernandes, D.; Prediger, R.D.; Takahashi, R.N. Adenosine Receptor Antagonists Improve Short-Term Object-Recognition Ability of Spontaneously Hypertensive Rats: A Rodent Model of Attention-Deficit Hyperactivity Disorder. *Behav. Pharmacol.* **2009**, *20*, 134–145. [CrossRef] [PubMed]
232. Alves, C.B.; Almeida, A.S.; Marques, D.M.; Faé, A.H.L.; Machado, A.C.L.; Oliveira, D.L.; Portela, L.V.C.; Porciúncula, L.O. Caffeine and Adenosine A(2A) Receptors Rescue Neuronal Development in Vitro of Frontal Cortical Neurons in a Rat Model of Attention Deficit and Hyperactivity Disorder. *Neuropharmacology* **2020**, *166*, 107782. [CrossRef]
233. Fraporti, T.T.; Contini, V.; Tovo-Rodrigues, L.; Recamonde-Mendoza, M.; Rovaris, D.L.; Rohde, L.A.; Hutz, M.H.; Salatino-Oliveira, A.; Genro, J.P. Synergistic Effects Between ADORA2A and DRD2 Genes on Anxiety Disorders in Children with ADHD. *Prog. Neuropsychopharmacol. Biol. Psychiatry* **2019**, *93*, 214–220. [CrossRef]
234. Hall, H.; Köhler, C.; Gawell, L.; Farde, L.; Sedvall, G. Raclopride, a New Selective Ligand for the Dopamine-D2 Receptors. *Prog. Neuropsychopharmacol. Biol. Psychiatry* **1988**, *12*, 559–568. [CrossRef]
235. Elsinga, P.H.; Hatano, K.; Ishiwata, K. PET Tracers for Imaging of the Dopaminergic System. *Curr. Med. Chem.* **2006**, *13*, 2139–2153. [PubMed]
236. Prante, O.; Maschauer, S.; Banerjee, A. Radioligands for the Dopamine Receptor Subtypes. *J. Labelled Comp. Radiopharm.* **2013**, *56*, 130–148. [CrossRef] [PubMed]
237. Banerjee, A.; Prante, O. Subtype-Selective Dopamine Receptor Radioligands for PET Imaging: Current Status and Recent Developments. *Curr. Med. Chem.* **2012**, *19*, 3957–3966. [CrossRef]
238. Mach, R.H.; Luedtke, R.R. Challenges in the Development of Dopamine D2- and D3-Selective Radiotracers for PET Imaging Studies. *J. Labelled Comp. Radiopharm.* **2018**, *61*, 291–298. [CrossRef]
239. Márián, T.; Boros, I.; Lengyel, Z.; Balkay, L.; Horváth, G.; Emri, M.; Sarkadi, E.; Szentmiklósi, A.J.; Fekete, I.; Trón, L. Preparation and Primary Evaluation of [11C]CSC As a Possible Tracer for Mapping Adenosine A2A Receptors by PET. *Appl. Radiat. Isot.* **1999**, *50*, 887–893. [CrossRef]

240. Ishiwata, K.; Shimada, J.; Wang, W.F.; Harakawa, H.; Ishii, S.; Kiyosawa, M.; Suzuki, F.; Senda, M. Evaluation of Iodinated and Brominated [11C]Styrylxanthine Derivatives As in Vivo Radioligands Mapping Adenosine A2A Receptor in the Central Nervous System. *Ann. Nucl. Med.* **2000**, *14*, 247–253. [[CrossRef](#)] [[PubMed](#)]
241. Lowe, P.T.; Dall’Angelo, S.; Mulder-Krieger, T.; IJzerman, A.P.; Zanda, M.; O’Hagan, D. A New Class of Fluorinated A(2A) Adenosine Receptor Agonist With Application to Last-Step Enzymatic [(18)F]Fluorination for PET Imaging. *Chembiochem* **2017**, *18*, 2156–2164. [[CrossRef](#)] [[PubMed](#)]
242. Bhattacharjee, A.K.; Lang, L.; Jacobson, O.; Shinkre, B.; Ma, Y.; Niu, G.; Trenkle, W.C.; Jacobson, K.A.; Chen, X.; Kiesewetter, D.O. Striatal Adenosine A(2A) Receptor-Mediated Positron Emission Tomographic Imaging in 6-Hydroxydopamine-Lesioned Rats Using [(18)F]-MRS5425. *Nucl. Med. Biol.* **2011**, *38*, 897–906. [[CrossRef](#)] [[PubMed](#)]
243. Schröder, S.; Lai, T.H.; Toussaint, M.; Kranz, M.; Chovsepian, A.; Shang, Q.; Dukic-Stefanovic, S.; Deuther-Conrad, W.; Teodoro, R.; Wenzel, B.; et al. PET Imaging of the Adenosine A(2A) Receptor in the Rotenone-Based Mouse Model of Parkinson’s Disease With [(18)F]FESCH Synthesized by a Simplified Two-Step One-Pot Radiolabeling Strategy. *Molecules* **2020**, *25*, 1633. [[CrossRef](#)] [[PubMed](#)]
244. Khanapur, S.; Paul, S.; Shah, A.; Vatakuti, S.; Koole, M.J.; Zijlma, R.; Dierckx, R.A.; Luurtsema, G.; Garg, P.; Van Waarde, A.; et al. Development of [18F]-Labeled Pyrazolo[4,3-e]-1,2,4-Triazolo[1,5-c]Pyrimidine (SCH442416) Analogs for the Imaging of Cerebral Adenosine A2A Receptors With Positron Emission Tomography. *J. Med. Chem.* **2014**, *57*, 6765–6780. [[CrossRef](#)] [[PubMed](#)]
245. Khanapur, S.; Van Waarde, A.; Dierckx, R.A.; Elsinga, P.H.; Koole, M.J. Preclinical Evaluation and Quantification of (18)F-Fluoroethyl and (18)F-Fluoropropyl Analogs of SCH442416 As Radioligands for PET Imaging of the Adenosine A(2A) Receptor in Rat Brain. *J. Nucl. Med.* **2017**, *58*, 466–472. [[CrossRef](#)] [[PubMed](#)]
246. Hirani, E.; Gillies, J.; Karasawa, A.; Shimada, J.; Kase, H.; Opacka-Juffry, J.; Osman, S.; Luthra, S.K.; Hume, S.P.; Brooks, D.J. Evaluation of [4-O-Methyl-(11)C]KW-6002 As a Potential PET Ligand for Mapping Central Adenosine A(2A) Receptors in Rats. *Synapse* **2001**, *42*, 164–176. [[CrossRef](#)] [[PubMed](#)]
247. Brooks, D.J.; Doder, M.; Osman, S.; Luthra, S.K.; Hirani, E.; Hume, S.; Kase, H.; Kilborn, J.; Martindill, S.; Mori, A. Positron Emission Tomography Analysis of [11C]KW-6002 Binding to Human and Rat Adenosine A2A Receptors in the Brain. *Synapse* **2008**, *62*, 671–681. [[CrossRef](#)]
248. Ishiwata, K.; Noguchi, J.; Toyama, H.; Sakiyama, Y.; Koike, N.; Ishii, S.; Oda, K.; Endo, K.; Suzuki, F.; Senda, M. Synthesis and Preliminary Evaluation of [11C]KF17837, a Selective Adenosine A2A Antagonist. *Appl. Radiat. Isot.* **1996**, *47*, 507–511. [[CrossRef](#)]
249. Ishiwata, K.; Sakiyama, Y.; Sakiyama, T.; Shimada, J.; Toyama, H.; Oda, K.; Suzuki, F.; Senda, M. Myocardial Adenosine A2a Receptor Imaging of Rabbit by PET With [11C]KF17837. *Ann. Nucl. Med.* **1997**, *11*, 219–225. [[CrossRef](#)]
250. Noguchi, J.; Ishiwata, K.; Wakabayashi, S.; Nariai, T.; Shumiya, S.; Ishii, S.; Toyama, H.; Endo, K.; Suzuki, F.; Senda, M. Evaluation of Carbon-11-Labeled KF17837: A Potential CNS Adenosine A2a Receptor Ligand. *J. Nucl. Med.* **1998**, *39*, 498–503. [[PubMed](#)]
251. Stone-Elander, S.; Thorell, J.O.; Eriksson, L.; Fredholm, B.B.; Ingvar, M. In Vivo Biodistribution of [N-11C-Methyl]KF 17837 Using 3-D-PET: Evaluation As a Ligand for the Study of Adenosine A2A Receptors. *Nucl. Med. Biol.* **1997**, *24*, 187–191. [[CrossRef](#)]
252. Ishiwata, K.; Noguchi, J.; Wakabayashi, S.; Shimada, J.; Ogi, N.; Nariai, T.; Tanaka, A.; Endo, K.; Suzuki, F.; Senda, M. 11C-Labeled KF18446: A Potential Central Nervous System Adenosine A2a Receptor Ligand. *J. Nucl. Med.* **2000**, *41*, 345–354. [[PubMed](#)]
253. Ishiwata, K.; Ogi, N.; Shimada, J.; Nonaka, H.; Tanaka, A.; Suzuki, F.; Senda, M. Further Characterization of a CNS Adenosine A2a Receptor Ligand [11C]KF18446 with in Vitro Autoradiography and in Vivo Tissue Uptake. *Ann. Nucl. Med.* **2000**, *14*, 81–89. [[CrossRef](#)] [[PubMed](#)]
254. Ishiwata, K.; Ogi, N.; Shimada, J.; Wang, W.; Ishii, K.; Tanaka, A.; Suzuki, F.; Senda, M. Search for PET Probes for Imaging the Globus Pallidus Studied With Rat Brain Ex Vivo Autoradiography. *Ann. Nucl. Med.* **2000**, *14*, 461–466. [[CrossRef](#)] [[PubMed](#)]
255. Ishiwata, K.; Ogi, N.; Hayakawa, N.; Oda, K.; Nagaoka, T.; Toyama, H.; Suzuki, F.; Endo, K.; Tanaka, A.; Senda, M. Adenosine A2A Receptor Imaging With [11C]KF18446 PET in the Rat Brain After Quinolinic Acid Lesion: Comparison With the Dopamine Receptor Imaging. *Ann. Nucl. Med.* **2002**, *16*, 467–475. [[CrossRef](#)] [[PubMed](#)]
256. Ishiwata, K.; Wang, W.F.; Kimura, Y.; Kawamura, K.; Ishii, K. Preclinical Studies on [11C]TMSX for Mapping Adenosine A2A Receptors by Positron Emission Tomography. *Ann. Nucl. Med.* **2003**, *17*, 205–211. [[CrossRef](#)] [[PubMed](#)]
257. Ishiwata, K.; Mizuno, M.; Kimura, Y.; Kawamura, K.; Oda, K.; Sasaki, T.; Nakamura, Y.; Muraoka, I.; Ishii, K. Potential of [11C]TMSX for the Evaluation of Adenosine A2A Receptors in the Skeletal Muscle by Positron Emission Tomography. *Nucl. Med. Biol.* **2004**, *31*, 949–956. [[CrossRef](#)] [[PubMed](#)]
258. Ishiwata, K.; Kawamura, K.; Kimura, Y.; Oda, K.; Ishii, K. Potential of an Adenosine A2A Receptor Antagonist [11C]TMSX for Myocardial Imaging by Positron Emission Tomography: A First Human Study. *Ann. Nucl. Med.* **2003**, *17*, 457–462. [[CrossRef](#)] [[PubMed](#)]
259. Ishiwata, K.; Mishina, M.; Kimura, Y.; Oda, K.; Sasaki, T.; Ishii, K. First Visualization of Adenosine A(2A) Receptors in the Human Brain by Positron Emission Tomography With [11C]TMSX. *Synapse* **2005**, *55*, 133–136. [[CrossRef](#)]
260. Mizuno, M.; Kimura, Y.; Tokizawa, K.; Ishii, K.; Oda, K.; Sasaki, T.; Nakamura, Y.; Muraoka, I.; Ishiwata, K. Greater Adenosine A(2A) Receptor Densities in Cardiac and Skeletal Muscle in Endurance-Trained Men: A [11C]TMSX PET Study. *Nucl. Med. Biol.* **2005**, *32*, 831–836. [[CrossRef](#)]
261. Mishina, M.; Ishiwata, K.; Kimura, Y.; Naganawa, M.; Oda, K.; Kobayashi, S.; Katayama, Y.; Ishii, K. Evaluation of Distribution of Adenosine A2A Receptors in Normal Human Brain Measured With [11C]TMSX PET. *Synapse* **2007**, *61*, 778–784. [[CrossRef](#)]

262. Mishina, M.; Ishiwata, K.; Naganawa, M.; Kimura, Y.; Kitamura, S.; Suzuki, M.; Hashimoto, M.; Ishibashi, K.; Oda, K.; Sakata, M.; et al. Adenosine A(2A) Receptors Measured With [<sup>11</sup>C]TMSX PET in the Striata of Parkinson's Disease Patients. *PLoS ONE* **2011**, *6*, e17338. [CrossRef] [PubMed]
263. Heinonen, I.; Nesterov, S.V.; Liukko, K.; Kemppainen, J.; Någren, K.; Luotolahti, M.; Virsu, P.; Oikonen, V.; Nuutila, P.; Kujala, U.M.; et al. Myocardial Blood Flow and Adenosine A2A Receptor Density in Endurance Athletes and Untrained Men. *J. Physiol.* **2008**, *586*, 5193–5202. [CrossRef]
264. Rissanen, E.; Virta, J.R.; Paavilainen, T.; Tuisku, J.; Helin, S.; Luoto, P.; Parkkola, R.; Rinne, J.O.; Airas, L. Adenosine A2A Receptors in Secondary Progressive Multiple Sclerosis: A [(<sup>11</sup>C)]TMSX Brain PET Study. *J. Cereb. Blood Flow Metab.* **2013**, *33*, 1394–1401. [CrossRef] [PubMed]
265. Rissanen, E.; Tuisku, J.; Luoto, P.; Arponen, E.; Johansson, J.; Oikonen, V.; Parkkola, R.; Airas, L.; Rinne, J.O. Automated Reference Region Extraction and Population-Based Input Function for Brain [(<sup>11</sup>C)]TMSX PET Image Analyses. *J. Cereb. Blood Flow Metab.* **2015**, *35*, 157–165. [CrossRef]
266. Naganawa, M.; Kimura, Y.; Mishina, M.; Manabe, Y.; Chihara, K.; Oda, K.; Ishii, K.; Ishiwata, K. Quantification of Adenosine A2A Receptors in the Human Brain Using [<sup>11</sup>C]TMSX and Positron Emission Tomography. *Eur. J. Nucl. Mol. Imaging* **2007**, *34*, 679–687. [CrossRef] [PubMed]
267. Naganawa, M.; Kimura, Y.; Yano, J.; Mishina, M.; Yanagisawa, M.; Ishii, K.; Oda, K.; Ishiwata, K. Robust Estimation of the Arterial Input Function for Logan Plots Using an Intersectional Searching Algorithm and Clustering in Positron Emission Tomography for Neuroreceptor Imaging. *Neuroimage* **2008**, *40*, 26–34. [CrossRef]
268. Naganawa, M.; Mishina, M.; Sakata, M.; Oda, K.; Hiura, M.; Ishii, K.; Ishiwata, K. Test-Retest Variability of Adenosine A2A Binding in the Human Brain With (<sup>11</sup>C)-TMSX and PET. *EJNMMI Res* **2014**, *4*, 76. [CrossRef] [PubMed]
269. Mishina, M.; Kimura, Y.; Naganawa, M.; Ishii, K.; Oda, K.; Sakata, M.; Toyohara, J.; Kobayashi, S.; Katayama, Y.; Ishiwata, K. Differential Effects of Age on Human Striatal Adenosine A1 and A2A Receptors. *Synapse* **2012**, *66*, 832–839. [CrossRef]
270. Lahesmaa, M.; Oikonen, V.; Helin, S.; Luoto, P.; Din, U.; Pfeifer, A.; Nuutila, P.; Virtanen, K.A. Regulation of Human Brown Adipose Tissue by Adenosine and A(2A) Receptors Studies with [(<sup>15</sup>O)]H<sub>2</sub>O and [(<sup>11</sup>C)]TMSX PET/CT. *Eur. J. Nucl. Mol. Imaging* **2019**, *46*, 743–750. [CrossRef]
271. Wang, W.F.; Ishiwata, K.; Nonaka, H.; Ishii, S.; Kiyosawa, M.; Shimada, J.; Suzuki, F.; Senda, M. Carbon-11-Labeled KF21213: A Highly Selective Ligand for Mapping CNS Adenosine A(2A) Receptors With Positron Emission Tomography. *Nucl. Med. Biol.* **2000**, *27*, 541–546. [CrossRef]
272. Barret, O.; Hannestad, J.; Alagille, D.; Vala, C.; Tavares, A.; Papin, C.; Morley, T.; Fowles, K.; Lee, H.; Seibyl, J.; et al. Adenosine 2A Receptor Occupancy by Tozadenant and Preladenant in Rhesus Monkeys. *J. Nucl. Med.* **2014**, *55*, 1712–1718. [CrossRef]
273. Vala, C.; Morley, T.J.; Zhang, X.; Papin, C.; Tavares, A.A.; Lee, H.S.; Constantinescu, C.; Barret, O.; Carroll, V.M.; Baldwin, R.M.; et al. Synthesis and in Vivo Evaluation of Fluorine-18 and Iodine-123 Pyrazolo[4,3-e]-1,2,4-Triazolo[1,5-c] Pyrimidine Derivatives As PET and SPECT Radiotracers for Mapping A2A Receptors. *ChemMedChem* **2016**, *11*, 1936–1943. [CrossRef] [PubMed]
274. Barret, O.; Hannestad, J.; Vala, C.; Alagille, D.; Tavares, A.; Laruelle, M.; Jennings, D.; Marek, K.; Russell, D.; Seibyl, J.; et al. Characterization in Humans of <sup>18</sup>F-MNI-444, a PET Radiotracer for Brain Adenosine 2A Receptors. *J. Nucl. Med.* **2015**, *56*, 586–591. [CrossRef] [PubMed]
275. Zhou, X.; Khanapur, S.; Huizing, A.P.; Zijlma, R.; Schepers, M.; Dierckx, R.A.; Van Waarde, A.; de Vries, E.F.; Elsinga, P.H. Synthesis and Preclinical Evaluation of 2-(2-Furanyl)-7-[2-[4-[4-(2-[<sup>11</sup>C]Methoxyethoxy)Phenyl]-1-Piperazinyl]Ethyl]-7H-Pyrazolo[4,3-e][1,2,4]Triazolo[1,5-c]Pyrimidine-5-Amine ([<sup>11</sup>C]Preladenant) As a PET Tracer for the Imaging of Cerebral Adenosine A2A Receptors. *J. Med. Chem.* **2014**, *57*, 9204–9210. [PubMed]
276. Zhou, X.; Elsinga, P.H.; Khanapur, S.; Dierckx, R.A.; de Vries, E.F.; de Jong, J.R. Radiation Dosimetry of a Novel Adenosine A(2A) Receptor Radioligand [(<sup>11</sup>C)]Preladenant Based on PET/CT Imaging and Ex Vivo Biodistribution in Rats. *Mol. Imaging Biol.* **2017**, *19*, 289–297. [CrossRef] [PubMed]
277. Zhou, X.; Khanapur, S.; de Jong, J.R.; Willemse, A.T.; Dierckx, R.A.; Elsinga, P.H.; de Vries, E.F. In Vivo Evaluation of [(<sup>11</sup>C)]Preladenant Positron Emission Tomography for Quantification of Adenosine A(2A) Receptors in the Rat Brain. *J. Cereb. Blood Flow Metab.* **2017**, *37*, 577–589. [CrossRef] [PubMed]
278. Zhou, X.; Doorduin, J.; Elsinga, P.H.; Dierckx, R.A.J.O.; de Vries, E.F.J.; Casteels, C. Altered Adenosine 2A and Dopamine D2 Receptor Availability in the 6-Hydroxydopamine-Treated Rats With and Without Levodopa-Induced Dyskinesia. *Neuroimage* **2017**, *157*, 209–218. [CrossRef] [PubMed]
279. Zhou, X.; Boellaard, R.; Ishiwata, K.; Sakata, M.; Dierckx, R.A.J.O.; de Jong, J.R.; Nishiyama, S.; Ohba, H.; Tsukada, H.; de Vries, E.F.J.; et al. In Vivo Evaluation of (<sup>11</sup>C)-Preladenant for PET Imaging of Adenosine A(2A) Receptors in the Conscious Monkey. *J. Nucl. Med.* **2017**, *58*, 762–767. [CrossRef]
280. Ishibashi, K.; Miura, Y.; Wagatsuma, K.; Toyohara, J.; Ishiwata, K.; Ishii, K. Occupancy of Adenosine A(2A) Receptors by Istradefylline in Patients With Parkinson's Disease Using (<sup>11</sup>C)-Preladenant PET. *Neuropharmacology* **2018**, *143*, 106–112. [CrossRef]
281. Sakata, M.; Ishibashi, K.; Imai, M.; Wagatsuma, K.; Ishii, K.; Zhou, X.; de Vries, E.F.J.; Elsinga, P.H.; Ishiwata, K.; Toyohara, J. Initial Evaluation of an Adenosine A(2A) Receptor Ligand, (<sup>11</sup>C)-Preladenant, in Healthy Human Subjects. *J. Nucl. Med.* **2017**, *58*, 1464–1470. [CrossRef]

282. Moresco, R.M.; Todde, S.; Belloli, S.; Simonelli, P.; Panzacchi, A.; Rigamonti, M.; Galli-Kienle, M.; Fazio, F. In Vivo Imaging of Adenosine A2A Receptors in Rat and Primate Brain Using [11C]SCH442416. *Eur. J. Nucl. Med. Mol. Imaging* **2005**, *32*, 405–413. [[CrossRef](#)]
283. Todde, S.; Moresco, R.M.; Simonelli, P.; Baraldi, P.G.; Cacciari, B.; Spalluto, G.; Varani, K.; Monopoli, A.; Matarrese, M.; Carpinelli, A.; et al. Design, Radiosynthesis, and Biodistribution of a New Potent and Selective Ligand for in Vivo Imaging of the Adenosine A(2A) Receptor System Using Positron Emission Tomography. *J. Med. Chem.* **2000**, *43*, 4359–4362. [[CrossRef](#)]
284. Mihara, T.; Noda, A.; Arai, H.; Mihara, K.; Iwashita, A.; Murakami, Y.; Matsuya, T.; Miyoshi, S.; Nishimura, S.; Matsuoka, N. Brain Adenosine A2A Receptor Occupancy by a Novel A1/A2A Receptor Antagonist, ASP5854, in Rhesus Monkeys: Relationship to Anticataleptic Effect. *J. Nucl. Med.* **2008**, *49*, 1183–1188. [[CrossRef](#)]
285. Ramlackhansingh, A.F.; Bose, S.K.; Ahmed, I.; Turkheimer, F.E.; Pavese, N.; Brooks, D.J. Adenosine 2A Receptor Availability in Dyskinetic and Nondyskinetic Patients With Parkinson Disease. *Neurology* **2011**, *76*, 1811–1816. [[CrossRef](#)]
286. Brooks, D.J.; Papapetropoulos, S.; Vandenbende, F.; Tomic, D.; He, P.; Coppell, A.; O'Neill, G. An Open-Label, Positron Emission Tomography Study to Assess Adenosine A2A Brain Receptor Occupancy of Vipadenant (BIIB014) at Steady-State Levels in Healthy Male Volunteers. *Clin. Neuropharmacol.* **2010**, *33*, 55–60. [[CrossRef](#)]
287. Arnett, C.D.; Shue, C.Y.; Wolf, A.P.; Fowler, J.S.; Logan, J.; Watanabe, M. Comparison of Three 18F-Labeled Butyrophenone Neuroleptic Drugs in the Baboon Using Positron Emission Tomography. *J. Neurochem.* **1985**, *44*, 835–844. [[CrossRef](#)]
288. Moerlein, S.M.; Perlmutter, J.S.; Welch, M.J. Specific, Reversible Binding of [18F]Benperidol to Baboon D2 Receptors: PET Evaluation of an Improved 18F-Labeled Ligand. *Nucl. Med. Biol.* **1995**, *22*, 809–815. [[CrossRef](#)]
289. Döbrössy, M.D.; Braun, F.; Klein, S.; Garcia, J.; Langen, K.J.; Weber, W.A.; Nikkhah, G.; Meyer, P.T. [18F]Desmethoxyfallypride As a Novel PET Radiotracer for Quantitative in Vivo Dopamine D2/D3 Receptor Imaging in Rat Models of Neurodegenerative Diseases. *Nucl. Med. Biol.* **2012**, *39*, 1077–1080. [[CrossRef](#)] [[PubMed](#)]
290. Mille, E.; Cumming, P.; Rominger, A.; La, F.C.; Tatsch, K.; Wängler, B.; Bartenstein, P.; Böning, G. Compensation for Cranial Spill-in into the Cerebellum Improves Quantitation of Striatal Dopamine D2/3 Receptors in Rats With Prolonged [18F]-DMFP Infusions. *Synapse* **2012**, *66*, 705–713. [[CrossRef](#)] [[PubMed](#)]
291. Rominger, A.; Wagner, E.; Mille, E.; Böning, G.; Esmaeilzadeh, M.; Wängler, B.; Gildehaus, F.J.; Nowak, S.; Bruche, A.; Tatsch, K.; et al. Endogenous Competition Against Binding of [(18)F]DMFP and [(18)F]Fallypride to Dopamine D(2/3) Receptors in Brain of Living Mouse. *Synapse* **2010**, *64*, 313–322. [[CrossRef](#)] [[PubMed](#)]
292. Heinz, A.; Siessmeier, T.; Wräse, J.; Buchholz, H.G.; Gründer, G.; Kumakura, Y.; Cumming, P.; Schreckenberger, M.; Smolka, M.N.; Röscher, F.; et al. Correlation of Alcohol Craving With Striatal Dopamine Synthesis Capacity and D2/3 Receptor Availability: A Combined [18F]DOPA and [18F]DMFP PET Study in Detoxified Alcoholic Patients. *Am. J. Psychiatry* **2005**, *162*, 1515–1520. [[CrossRef](#)]
293. Schreckenberger, M.; Hägele, S.; Siessmeier, T.; Buchholz, H.G.; Armbrust-Henrich, H.; Röscher, F.; Gründer, G.; Bartenstein, P.; Vogt, T. The Dopamine D2 Receptor Ligand 18F-Desmethoxyfallypride: An Appropriate Fluorinated PET Tracer for the Differential Diagnosis of Parkinsonism. *Eur. J. Nucl. Med. Mol. Imaging* **2004**, *31*, 1128–1135. [[CrossRef](#)]
294. Halldin, C.; Farde, L.; Höglberg, T.; Hall, H.; Sedvall, G. Carbon-11 Labelling of Eticlopride in Two Different Positions—a Selective High-Affinity Ligand for the Study of Dopamine D-2 Receptors Using PET. *Int. J. Rad. Appl. Instrum. A* **1990**, *41*, 669–674. [[CrossRef](#)]
295. Mukherjee, J.; Shi, B.; Christian, B.T.; Chattopadhyay, S.; Narayanan, T.K. 11C-Fallypride: Radiosynthesis and Preliminary Evaluation of a Novel Dopamine D2/D3 Receptor PET Radiotracer in Non-Human Primate Brain. *Bioorg. Med. Chem.* **2004**, *12*, 95–102. [[CrossRef](#)] [[PubMed](#)]
296. Narendran, R.; Frankle, W.G.; Mason, N.S.; Rabiner, E.A.; Gunn, R.N.; Searle, G.E.; Vora, S.; Litschge, M.; Kendro, S.; Cooper, T.B.; et al. Positron Emission Tomography Imaging of Amphetamine-Induced Dopamine Release in the Human Cortex: A Comparative Evaluation of the High Affinity Dopamine D2/3 Radiotracers [11C]FLB 457 and [11C]Fallypride. *Synapse* **2009**, *63*, 447–461. [[CrossRef](#)] [[PubMed](#)]
297. Mukherjee, J.; Yang, Z.Y.; Brown, T.; Lew, R.; Wernick, M.; Ouyang, X.; Yasillo, N.; Chen, C.T.; Mintzer, R.; Cooper, M. Preliminary Assessment of Extrastriatal Dopamine D-2 Receptor Binding in the Rodent and Nonhuman Primate Brains Using the High Affinity Radioligand, 18F-Fallypride. *Nucl. Med. Biol.* **1999**, *26*, 519–527. [[CrossRef](#)]
298. Vuckovic, M.G.; Li, Q.; Fisher, B.; Nacca, A.; Leahy, R.M.; Walsh, J.P.; Mukherjee, J.; Williams, C.; Jakowec, M.W.; Petzinger, G.M. Exercise Elevates Dopamine D2 Receptor in a Mouse Model of Parkinson’s Disease: In Vivo Imaging With [18F]Fallypride. *Mov. Disord.* **2010**, *25*, 2777–2784. [[CrossRef](#)] [[PubMed](#)]
299. Ota, M.; Ogawa, S.; Kato, K.; Masuda, C.; Kunugi, H. Striatal and Extrastriatal Dopamine Release in the Common Marmoset Brain Measured by Positron Emission Tomography and [(18)F]Fallypride. *Neurosci. Res.* **2015**, *101*, 1–5. [[CrossRef](#)]
300. Mukherjee, J.; Yang, Z.Y.; Lew, R.; Brown, T.; Kronmal, S.; Cooper, M.D.; Seiden, L.S. Evaluation of D-Amphetamine Effects on the Binding of Dopamine D-2 Receptor Radioligand, 18F-Fallypride in Nonhuman Primates Using Positron Emission Tomography. *Synapse* **1997**, *27*, 1–13. [[CrossRef](#)]
301. Slifstein, M.; Narendran, R.; Hwang, D.R.; Sudo, Y.; Talbot, P.S.; Huang, Y.; Laruelle, M. Effect of Amphetamine on [(18)F]Fallypride in Vivo Binding to D(2) Receptors in Striatal and Extrastriatal Regions of the Primate Brain: Single Bolus and Bolus Plus Constant Infusion Studies. *Synapse* **2004**, *54*, 46–63. [[CrossRef](#)] [[PubMed](#)]

302. Slifstein, M.; Hwang, D.R.; Huang, Y.; Guo, N.; Sudo, Y.; Narendran, R.; Talbot, P.; Laruelle, M. In Vivo Affinity of [18F]Fallypride for Striatal and Extrastriatal Dopamine D2 Receptors in Nonhuman Primates. *Psychopharmacology* **2004**, *175*, 274–286. [[CrossRef](#)] [[PubMed](#)]
303. Mukherjee, J.; Christian, B.T.; Narayanan, T.K.; Shi, B.; Collins, D. Measurement of D-Amphetamine-Induced Effects on the Binding of Dopamine D-2/D-3 Receptor Radioligand, 18F-Fallypride in Extrastriatal Brain Regions in Non-Human Primates Using PET. *Brain Res.* **2005**, *1032*, 77–84. [[CrossRef](#)] [[PubMed](#)]
304. Naylor, J.E.; Hiranita, T.; Matazel, K.S.; Zhang, X.; Paule, M.G.; Goodwin, A.K. Positron Emission Tomography (PET) Imaging of Nicotine-Induced Dopamine Release in Squirrel Monkeys Using [(18)F]Fallypride. *Drug Alcohol Depend.* **2017**, *179*, 254–259. [[CrossRef](#)] [[PubMed](#)]
305. Riccardi, P.; Li, R.; Ansari, M.S.; Zald, D.; Park, S.; Dawant, B.; Anderson, S.; Doop, M.; Woodward, N.; Schoenberg, E.; et al. Amphetamine-Induced Displacement of [18F] Fallypride in Striatum and Extrastriatal Regions in Humans. *Neuropsychopharmacology* **2006**, *31*, 1016–1026. [[CrossRef](#)]
306. Riccardi, P.; Zald, D.; Li, R.; Park, S.; Ansari, M.S.; Dawant, B.; Anderson, S.; Woodward, N.; Schmidt, D.; Baldwin, R.; et al. Sex Differences in Amphetamine-Induced Displacement of [(18)F]Fallypride in Striatal and Extrastriatal Regions: A PET Study. *Am. J. Psychiatry* **2006**, *163*, 1639–1641. [[CrossRef](#)]
307. Cropley, V.L.; Innis, R.B.; Nathan, P.J.; Brown, A.K.; Sangare, J.L.; Lerner, A.; Ryu, Y.H.; Sprague, K.E.; Pike, V.W.; Fujita, M. Small Effect of Dopamine Release and No Effect of Dopamine Depletion on [18F]Fallypride Binding in Healthy Humans. *Synapse* **2008**, *62*, 399–408. [[CrossRef](#)]
308. Lehrer, D.S.; Christian, B.T.; Kirbas, C.; Chiang, M.; Sidhu, S.; Short, H.; Wang, B.; Shi, B.; Chu, K.W.; Merrill, B.; et al. 18F-Fallypride Binding Potential in Patients With Schizophrenia Compared to Healthy Controls. *Schizophr. Res.* **2010**, *122*, 43–52. [[CrossRef](#)]
309. Slifstein, M.; Kegeles, L.S.; Xu, X.; Thompson, J.L.; Urban, N.; Castrillon, J.; Hackett, E.; Bae, S.A.; Laruelle, M.; bi-Dargham, A. Striatal and Extrastriatal Dopamine Release Measured With PET and [(18)F] Fallypride. *Synapse* **2010**, *64*, 350–362. [[CrossRef](#)]
310. Stark, A.J.; Smith, C.T.; Petersen, K.J.; Trujillo, P.; van Wouwe, N.C.; Donahue, M.J.; Kessler, R.M.; Deutch, A.Y.; Zald, D.H.; Claassen, D.O. [(18)F]Fallypride Characterization of Striatal and Extrastriatal D(2/3) Receptors in Parkinson’s Disease. *Neuroimage Clin.* **2018**, *18*, 433–442. [[CrossRef](#)]
311. Joo, Y.H.; Kim, J.H.; Son, Y.D.; Kim, H.K.; Shin, Y.J.; Lee, S.Y.; Kim, J.H. The Relationship Between Excitement Symptom Severity and Extrastriatal Dopamine D(2/3) Receptor Availability in Patients With Schizophrenia: A High-Resolution PET Study With [(18)F]Fallypride. *Eur. Arch. Psychiatry Clin. Neurosci.* **2018**, *268*, 529–540. [[CrossRef](#)]
312. Mach, R.H.; Nader, M.A.; Ehrenkaufer, R.L.; Line, S.W.; Smith, C.R.; Luedtke, R.R.; Kung, M.P.; Kung, H.F.; Lyons, D.; Morton, T.E. Comparison of Two Fluorine-18 Labeled Benzamide Derivatives That Bind Reversibly to Dopamine D2 Receptors: In Vitro Binding Studies and Positron Emission Tomography. *Synapse* **1996**, *24*, 322–333. [[CrossRef](#)]
313. Takao, F.; Sasaki, S.; Maeda, M.; Fukumura, T.; Tahara, T.; Masuda, K.; Ichiya, Y. Synthesis and in Vivo Evaluation of a New Fluorine-18 Labeled Dopamine D2 Radioligand with Benzofuran Benzamide Skeleton. *J. Labelled Comp. Radiopharm.* **1993**, *33*, 1107–1112. [[CrossRef](#)]
314. Barrio, J.R.; Satyamurthy, N.; Huang, S.C.; Keen, R.E.; Nissenson, C.H.; Hoffman, J.M.; Ackermann, R.F.; Bahn, M.M.; Mazziotta, J.C.; Phelps, M.E. 3-(2’-[18F]Fluoroethyl) Spiperone: In Vivo Biochemical and Kinetic Characterization in Rodents, Nonhuman Primates, and Humans. *J. Cereb. Blood Flow Metab.* **1989**, *9*, 830–839. [[CrossRef](#)] [[PubMed](#)]
315. Coenen, H.H.; Laufer, P.; Stöcklin, G.; Wienhard, K.; Pawlik, G.; Böcher-Schwarz, H.G.; Heiss, W.D. 3-N-(2-[18F]-Fluoroethyl)-Spiperone: A Novel Ligand for Cerebral Dopamine Receptor Studies With PET. *Life Sci.* **1987**, *40*, 81–88. [[CrossRef](#)]
316. Satyamurthy, N.; Bida, G.T.; Barrio, J.R.; Luxen, A.; Mazziotta, J.C.; Huang, S.C.; Phelps, M.E. No-Carrier-Added 3-(2’-[18F]Fluoroethyl)Spiperone, a New Dopamine Receptor-Binding Tracer for Positron Emission Tomography. *Int. J. Rad. Appl. Instrum. B* **1986**, *13*, 617–624. [[CrossRef](#)]
317. Satyamurthy, N.; Barrio, J.R.; Bida, G.T.; Huang, S.C.; Mazziotta, J.C.; Phelps, M.E. 3-(2’-[18F]Fluoroethyl)Spiperone, a Potent Dopamine Antagonist: Synthesis, Structural Analysis and in-Vivo Utilization in Humans. *Int. J. Rad. Appl. Instrum. A* **1990**, *41*, 113–129. [[CrossRef](#)]
318. Aung, W.; Okauchi, T.; Sato, M.; Saito, T.; Nakagawa, H.; Ishihara, H.; Ikota, N.; Suhara, T.; Anzai, K. In-Vivo PET Imaging of Inducible D2R Reporter Transgene Expression Using [11C]FLB 457 As Reporter Probe in Living Rats. *Nucl. Med. Commun.* **2005**, *26*, 259–268. [[CrossRef](#)]
319. Halldin, C.; Farde, L.; Höglberg, T.; Mohell, N.; Hall, H.; Suhara, T.; Karlsson, P.; Nakashima, Y.; Swahn, C.G. Carbon-11-FLB 457: A Radioligand for Extrastriatal D2 Dopamine Receptors. *J. Nucl. Med.* **1995**, *36*, 1275–1281. [[PubMed](#)]
320. Narendran, R.; Jedema, H.P.; Lopresti, B.J.; Mason, N.S.; Gurnsey, K.; Ruszkiewicz, J.; Chen, C.M.; Deutitch, L.; Frankle, W.G.; Bradberry, C.W. Imaging Dopamine Transmission in the Frontal Cortex: A Simultaneous Microdialysis and [11C]FLB 457 PET Study. *Mol. Psychiatry* **2014**, *19*, 302–310. [[CrossRef](#)] [[PubMed](#)]
321. Olsson, H.; Halldin, C.; Swahn, C.G.; Farde, L. Quantification of [11C]FLB 457 Binding to Extrastriatal Dopamine Receptors in the Human Brain. *J. Cereb. Blood Flow Metab.* **1999**, *19*, 1164–1173. [[CrossRef](#)]
322. Vilkman, H.; Kajander, J.; Någren, K.; Oikonen, V.; Syvälahti, E.; Hietala, J. Measurement of Extrastriatal D2-Like Receptor Binding With [11C]FLB 457—A Test-Retest Analysis. *Eur. J. Nucl. Med.* **2000**, *27*, 1666–1673. [[CrossRef](#)]

323. Ito, H.; Sudo, Y.; Suhara, T.; Okubo, Y.; Halldin, C.; Farde, L. Error Analysis for Quantification of [(11)C]FLB 457 Binding to Extrastriatal D(2) Dopamine Receptors in the Human Brain. *Neuroimage* **2001**, *13*, 531–539. [CrossRef]
324. Sudo, Y.; Suhara, T.; Inoue, M.; Ito, H.; Suzuki, K.; Saijo, T.; Halldin, C.; Farde, L. Reproducibility of [11 C]FLB 457 Binding in Extrastriatal Regions. *Nucl. Med. Commun.* **2001**, *22*, 1215–1221. [CrossRef]
325. Talvik, M.; Nordström, A.L.; Olsson, H.; Halldin, C.; Farde, L. Decreased Thalamic D2/D3 Receptor Binding in Drug-Naïve Patients With Schizophrenia: A PET Study With [11C]FLB 457. *Int. J. Neuropsychopharmacol.* **2003**, *6*, 361–370. [CrossRef]
326. Aalto, S.; Brück, A.; Laine, M.; Någren, K.; Rinne, J.O. Frontal and Temporal Dopamine Release During Working Memory and Attention Tasks in Healthy Humans: A Positron Emission Tomography Study Using the High-Affinity Dopamine D2 Receptor Ligand [11C]FLB 457. *J. Neurosci.* **2005**, *25*, 2471–2477. [CrossRef] [PubMed]
327. Talvik, M.; Nordström, A.L.; Okubo, Y.; Olsson, H.; Borg, J.; Halldin, C.; Farde, L. Dopamine D2 Receptor Binding in Drug-Naïve Patients With Schizophrenia Examined With Raclopride-C11 and Positron Emission Tomography. *Psychiatry Res.* **2006**, *148*, 165–173. [CrossRef] [PubMed]
328. Kimura, Y.; Ito, H.; Shiraishi, T.; Fujiwara, H.; Kodaka, F.; Takano, H.; Shimada, H.; Kanno, I.; Suhara, T. Biodistribution and Radiation Dosimetry in Humans of [11C]FLB 457, a Positron Emission Tomography Ligand for the Extrastriatal Dopamine D2 Receptor. *Nucl. Med. Biol.* **2014**, *41*, 102–105. [CrossRef] [PubMed]
329. Farde, L.; Ehrin, E.; Eriksson, L.; Greitz, T.; Hall, H.; Hedström, C.G.; Litton, J.E.; Sedvall, G. Substituted Benzamides As Ligands for Visualization of Dopamine Receptor Binding in the Human Brain by Positron Emission Tomography. *Proc. Natl. Acad. Sci. USA* **1985**, *82*, 3863–3867. [CrossRef] [PubMed]
330. Kessler, R.M.; Votaw, J.R.; de Paulis, T.; Bingham, D.R.; Ansari, M.S.; Mason, N.S.; Holburn, G.; Schmidt, D.E.; Votaw, D.B.; Manning, R.G. Evaluation of 5-[18F]Fluoropropylpeptide As a Potential PET Radioligand for Imaging Dopamine D2 Receptors. *Synapse* **1993**, *15*, 169–176. [CrossRef]
331. Welch, M.J.; Katzenellenbogen, J.A.; Mathias, C.J.; Brodack, J.W.; Carlson, K.E.; Chi, D.Y.; Dence, C.S.; Kilbourn, M.R.; Perlmutter, J.S.; Raichle, M.E. N-(3-[18F]Fluoropropyl) Spiperone: The Preferred 18F Labeled Spiperone Analog for Positron Emission Tomographic Studies of the Dopamine Receptor. *Int. J. Rad. Appl. Instrum. B* **1988**, *15*, 83–97. [CrossRef]
332. Hashizume, K.; Tamakawa, H.; Hashimoto, N.; Miyake, Y. Single-Step Synthesis of [18F]Haloperidol From the Chloro-Precursor and Its Applications in PET Imaging of a Cat's Brain. *Appl. Radiat. Isot.* **1997**, *48*, 1179–1185. [CrossRef]
333. Zanzonico, P.B.; Bigler, R.E.; Schmall, B. Neuroleptic Binding Sites: Specific Labeling in Mice With [18F]Haloperidol, a Potential Tracer for Positron Emission Tomography. *J. Nucl. Med.* **1983**, *24*, 408–416.
334. Yousef, K.A.; Fowler, J.S.; Volkow, N.D.; Dewey, S.L.; Shea, C.; Schlyer, D.J.; Gatley, S.J.; Logan, J.; Wolf, A.P. [18F]Haloperidol Binding in Baboon Brain in Vivo. *Nucl. Med. Biol.* **1996**, *23*, 47–52. [CrossRef]
335. Mach, R.H.; Luedtke, R.R.; Unsworth, C.D.; Boundy, V.A.; Nowak, P.A.; Scripko, J.G.; Elder, S.T.; Jackson, J.R.; Hoffman, P.L.; Evora, P.H. 18F-Labeled Benzamides for Studying the Dopamine D2 Receptor With Positron Emission Tomography. *J. Med. Chem.* **1993**, *36*, 3707–3720. [CrossRef]
336. Mach, R.H.; Ehrenkaufer, R.L.; Greenberg, J.H.; Shao, L.; Morton, T.E.; Evora, P.H.; Nowak, P.A.; Luedtke, R.R.; Cohen, D.; Reivich, M. PET Imaging Studies of Dopamine D2 Receptors: Comparison of [18F]N-Methylspiperone and the Benzamide Analogue [18F]MABN and [18F]MBP in Baboon Brain. *Synapse* **1995**, *19*, 177–187. [CrossRef] [PubMed]
337. Seneca, N.; Zoghbi, S.S.; Skinbjerg, M.; Liow, J.S.; Hong, J.; Sibley, D.R.; Pike, V.W.; Halldin, C.; Innis, R.B. Occupancy of Dopamine D2/3 Receptors in Rat Brain by Endogenous Dopamine Measured With the Agonist Positron Emission Tomography Radioligand [11C]MNPA. *Synapse* **2008**, *62*, 756–763. [CrossRef]
338. Finnema, S.J.; Seneca, N.; Farde, L.; Shchukin, E.; Sóvágó, J.; Gulyás, B.; Wikström, H.V.; Innis, R.B.; Neumeyer, J.L.; Halldin, C.; et al. A Preliminary PET Evaluation of the New Dopamine D2 Receptor Agonist [11C]MNPA in Cynomolgus Monkey. *Nucl. Med. Biol.* **2005**, *32*, 353–360. [CrossRef]
339. Seneca, N.; Skinbjerg, M.; Zoghbi, S.S.; Liow, J.S.; Gladding, R.L.; Hong, J.; Kannan, P.; Tuan, E.; Sibley, D.R.; Halldin, C.; et al. Kinetic Brain Analysis and Whole-Body Imaging in Monkey of [11C]MNPA: A Dopamine Agonist Radioligand. *Synapse* **2008**, *62*, 700–709. [CrossRef] [PubMed]
340. Kodaka, F.; Ito, H.; Kimura, Y.; Fujie, S.; Takano, H.; Fujiwara, H.; Sasaki, T.; Nakayama, K.; Halldin, C.; Farde, L.; et al. Test-Retest Reproducibility of Dopamine D2/3 Receptor Binding in Human Brain Measured by PET With [11C]MNPA and [11C]Raclopride. *Eur. J. Nucl. Med. Mol. Imaging* **2013**, *40*, 574–579. [CrossRef] [PubMed]
341. Ishiwata, K.; Hayakawa, N.; Ogi, N.; Oda, K.; Toyama, H.; Endo, K.; Tanaka, A.; Senda, M. Comparison of Three PET Dopamine D2-Like Receptor Ligands, [11C]Raclopride, [11C]Nemonapride and [11C]N-Methylspiperone, in Rats. *Ann. Nucl. Med.* **1999**, *13*, 161–167. [CrossRef] [PubMed]
342. Ishiwata, K.; Senda, M. In Vivo Binding of [11C]Nemonapride to Sigma Receptors in the Cortex and Cerebellum. *Nucl. Med. Biol.* **1999**, *26*, 627–631. [CrossRef]
343. Ishiwata, K.; Ogi, N.; Hayakawa, N.; Umegaki, H.; Nagaoka, T.; Oda, K.; Toyama, H.; Endo, K.; Tanaka, A.; Senda, M. Positron Emission Tomography and Ex Vivo and in Vitro Autoradiography Studies on Dopamine D2-Like Receptor Degeneration in the Quinolinic Acid-Lesioned Rat Striatum: Comparison of [11C]Raclopride, [11C]Nemonapride and [11C]N-Methylspiperone. *Nucl. Med. Biol.* **2002**, *29*, 307–316. [CrossRef]

344. Burns, H.D.; Dannals, R.F.; Langström, B.; Ravert, H.T.; Zemyan, S.E.; Duelfer, T.; Wong, D.F.; Frost, J.J.; Kuhar, M.J.; Wagner, H.N. (3-N-[11C]Methyl)Spiperone, a Ligand Binding to Dopamine Receptors: Radiochemical Synthesis and Biodistribution Studies in Mice. *J. Nucl. Med.* **1984**, *25*, 1222–1227.
345. Yanai, K.; Ido, T.; Ishiwata, K.; Hatazawa, J.; Watanuki, S.; Takahashi, T.; Ujiie, A.; Ito, M.; Matsuzawa, T. Characteristics of Specific in Vivo Labeling of Neuroleptic Binding Sites With 3-N-[11C]Methylspiperone. *Eur. J. Nucl. Med.* **1986**, *11*, 438–443. [CrossRef]
346. Wagner, H.N., Jr.; Burns, H.D.; Dannals, R.F.; Wong, D.F.; Långström, B.; Duelfer, T.; Frost, J.J.; Ravert, H.T.; Links, J.M.; Rosenbloom, S.B. Assessment of Dopamine Receptor Densities in the Human Brain With Carbon-11-Labeled N-Methylspiperone. *Ann. Neurol.* **1984**, *15*, S79–S84. [CrossRef] [PubMed]
347. Wong, D.F.; Wagner, H.N., Jr.; Dannals, R.F.; Links, J.M.; Frost, J.J.; Ravert, H.T.; Wilson, A.A.; Rosenbaum, A.E.; Gjedde, A.; Douglass, K.H. Effects of Age on Dopamine and Serotonin Receptors Measured by Positron Tomography in the Living Human Brain. *Science* **1984**, *226*, 1393–1396. [CrossRef] [PubMed]
348. Andersson, U.; Eckernås, S.A.; Hartvig, P.; Ulin, J.; Långström, B.; Häggström, J.E. Striatal Binding of 11C-NMSP Studied With Positron Emission Tomography in Patients With Persistent Tardive Dyskinesia: No Evidence for Altered Dopamine D2 Receptor Binding. *J. Neural Transm. Gen. Sect.* **1990**, *79*, 215–226. [CrossRef]
349. Nordström, A.L.; Farde, L.; Eriksson, L.; Halldin, C. No Elevated D2 Dopamine Receptors in Neuroleptic-Naïve Schizophrenic Patients Revealed by Positron Emission Tomography and [11C]N-Methylspiperone. *Psychiatry Res.* **1995**, *61*, 67–83. [CrossRef]
350. Arnett, C.D.; Fowler, J.S.; Wolf, A.P.; Shiue, C.Y.; McPherson, D.W. [18F]-N-Methylspiroperidol: The Radioligand of Choice for PETT Studies of the Dopamine Receptor in Human Brain. *Life Sci.* **1985**, *36*, 1359–1366. [CrossRef]
351. Arnett, C.D.; Wolf, A.P.; Shiue, C.Y.; Fowler, J.S.; MacGregor, R.R.; Christman, D.R.; Smith, M.R. Improved Delineation of Human Dopamine Receptors Using [18F]-N-Methylspiroperidol and PET. *J. Nucl. Med.* **1986**, *27*, 1878–1882. [PubMed]
352. Hwang, D.R.; Kegeles, L.S.; Laruelle, M. (-)-N-[(11)C]Propyl-Norapomorphine: A Positron-Labeled Dopamine Agonist for PET Imaging of D(2) Receptors. *Nucl. Med. Biol.* **2000**, *27*, 533–539. [CrossRef]
353. Cumming, P.; Gillings, N.M.; Jensen, S.B.; Bjarkam, C.; Gjedde, A. Kinetics of the Uptake and Distribution of the Dopamine D(2,3) Agonist (R)-N-[1-(11)C]n-Propyl-norapomorphine in Brain of Healthy and MPTP-Treated Göttingen Miniature Pigs. *Nucl. Med. Biol.* **2003**, *30*, 547–553. [CrossRef]
354. Narendran, R.; Hwang, D.R.; Slifstein, M.; Talbot, P.S.; Erritzoe, D.; Huang, Y.; Cooper, T.B.; Martinez, D.; Kegeles, L.S.; bi-Dargham, A.; et al. In Vivo Vulnerability to Competition by Endogenous Dopamine: Comparison of the D2 Receptor Agonist Radiotracer (-)-N-[11C]Propyl-Norapomorphine ([11C]NPA) With the D2 Receptor Antagonist Radiotracer [11C]-Raclopride. *Synapse* **2004**, *52*, 188–208. [CrossRef] [PubMed]
355. Narendran, R.; Hwang, D.R.; Slifstein, M.; Hwang, Y.; Huang, Y.; Ekelund, J.; Guillen, O.; Scher, E.; Martinez, D.; Laruelle, M. Measurement of the Proportion of D2 Receptors Configured in State of High Affinity for Agonists in Vivo: A Positron Emission Tomography Study Using [11C]N-Propyl-Norapomorphine and [11C]Raclopride in Baboons. *J. Pharmacol. Exp. Ther.* **2005**, *315*, 80–90. [CrossRef] [PubMed]
356. Narendran, R.; Frankle, W.G.; Mason, N.S.; Laymon, C.M.; Lopresti, B.J.; Price, J.C.; Kendro, S.; Vora, S.; Litschge, M.; Mountz, J.M.; et al. Positron Emission Tomography Imaging of D(2/3) Agonist Binding in Healthy Human Subjects With the Radiotracer [(11)C]-N-Propyl-Norapomorphine: Preliminary Evaluation and Reproducibility Studies. *Synapse* **2009**, *63*, 574–584. [CrossRef] [PubMed]
357. Narendran, R.; Mason, N.S.; Laymon, C.M.; Lopresti, B.J.; Velasquez, N.D.; May, M.A.; Kendro, S.; Martinez, D.; Mathis, C.A.; Frankle, W.G. A Comparative Evaluation of the Dopamine D(2/3) Agonist Radiotracer [11C](-)-N-Propyl-Norapomorphine and Antagonist [11C]Raclopride to Measure Amphetamine-Induced Dopamine Release in the Human Striatum. *J. Pharmacol. Exp. Ther.* **2010**, *333*, 533–539. [CrossRef] [PubMed]
358. Narendran, R.; Martinez, D.; Mason, N.S.; Lopresti, B.J.; Himes, M.L.; Chen, C.M.; May, M.A.; Price, J.C.; Mathis, C.A.; Frankle, W.G. Imaging of Dopamine D2/3 Agonist Binding in Cocaine Dependence: A [11C]NPA Positron Emission Tomography Study. *Synapse* **2011**, *65*, 1344–1349. [CrossRef] [PubMed]
359. Mukherjee, J.; Narayanan, T.K.; Christian, B.T.; Shi, B.; Yang, Z.Y. Binding Characteristics of High-Affinity Dopamine D2/D3 Receptor Agonists, 11C-PPHT and 11C-ZYY-339 in Rodents and Imaging in Non-Human Primates by PET. *Synapse* **2004**, *54*, 83–91. [CrossRef]
360. Shi, B.; Narayanan, T.K.; Yang, Z.Y.; Christian, B.T.; Mukherjee, J. Radiosynthesis and in Vitro Evaluation of 2-(N-Alkyl-N-1'-11C-Propyl)Amino-5-Hydroxytetralin Analogs As High Affinity Agonists for Dopamine D-2 Receptors. *Nucl. Med. Biol.* **1999**, *26*, 725–735. [CrossRef]
361. Hume, S.P.; Myers, R.; Bloomfield, P.M.; Opacka-Juffry, J.; Cremer, J.E.; Ahier, R.G.; Luthra, S.K.; Brooks, D.J.; Lammertsma, A.A. Quantitation of Carbon-11-Labeled Raclopride in Rat Striatum Using Positron Emission Tomography. *Synapse* **1992**, *12*, 47–54. [CrossRef] [PubMed]
362. Hume, S.P.; Opacka-Juffry, J.; Myers, R.; Ahier, R.G.; Ashworth, S.; Brooks, D.J.; Lammertsma, A.A. Effect of L-Dopa and 6-Hydroxydopamine Lesioning on [11C]Raclopride Binding in Rat Striatum, Quantified Using PET. *Synapse* **1995**, *21*, 45–53. [CrossRef] [PubMed]
363. Ehrin, E.; Farde, L.; de Paulis, T.; Eriksson, L.; Greitz, T.; Johnström, P.; Litton, J.E.; Nilsson, J.L.; Sedvall, G.; Stone-Elander, S. Preparation of 11C-Labelled Raclopride, a New Potent Dopamine Receptor Antagonist: Preliminary PET Studies of Cerebral Dopamine Receptors in the Monkey. *Int. J. Appl. Radiat. Isot.* **1985**, *36*, 269–273. [CrossRef]

364. Hartvig, P.; Eckernås, S.A.; Ekblom, B.; Lindström, L.; Lundqvist, H.; Axelsson, S.; Fasth, K.J.; Gullberg, P.; Långström, B. Receptor Binding and Selectivity of Three <sup>11</sup>C-Labelled Dopamine Receptor Antagonists in the Brain of Rhesus Monkeys Studied With Positron Emission Tomography. *Acta Neurol. Scand.* **1988**, *77*, 314–321. [CrossRef] [PubMed]
365. Dewey, S.L.; Smith, G.S.; Logan, J.; Brodie, J.D.; Fowler, J.S.; Wolf, A.P. Striatal Binding of the PET Ligand <sup>11</sup>C-Raclopride Is Altered by Drugs That Modify Synaptic Dopamine Levels. *Synapse* **1993**, *13*, 350–356. [CrossRef] [PubMed]
366. Endres, C.J.; Kolachana, B.S.; Saunders, R.C.; Su, T.; Weinberger, D.; Breier, A.; Eckelman, W.C.; Carson, R.E. Kinetic Modeling of <sup>11</sup>C-Raclopride: Combined PET-Microdialysis Studies. *J. Cereb. Blood Flow Metab.* **1997**, *17*, 932–942. [CrossRef] [PubMed]
367. Tsukada, H.; Nishiyama, S.; Kakiuchi, T.; Ohba, H.; Sato, K.; Harada, N. Is Synaptic Dopamine Concentration the Exclusive Factor Which Alters the in Vivo Binding of <sup>11</sup>C-Raclopride? PET Studies Combined With Microdialysis in Conscious Monkeys. *Brain Res.* **1999**, *841*, 160–169. [CrossRef]
368. Farde, L.; Hall, H.; Ehrin, E.; Sedvall, G. Quantitative Analysis of D2 Dopamine Receptor Binding in the Living Human Brain by PET. *Science* **1986**, *231*, 258–261. [CrossRef]
369. Farde, L.; Pauli, S.; Hall, H.; Eriksson, L.; Halldin, C.; Höglberg, T.; Nilsson, L.; Sjögren, I.; Stone-Elander, S. Stereoselective Binding of <sup>11</sup>C-Raclopride in Living Human Brain—a Search for Extrastriatal Central D2-Dopamine Receptors by PET. *Psychopharmacology* **1988**, *94*, 471–478. [CrossRef]
370. Farde, L.; Wiesel, F.A.; Stone-Elander, S.; Halldin, C.; Nordström, A.L.; Hall, H.; Sedvall, G. D2 Dopamine Receptors in Neuroleptic-Naïve Schizophrenic Patients. A Positron Emission Tomography Study With <sup>11</sup>C-Raclopride. *Arch. Gen. Psychiatry* **1990**, *47*, 213–219. [CrossRef]
371. Laihinen, A.; Rinne, J.O.; Någren, K.; Bergman, J.; Haaparanta, M.; Solin, O.; Ruotsalainen, U.; Rinne, U.K. Positron Emission Tomography of Brain Dopamine D-2 Receptors With <sup>11</sup>C-Raclopride in Early Parkinson's Disease. *Acta Radiol. Suppl.* **1991**, *376*, 151.
372. Brooks, D.J.; Ibanez, V.; Sawle, G.V.; Playford, E.D.; Quinn, N.; Mathias, C.J.; Lees, A.J.; Marsden, C.D.; Bannister, R.; Frackowiak, R.S. Striatal D2 Receptor Status in Patients With Parkinson's Disease, Striatonigral Degeneration, and Progressive Supranuclear Palsy, Measured With <sup>11</sup>C-Raclopride and Positron Emission Tomography. *Ann. Neurol.* **1992**, *31*, 184–192. [CrossRef] [PubMed]
373. Antonini, A.; Leenders, K.L. Dopamine D2 Receptors in Normal Human Brain: Effect of Age Measured by Positron Emission Tomography (PET) and <sup>11</sup>C-Raclopride. *Ann. N. Y. Acad. Sci.* **1993**, *695*, 81–85. [CrossRef] [PubMed]
374. Antonini, A.; Leenders, K.L.; Reist, H.; Thomann, R.; Beer, H.F.; Locher, J. Effect of Age on D2 Dopamine Receptors in Normal Human Brain Measured by Positron Emission Tomography and <sup>11</sup>C-Raclopride. *Arch. Neurol.* **1993**, *50*, 474–480. [CrossRef]
375. Volkow, N.D.; Wang, G.J.; Fowler, J.S.; Logan, J.; Schlyer, D.; Hitzemann, R.; Lieberman, J.; Angrist, B.; Pappas, N.; MacGregor, R. Imaging Endogenous Dopamine Competition With <sup>11</sup>C-Raclopride in the Human Brain. *Synapse* **1994**, *16*, 255–262. [CrossRef] [PubMed]
376. Rinne, J.O.; Laihinen, A.; Ruottinen, H.; Ruotsalainen, U.; Någren, K.; Lehikoinen, P.; Oikonen, V.; Rinne, U.K. Increased Density of Dopamine D2 Receptors in the Putamen, but Not in the Caudate Nucleus in Early Parkinson's Disease: A PET Study With <sup>11</sup>C-Raclopride. *J. Neurol. Sci.* **1995**, *132*, 156–161. [CrossRef]
377. Volkow, N.D.; Wang, G.J.; Fowler, J.S.; Logan, J.; Gatley, S.J.; MacGregor, R.R.; Schlyer, D.J.; Hitzemann, R.; Wolf, A.P. Measuring Age-Related Changes in Dopamine D2 Receptors With <sup>11</sup>C-Raclopride and <sup>18</sup>F-N-Methylspiroperidol. *Psychiatry Res.* **1996**, *67*, 11–16. [CrossRef]
378. Antonini, A.; Schwarz, J.; Oertel, W.H.; Pogarell, O.; Leenders, K.L. Long-Term Changes of Striatal Dopamine D2 Receptors in Patients With Parkinson's Disease: A Study With Positron Emission Tomography and <sup>11</sup>C-Raclopride. *Mov. Disord.* **1997**, *12*, 33–38. [CrossRef] [PubMed]
379. Hietala, J.; Någren, K.; Lehikoinen, P.; Ruotsalainen, U.; Syvälahti, E. Measurement of Striatal D2 Dopamine Receptor Density and Affinity With <sup>11</sup>C-Raclopride in Vivo: A Test-Retest Analysis. *J. Cereb. Blood Flow Metab.* **1999**, *19*, 210–217. [CrossRef]
380. Xu, J.; Vangveravong, S.; Li, S.; Fan, J.; Jones, L.A.; Cui, J.; Wang, R.; Tu, Z.; Chu, W.; Perlmutter, J.S.; et al. Positron Emission Tomography Imaging of Dopamine D2 Receptors Using a Highly Selective Radiolabeled D2 Receptor Partial Agonist. *Neuroimage* **2013**, *71*, 168–174. [CrossRef]
381. Volkow, N.D.; Wang, G.J.; Logan, J.; Alexoff, D.; Fowler, J.S.; Thanos, P.K.; Wong, C.; Casado, V.; Ferre, S.; Tomasi, D. Caffeine Increases Striatal Dopamine D2/D3 Receptor Availability in the Human Brain. *Transl. Psychiatry* **2015**, *5*, e549. [CrossRef] [PubMed]
382. Kaasinen, V.; Aalto, S.; Någren, K.; Rinne, J.O. Dopaminergic Effects of Caffeine in the Human Striatum and Thalamus. *Neuroreport* **2004**, *15*, 281–285. [CrossRef]
383. Prasad, K.; de Vries, E.F.J.; Sijbesma, J.W.A.; Attia, K.; Kwizera, C.; García Varéla, L.; Vazquez-Matias, D.A.; Moraga Amaro, R.; Vállez-García, D.; Dierckx, R.A.J.O.; et al. Impact of A2AR Agonist and Antagonist on Binding of the Dopamine D2 Receptor Ligand <sup>11</sup>C-Raclopride in the Rodent Striatum. *Eur. J. Nucl. Med. Mol. Imaging* **2020**, *47*, S114–S115.
384. Golembiowska, K.; Dziubina, A. Striatal Adenosine A(2A) Receptor Blockade Increases Extracellular Dopamine Release Following L-DOPA Administration in Intact and Dopamine-Denervated Rats. *Neuropharmacology* **2004**, *47*, 414–426. [CrossRef] [PubMed]
385. Petzer, J.P.; Steyn, S.; Castagnoli, K.P.; Chen, J.F.; Schwarzschild, M.A.; Van der Schyf, C.J.; Castagnoli, N. Inhibition of Monoamine Oxidase B by Selective Adenosine A2A Receptor Antagonists. *Bioorg. Med. Chem.* **2003**, *11*, 1299–1310. [CrossRef]
386. Morelli, M.; Di, P.T.; Wardas, J.; Calon, F.; Xiao, D.; Schwarzschild, M.A. Role of Adenosine A2A Receptors in Parkinsonian Motor Impairment and L-DOPA-Induced Motor Complications. *Prog. Neurobiol.* **2007**, *83*, 293–309. [CrossRef] [PubMed]

387. Casadó, V.; Cortés, A.; Mallol, J.; Pérez-Capote, K.; Ferré, S.; Lluis, C.; Franco, R.; Canela, E.I. GPCR Homomers and Heteromers: A Better Choice As Targets for Drug Development Than GPCR Monomers? *Pharmacol. Ther.* **2009**, *124*, 248–257. [CrossRef] [PubMed]
388. Shao, Y.M.; Ma, X.; Paira, P.; Tan, A.; Herr, D.R.; Lim, K.L.; Ng, C.H.; Venkatesan, G.; Klotz, K.N.; Federico, S.; et al. Discovery of Indolylpiperazinylpyrimidines With Dual-Target Profiles at Adenosine A2A and Dopamine D2 Receptors for Parkinson’s Disease Treatment. *PLoS ONE* **2018**, *13*, e0188212. [CrossRef]
389. Daniels, D.J.; Lenard, N.R.; Etienne, C.L.; Law, P.Y.; Roerig, S.C.; Portoghesi, P.S. Opioid-Induced Tolerance and Dependence in Mice Is Modulated by the Distance Between Pharmacophores in a Bivalent Ligand Series. *Proc. Natl. Acad. Sci. USA* **2005**, *102*, 19208–19213. [CrossRef]
390. Akgün, E.; Javed, M.I.; Lunzer, M.M.; Smeester, B.A.; Beitz, A.J.; Portoghesi, P.S. Ligands That Interact With Putative MOR-MGluR5 Heteromer in Mice With Inflammatory Pain Produce Potent Antinociception. *Proc. Natl. Acad. Sci. USA* **2013**, *110*, 11595–11599. [CrossRef]
391. Soriano, A.; Ventura, R.; Molero, A.; Hoen, R.; Casadó, V.; Cortés, A.; Fanelli, F.; Albericio, F.; Lluís, C.; Franco, R.; et al. Adenosine A2A Receptor-Antagonist / Dopamine D2 Receptor-Agonist Bivalent Ligands as Pharmacological Tools to Detect A2A-D2 Receptor Heteromers. *J. Med. Chem.* **2009**, *52*, 5590–5602. [CrossRef]
392. Shonberg, J.; Scammells, P.J.; Capuano, B. Design Strategies for Bivalent Ligands Targeting GPCRs. *ChemMedChem* **2011**, *6*, 963–974. [CrossRef] [PubMed]
393. Clark, D.E. Computational Prediction of Blood-Brain Barrier Permeation. *Annu. Rep. Med. Chem.* **2005**, *40*, 403–415.
394. Dalpiaz, A.; Cacciari, B.; Vicentini, C.B.; Bortolotti, F.; Spalluto, G.; Federico, S.; Pavan, B.; Vincenzi, F.; Borea, P.A.; Varani, K. A Novel Conjugated Agent between Dopamine and an A2A Adenosine Receptor Antagonist as a Potential Anti-Parkinson Multitarget Approach. *Mol. Pharm.* **2012**, *9*, 591–604. [CrossRef] [PubMed]
395. Jörg, M.; May, L.T.; Mak, F.S.; Lee, K.C.; Miller, N.D.; Scammells, P.J.; Capuano, B. Synthesis and Pharmacological Evaluation of Dual Acting Ligands Targeting the Adenosine A2A and Dopamine D2 Receptors for the Potential Treatment of Parkinson’s Disease. *J. Med. Chem.* **2015**, *58*, 718–738. [CrossRef]
396. Agnati, L.F.; Ferré, S.; Genedani, S.; Leo, G.; Guidolin, D.; Filaferro, M.; Carriba, P.; Casadó, V.; Lluís, C.; Franco, R.; et al. Allosteric Modulation of Dopamine D2 Receptors by Homocysteine. *J. Proteome Res.* **2006**, *5*, 3077–3083. [CrossRef] [PubMed]
397. Cervetto, C.; Venturini, A.; Guidolin, D.; Maura, G.; Passalacqua, M.; Tacchetti, C.; Cortelli, P.; Genedani, S.; Candiani, S.; Ramoino, P.; et al. Homocysteine and A2A-D2 Receptor-Receptor Interaction at Striatal Astrocyte Processes. *J. Mol. Neurosci.* **2018**, *65*, 456–466. [CrossRef] [PubMed]
398. Schwarzschild, M.A.; Xu, K.; Oztas, E.; Petzer, J.P.; Castagnoli, K.; Castagnoli, N., Jr.; Chen, J.F. Neuroprotection by Caffeine and More Specific A2A Receptor Antagonists in Animal Models of Parkinson’s Disease. *Neurology* **2003**, *61*, S55–S61. [CrossRef] [PubMed]
399. Yu, L.; Shen, H.Y.; Coelho, J.E.; Araújo, I.M.; Huang, Q.Y.; Day, Y.J.; Rebola, N.; Canas, P.M.; Rapp, E.K.; Ferrara, J.; et al. Adenosine A2A Receptor Antagonists Exert Motor and Neuroprotective Effects by Distinct Cellular Mechanisms. *Ann. Neurol.* **2008**, *63*, 338–346. [CrossRef]
400. Cerri, S.; Levandis, G.; Ambrosi, G.; Montepeloso, E.; Antoninetti, G.F.; Franco, R.; Lanciego, J.L.; Baqi, Y.; Müller, C.E.; Pinna, A.; et al. Neuroprotective Potential of Adenosine A2A and Cannabinoid CB1 Receptor Antagonists in an Animal Model of Parkinson Disease. *J. Neuropathol. Exp. Neurol.* **2014**, *73*, 414–424. [CrossRef] [PubMed]
401. Fathalla, A.M.; Soliman, A.M.; Ali, M.H.; Moustafa, A.A. Adenosine A2A Receptor Blockade Prevents Rotenone-Induced Motor Impairment in a Rat Model of Parkinsonism. *Front. Behav. Neurosci.* **2016**, *10*, 35. [CrossRef] [PubMed]
402. Xu, K.; Di Luca, D.G.; Orrú, M.; Xu, Y.; Chen, J.F.; Schwarzschild, M.A. Neuroprotection by Caffeine in the MPTP Model of Parkinson’s Disease and Its Dependence on Adenosine A2A Receptors. *Neuroscience* **2016**, *322*, 129–137. [CrossRef] [PubMed]
403. Fathalla, A.M.; Soliman, A.M.; Moustafa, A.A. Selective A(2A) Receptors Blockade Reduces Degeneration of Substantia Nigra Dopamine Neurons in a Rotenone-Induced Rat Model of Parkinson’s Disease: A Histological Study. *Neurosci. Lett.* **2017**, *643*, 89–96. [CrossRef] [PubMed]