

Insights into the Structure and Protein Composition of *Moorella thermoacetica* Spores Formed at Different Temperatures

Tiffany Malleck ^{1,2}, Fatima Fekraoui ², Isabelle Bornard ³, Céline Henry ⁴, Eloi Haudebourg ⁴, Stella Planchon ^{2,*} and Véronique Broussolle ^{1,*}

¹ INRAE, Avignon Université, UMR SQPOV, F-84000 Avignon, France

² Unité EMaiRIT'S, CTCPA, F-84911 Avignon, France

³ INRAE, Pathologie végétale, F-84143 Montfavet, France

⁴ PAPPSO, Micalis Institute, INRAE, AgroParisTech, Université Paris-Saclay, F-78352 Jouy-en-Josas, France

* Correspondence: veronique.broussolle@inrae.fr (V.B.); splanchon@ctcpa.org (S.P.); Tel.: +33-(0)432722518 (V.B.); +33-(0)490843246 (S.P.)





Figure S1. Spores of *Moorella thermoacetica* strain ATCC 39073, produced on agar plates at 45 °C and observed by scanning electron microscopy. The exosporium (Ex) is indicated by the white arrow.




Figure S2. Heterogeneity of spore diameters of *Moorella thermoacetica* at 45 °C (blue bars) and 55 °C (orange bars), measured on SEM images.




Figure S3. SDS-PAGE gel (short migration) of *Moorella thermoacetica* spore protein extracts (three biological replicates for 45 °C and for 55 °C) used for LC-MS/MS analysis. Mw, molecular weight ladder.