



Review

Role of Leptin and Adiponectin in Endometrial Cancer

Aneta Słabuszewska-Jóźwiak ^{1,*}, Aron Lukaszuk ^{2,3}, Marta Janicka-Kośnik ², Artur Wdowiak ⁴
and Grzegorz Jakiel ¹

¹ First Department of Obstetrics and Gynecology, Centre of Postgraduate Medical Education, Żelazna 90 Street, 01-004 Warsaw, Poland; grzegorz.jakiel1@o2.pl

² Saint Sophia Hospital, Żelazna 90 Street, 01-004 Warsaw, Poland; aron@gumed.edu.pl (A.L.); lek.martajanicka@gmail.com (M.J.-K.)

³ Invicta Research and Development Center, Polna 64 Street, 81-710 Sopot, Poland

⁴ Chair of Obstetrics and Gynecology, Faculty of Health Sciences, Medical University of Lublin, 4-6 Staszica St., 20-081 Lublin, Poland; wdowiakartur@gmail.com

* Correspondence: anetaslabuszewska@gmail.com

Abstract: Endometrial cancer is the most common malignancy of the female genital tract. Obesity is a strong risk factor for endometrial cancer. Adipose tissue is an active endocrine organ that synthesizes biologically active cytokine peptides, called adipokines. Adiponectin and leptin are the main cytokines of adipose tissue, which may influence the development of metabolic diseases and carcinogenesis. In this scenario, we describe the role of leptin and adiponectin in the development of endometrial cancer. A better understanding of the signalling pathway of these cytokines in endometrial cancerogenesis will provide an opportunity for effective target therapy and may be usable in fertility-sparing treatment. In the future, clinical trials focusing on adipokines, molecular biology, and genetics of the tumour will be needed.



Citation: Słabuszewska-Jóźwiak, A.; Lukaszuk, A.; Janicka-Kośnik, M.; Wdowiak, A.; Jakiel, G. Role of Leptin and Adiponectin in Endometrial Cancer. *Int. J. Mol. Sci.* **2022**, *23*, 5307. <https://doi.org/10.3390/ijms23105307>

Academic Editors: Antonio Simone Laganà, Gaspare Cucinella and Giuseppe Gullo

Received: 22 March 2022

Accepted: 7 May 2022

Published: 10 May 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Keywords: leptin; adiponectin; endometrial cancer; signalling pathway

1. Introduction

According to world statistics, endometrial cancer is the second most frequently detected malignant neoplasm of the female genital tract after cervical cancer [1]. It mainly occurs in women over the age of 50 [2], and is associated with obesity, diabetes, and metabolic syndrome. Based on clinical, endocrine, and histopathological features, two types of endometrial cancer are distinguished.

Type I occurs in 80% of cases, includes the histological type of endometrioid carcinoma, and is formed on endometrial hyperplasia, with the presence of atypia, as a result of estrogenic overstimulation. Type II is hormone-independent and includes serous, clear-cell and poorly differentiated carcinomas. It mainly occurs in women in the 7th and 8th decade of life, and is characterized by dynamic growth and poor prognosis [3].

Multiple genetic disorders were found in endometrial neoplasms. Mutations of PTEN, K-ras, β -catenin, and PIK3CA, as well as mutations in mismatch genes (MMR) and microsatellite instability (MSI), are common in endometrial cancer. Type II tumours are dominated by mutations within the p53 and p16 genes, and overexpression or amplification of the HER-2/neu gene. The Genome Atlas project identified four molecular subtypes of endometrial carcinomas: POLE ultramutated with a mutation in the POLE gene, MSI hypermutated with microsatellite instability MSI-H, subtype with a small number of copy alterations—SNCA without genetic markers, and subtype with a high number of copy alterations—SNCA with the TP53 mutation. These subtypes differ not only in genetic disorders, but also in their clinical course and prognosis. It should be noted that obesity is recognized as an independent risk factor for the development of endometrial cancer, regardless of the histopathological type [4].

Adipose tissue is an active endocrine organ that synthesizes biologically active cytokine peptides, called adipokines, that act within the adipose tissue (auto and paracrine), and on distant organs and tissues. Adiponectin and leptin are the main cytokines of adipose tissue, affecting insulin resistance, lipolysis [5], and inflammatory pathways, which may influence the development of metabolic diseases and carcinogenesis [6]. An increased leptin-to-adiponectin ratio is associated with an increased risk of cancer, including endometrial cancer [7–9].

2. Leptin and Adiponectin

2.1. Leptin

Leptin is a 16 kDa adipocytokine encoded by the LEP gene, mainly produced in white adipose tissue [8]. Similarly to many hormones, it is secreted in a free form, responsible for controlling the mass of adipose tissue in the body, and in a form associated with the protein being its soluble receptor. The complex form of the adipokine determines energy expenditure.

The structure of leptin is a quadruple helix, very similar to proteins from the cytokine family, such as IL-2, IL-6, IL-12, IL-15, IF, and G-CSF. By acting directly on the nervous system, leptin regulates food intake, and through its peripheral action, it inhibits insulin secretion from pancreatic β -cells stimulated by glucagon-like peptide-1 (GLP-1). In addition, it stimulates lipolysis, glucose utilization, and carbohydrate transport in adipocytes. It also takes part in thermogenesis, haematopoiesis, and osteogenesis, and has pro-angiogenic effects. It has been proven that leptin affects the functioning of the immune system by increasing the pro-inflammatory effects, including the activation of T lymphocytes, increasing the synthesis and release of IL-6 and TNF α , stimulating the response to Th1 lymphocytes, and increasing the activity of NK cells, macrophages, and neutrophils by intensifying their chemotaxis [10]. The action of leptin at the cellular level is related to the presence of membrane receptors (Ob-R), widespread in target organs (brain tissue, heart, lungs, kidneys, liver, pancreas, intestines, placenta, gonads, spleen, and thymus). The membrane receptor is a glycoprotein that is similar in structure to class I cytokine receptors. The gene encoding the leptin receptor is located on the 1p32 chromosome. It is responsible for the synthesis of at least five isoforms: long form—OB-Rb—showing the ability to transmit intracellular signals, three short forms—OB-Ra, OB-Rc, and OB-Rd, and one soluble receptor isoform—OB-Re. Within the OB-Rb receptor, the presence of an extracellular domain, with leptin binding sites, and a transcellular domain, with interaction sites for Janus kinase (JAK) and STAT (signal transducers and activators of transcription) proteins, has been demonstrated. The mechanism of action of leptin is, among others, based on the activation of STAT signalling proteins. Within the OB-Rb receptor, it homodimerizes and activates JAK protein kinases, which are responsible for the phosphorylation of STAT signalling proteins. After phosphorylation, these proteins move to the cell nucleus, where they bind to the appropriate gene, stimulating its transcription. Moreover, leptin stimulates cancer stem cells, which leads to cell proliferation and tumour growth [11–13]. There is a positive correlation between leptin levels and body mass index (BMI). Elevated levels of leptin and its receptors may potentially contribute to the progression of cancer [14]. The role of leptin and adiponectin in signalling pathways in carcinogenesis is presented in Figure 1.

2.2. Adiponectin

Adiponectin, apM1 (adipose most abundant gene transcript 1), is a peptide hormone that is mainly synthesized in white adipose tissue, which plays an important role in glycaemic and lipidemic homeostasis. Adiponectin is made of 244 amino acids and comes in two forms: homotrimers with a low molecular weight (LMW—low molecular weight), and in the form of high-molecular-weight oligomers (HMW—high molecular weight). This cytokine has an affinity for two receptors—AdipoR1, found mainly in skeletal muscle, and AdipoR2, found in liver tissue [15–17]. Both receptors are in the endometrium, in the placenta [18], and in neoplastic tissue, including endometrial cancer [19]. Adiponectin, in

contrast to leptin, has antiproliferative, anti-angiogenic and anti-atherosclerotic effects [20]. It has a selective influence on mitogenic growth factors, i.e., fibroblast growth factor, platelet-driven growth factor BB (PDGFBB), fibroblast growth factor (FGF), and heparin-binding epidermal growth factor-like growth factor (HB EGF), which causes a reduction in the bioavailability at the pre-receptor stage, thus weakening the DNA synthesis and cell proliferation induced by these factors [21]. Hence, the decreased levels of adiponectin observed in obese individuals may become a carcinogenic factor.




Figure 1. Schematic diagram shows the interaction of leptin and adiponectin and their downstream signalling pathways in endometrial cancer. The black arrows indicate stimulatory effects. The red blunt arrows indicate inhibitory effects. AMPK—AMP-activated protein kinase; ERK1/2—extracellular regulated kinase 1 or 2; JAK—Janus kinase; mTOR—mammalian target of rapamycin; PI3K/Akt—phosphatidylinositol 3-kinase/protein kinase B; STAT3—signal transducer and activator of transcription; Lep-R—leptin receptor; AdipoR1/AdipoR2—adipokine receptors.

The major signalling pathway for adiponectin involves AMP-activated protein kinase (AMPK) [22]. AMPK inhibits key kinase signalling pathways, such as extracellular 1/2-regulated kinase (ERK1/2), phosphatidylinositol-3-kinase (PI3K), protein kinase B (Akt), c-Jun N-terminal kinase (JNK), terminal kinase (cJNK), and STAT3, which are involved in the cell cycle that guarantees proper cell growth and survival [23–26]. The AMPK pathway is also crucial for cell growth by regulating Akt/mTOR/S6K transmission. It is presented in Figure 1. Thus, decreased AMPK phosphorylation leads to the stimulation of cell proliferation [27], and its activation can inhibit cell growth and initiate apoptosis [28,29]. AMPK can be phosphorylated and activated by LKB1 and by calmodulin-dependent protein kinase kinase- β (CaMKK- β) [30–32]. It should be emphasized that LKB1 is a tumour suppressor protein, and is an upstream kinase of tumour energy [33].

Moreover, adiponectin can directly regulate the expression of proteins involved in the cell cycle and apoptosis by affecting p53 and Bax (up-regulating), as well as c-myc, cyclin D1, and Bcl-2 (down-regulating) [34]. This cytokine also has the ability to inhibit the phosphorylation of the nuclear transcription factor κ B (NF- κ B), thus influencing the activity of various pro-inflammatory mediators [35].

Adiponectin is also involved in the energy metabolism of the organism, intensifying the process of lipid oxidation in peripheral tissues (white adipose tissue and muscle tissue) by stimulating the synthesis and activity of enzymes involved in triglyceride metabolism (Acyl-CoA oxidase, 5-protein kinase, and triglyceride lipase). In addition, it inhibits triglyceride synthesis and gluconeogenesis in the liver, and plays a significant hypoglycaemic and lipid-lowering role [16,36]. By activating the nuclear receptor for PPAR γ , which is associated with insulin resistance, adiponectin has become a marker of both insulin sensitivity and insulin resistance.

3. Leptin Effect on the Development of Endometrial Cancer

Neoplastic tumour progression is based on the unlimited ability of its cells to proliferate and their potential immortality. The continuous growth and development of the tumour depends on providing it with increased amounts of oxygen and nutrients, and those can only be sustained by neoangiogenesis. At a later stage, epithelial–mesenchymal transformations are initiated, aimed at cancer cells acquiring the ability to move to blood and lymphatic vessels, and then to spread all over the body. The emerging hypoxia in neoplastic tissue regulates the expression of various growth factors and their receptors.

HIF-1 is a marker of oxygen deficiency in any tissue, including neoplastic; hence, the overexpression of this factor in endometrial cancer is associated with stimulated angiogenesis and an unfavourable prognosis [37]. HIF-1 overexpression is associated with increased transcription of the leptin gene and its receptor (ObR) [38–40]. Both leptin and HIF-1 activate STAT (signal transducer and activator of transcription) proteins, which are related to the invasiveness of the endometrial cancer cell line [41]. STAT proteins play a role in the transduction of extracellular signals, and belong to the direct transcription factors regulating gene expression [42]. The STAT protein family consists of seven members (STAT1, STAT2, STAT3, STAT4, STAT5a, STAT5b, and STAT6). These proteins are associated with the inflammation, survival, proliferation, metastasis, angiogenesis, and chemoresistance of neoplastic cells [43].

Initially, they are present in the cytoplasm in an inactive form and become activated after binding to signalling peptides, which include cytokines, growth factors, and hormones. When activated by cytokines, the cytokine receptor dimerizes and then binds to the cell membrane, leading to the activation of JAK (Janus-activated kinase) by rapid phosphorylation of cytoplasmic domains. Phosphorylation generates a recognition site for STAT proteins with a homologous Src-binding domain (SH2—Src homology2), allowing them to form homo- or heterodimers and rapidly translocate to the cell nucleus. In the nucleus, dimers bind to specific sequences (often referred to as activated sequences) in gene promoter regions to stimulate their transcription [44]. In endometrial cancer, leptin activates STAT3 proteins, which increase their activity in the process of oncogenesis by stimulating proliferation [45], promoting angiogenesis, and avoiding the control of the immune system. A study on an endometrial cancer cell line showed that activated STAT3 proteins overexpress anti-apoptotic genes, such as Bcl-xL and Mcl-1, rendering these cells immortal and chemo-resistant [46]. In a study on SPEC-2 cell lines, leptin, as opposed to adiponectin, led to an increase in the activity of metalloproteinases MMP-2 and MMP-9, which are associated with the aggressiveness of the neoplastic process. Moreover, a higher tissue concentration of leptin positively correlates with the degree of myometrial invasion and lymph node metastases, eventually leading to a worse prognosis [47]. Similar conclusions were provided by the study by Cymbaluk-Płoska et al., which showed that higher levels of serum leptin correlate with the involvement of lymphatic vessels and with

a low degree of neoplastic differentiation [48]. Hence, inhibition of the leptin-promoted STAT3/JAK signalling pathway may limit tumour metastasis.

Notch is an evolutionarily conserved cellular information pathway that influences the differentiation, proliferation, and apoptosis of cells at various stages of their development [49]. Changes in the Notch signal are closely related to the occurrence of genetic and autoimmune diseases, and the development of neoplasms, including breast, pancreatic and endometrial cancer [50,51]. Notch receptors are activated by contact with a ligand (Jagged or Delta) from a neighbouring cell. After ligand binding, the receptor undergoes proteolytic processing, as a result of which the NICD (notch intracellular domain) is detached from it and travels to the cell nucleus, where it acts as a transcription factor. In the nucleus, NICD forms a complex with the DNA binding protein of CSL, displacing the histone deacetylase (HDAC)–corepressor (CoR) complex with CSL. Subsequently, components of the activation complex, such as MAML1 and histone acetyltransferase (HAT), are recruited into the NICD-CSL complex, leading to transcriptional activation of Notch target genes. All four Notch receptors were found to be present in the endometrium. Notch 1 and Notch 3 are mainly located in the glandular epithelium, while Notch 4 is only in the stroma of the endometrium [52]. It should be emphasized that the expression of these signalling pathways changes with the phase of the menstrual cycle [52,53] and decreases after menopause [54]. Many years of research on the activity of individual receptors and ligands of this evolutionarily old signalling pathway in endometrial cancer have provided different, sometimes contradictory, results. However, most authors agree that both mRNA and Notch 4 protein expression are reduced [52,54–61]. The role of leptin in the regulation of Notch pathways has been well established in breast cancer, where, together with IL-1, Notch upregulates ligands, receptors, and relevant genes, thus enhancing the induction of proliferation and migration of cancer cells, as well as chemoresistance [62–66]. In a study on the HEC-1A and Ishikawa cell lines (type I endometrial cancer), and KLE and An3Ca (type II endometrial cancer), the same crosstalk of leptin and IL-1 was associated with greater invasiveness and chemoresistance by inducing all Notch receptors (NOTCH1-4), their ligands, JAG1 and DLL4, and downstream effectors (survivin and HEY2). This mechanism is described in both types of endometrial cancer, although it is stronger in type II, in which leptin increased the proliferation of cancer cells [67]. Thus, leptin becomes the cytokine responsible for the proliferation of endometrial cancer. Therefore, the implementation of a treatment that reduces the blood levels of leptin or inhibits the cascades of signalling dependent on this adipokine may limit the development of neoplastic disease [68].

4. The Role of Adiponectin in the Development of Endometrial Cancer

Epidemiological studies confirm the association between low adiponectin levels, obesity, and increased incidence of cancer [69]. The European Prospective Investigation into Cancer and Nutrition (EPIC) study found that lower plasma levels of adiponectin in peri- and postmenopausal women predispose them to an increased risk of developing endometrial cancer, regardless of BMI status and other obesity-related risk factors, such as circulating C-peptide levels, sex hormones, IGF-1 binding protein (IGFBP-1), and peri- and postmenopausal IGFBP-2 [70].

Low levels of adiponectin measured in blood serum (<8 mg/L) are associated with greater aggressiveness of the neoplastic process, more frequent metastases in the lymph nodes, and a low degree of histopathological differentiation [71]. It should be emphasized that an elevated leptin-to-adiponectin ratio is also an independent factor in the development of endometrial cancer [72]. In contrast, high preoperative levels of adiponectin are a positive prognostic factor, in terms of disease-free time and overall survival, in endometrial cancer patients [73].

The suppressor protein PTEN is a lipid phosphatase with dual specificity for both proteins and lipids, being a negative regulator of the PI3K-Akt-mTOR signal axis, thereby controlling cell survival, proliferation, and growth processes. This protein plays a key role in silencing the stimulation transmitted from EGFR, HER-2 and IGFR receptors to

the Akt signalling cascade. Somatic mutation of *PTEN*, leading to complete or partial loss of the *PTEN* protein, results in the development of a neoplastic process characterized by high dynamics of tumour growth, worse prognosis, and faster clinical manifestation of the proliferative process. A study conducted in an animal model confirmed that a heterozygous *PTEN* mutation leads to the formation of endometrial cancer in as many as 30% of cases, but this effect may be exacerbated by adiponectin deficiency, due to activation of the mitogen-activated protein kinase (MAPK) pathway [74].

The antiproliferative effect of adiponectin is related to the presence and functionality of AdipoR receptors, as well as the expression of the *LKB1* suppressor gene [32,75]. Studies on endometrial cancer cell lines have shown that knockdown of the *LKB1* gene abolishes adiponectin-induced reductions in cellular activity, including cell proliferation, colony formation, adhesion, and invasion, in endometrial cancer cell lines [76]. *LKB1*-mediated adiponectin signalling may also interact with the protein *PTEN* involved in the regulation of the cell cycle. It should be noted that a large proportion of endometrial neoplasms contain mutations in *PTEN*. Moreover, the inhibition of cell cycle regulators, cyclin D1 and E2, as well as ERK1/2 and Akt signalling proteins, by adiponectin confirms its antiproliferative activity [77]. In studies on endometrial cancer cell lines, adiponectin has an inhibitory effect on the signalling pathways stimulated by leptin, which makes its antiproliferative and pro-apoptotic effects stronger [78].

5. The Role of Oestrogens in the Development of Endometrial Cancer

Oestrogens play a mitogenic role in the normal endometrium, which consists of the functional and basal layers. The functional layer covers two-thirds of the thickness of the endometrium and peels off at the end of the luteal phase of the menstrual cycle. It separates from the basal layer, which remains intact and forms the basis for endometrial regeneration. The ovaries are the main source of oestrogen, and the levels of oestrogen fluctuate with the phases of the menstrual cycle. In the follicular phase, oestrogens, i.e., estrone and 17 β -oestradiol, are produced by the granulosa cells of a growing ovarian follicle and lead to the growth of the endometrium [79], mainly its glands, which are initially thin, tubular and lined with low cells of the cylindrical epithelium. With time, as a result of numerous mitotic divisions, the adjacent glands fuse with each other. The highest levels of oestradiol are reached at the end of the follicular phase, and then decline sharply after ovulation. This is followed by a re-rise in oestrogen levels in the middle luteal phase with a decline towards the end of the menstrual cycle.

In the following days, oestrogens are produced by the corpus luteum formed after ovulation, but their levels are already lower and drop significantly just before menstruation [80,81]. In the second phase of the cycle, because of the action of progesterone produced by the corpus luteum, the growth of the endometrium is inhibited, which is associated with a decrease in mitotic activity and DNA synthesis. Progesterone also interferes with the expression of oestrogen receptors, and stimulates the activity of 17 β -hydroxysteroid dehydrogenase and sulfotransferases, which convert oestradiol into estrone sulphate, which is quickly excreted from the cell. An imbalance between the oestrogens that lead to the growth of the endometrium and the progesterone that inhibits them can lead to endometrial hyperplasia and endometrial cancer. In menstruating women, the ovaries are the main source of oestrogen, and after menopause, these hormones mainly come from the extra-glandular conversion of androstenedione and testosterone.

Adipose tissue is a source of enzymes, such as cytochrome P450-dependent aromatase, 11 β -hydroxysteroid dehydrogenase type 1 (11 β HSD 1), and 17 β -hydroxysteroid dehydrogenase (17 β -HSD), involved in the synthesis and metabolism of steroid hormones [82]. Cytochrome P450 and 17 β -HSD-dependent aromatase are highly active in adipose tissue stromal cells and pre-adipocytes. Cytochrome P450-dependent aromatase catalyses the conversion of androstenedione to estrone, and testosterone to oestradiol, while 17 β -HSD is involved in the conversion of androstenedione to testosterone, and estrone to oestradiol [83]. The 17 β -HSD aromatase enzyme activity ratio positively correlates with abdominal obesity.

The percentage of androstenedione converted to oestrogens positively correlates with body weight. Most of the major sex steroids, such as oestradiol and testosterone, circulate in the blood in the form of a protein carrier, sex hormone-binding globulin (SHBG). In obese women, the reduction in SHBG protein leads to an increase in the concentration of free oestrogens, contributing to the development of endometrial hyperplasia and cancer. Thus, obese women have a three times higher risk of developing endometrial cancer [84].

Free oestrogens are active hormones that bind to ER α or ER β receptors to produce a biological effect [85]. These genomic receptors belong to the nuclear receptor family and act as transcription factors, or interact with other transcription factors, such as Sp-1, AP-1 and NF- κ B, to express genes related to cell proliferation, differentiation, and normal function [86,87]. Oestrogen receptors show activity dependent on the binding of the oestrogen ligand, increased amounts of which contribute to the development of endometrial hyperplasia or cancer [88]. Endometrial cancer has been divided into oestrogen-dependent type I, which accounts for approximately 80% of all ER α -predominant cases, and into poorly differentiated and more aggressive type II. Although, for many years, it was believed that type II is oestrogen independent [89,90], more and more experimental studies prove the lack of differences in the expression of oestrogen receptors and the levels of sex hormones between these types of neoplasms, which would suggest estrogenic aetiology in both cases [91,92].

The results of studies on the expression and role of ER β in endometrial cancer are ambiguous. Studies conducted on animal and human models indicate the suppressive role of ER β in the development of endometrial cancer [93–96]. However, an elevated, altered ER β , ER β 5 isoform is found in endometrial cancer, which is associated with the expression of HER 2 and MyBL2 oncogenes [97,98]. The analysis of endometrial cancer data from the Genome Atlas showed that the mean expression level of ER α was 2.9 times higher than ER β [99], which may be due to the higher levels of gonadotrophins in the postmenopausal period. Since ER β acts as a negative regulator of ER α , postmenopausal low expression of ER β may promote the proliferative effects of ER α [93].

ER α and ER β are highly homologous in their DNA-binding domains (DBDs) and possess moderate sequence identity in their ligand-binding domains (LBDs). Within the oestrogen receptor, there are six regions located in five domains. In the regulatory domain, there is an A/B region whose function is to activate TAF-1 transcription. TAF-1 can stimulate transcription without binding to the hormone. The C region, located in the DNA-binding domain, is essential for the activation of transcription. The D region is located between the DNA-binding domain and the hormone-binding domain, and plays an important role in the transport of the cytoplasm-synthesized receptor into the cell nucleus. The E region is in the hormone-binding domain and is responsible for dimerization and the activation of TAF-2 transcription. In addition, it is a binding site for heat shock proteins (hsp90), which may interfere with the membrane transport of the receptor and the dimerization process of DNA binding. The F region modulates gene transcription and influences its conformation. Both ER α and ER β have nearly identical DNA-binding domains, but differ in the ligand-binding domain [86].

The transcriptional activity of oestrogen receptors may be increased by phosphorylation derived from the cAMP and kinase A pathways. Thus, growth factors, such as epidermal growth factor-EGF, IGF-I and transforming growth factor alpha (TGF α), by stimulating phosphorylation of the kinase, may activate oestrogen receptors in the absence of oestrogen. In addition to nuclear receptors ER α and ER β , a G-protein-blinded oestrogen receptor, called GPER or GPR 30, has been identified, which is present within the cell membrane, endoplasmic reticulum and in the cell nucleus [100]. This receptor has a high affinity for oestradiol and a low affinity for estrone and estriol [87,101], with estrone and oestradiol being its agonists and estriol being its antagonist. The GPER receptor mediates non-genomic oestrogen signalling via cAMP or via calcium ions released through it from the endoplasmic reticulum by PLC. By activating the Src protein, GPER promotes the activation of metalloproteinases 2/9, causing EGFR transcription, and by stimulating the

MAPK and PI3K/Akt signalling pathways, it can influence the cell cycle, proliferation, differentiation, apoptosis, migration, and cell invasion. Hence, activation of this receptor may modulate carcinogenesis. Endometrial cancer studies have shown GPER overexpression, which correlated with the aggressiveness and advancement of the neoplastic process, as well as a worse prognosis, regardless of the menopausal status. In the case of non-endometrioid types of tumours, a loss of GPER expression was observed, as compared to the normal endometrium, which correlated with the advancement of the disease, low histopathological differentiation, and low expression of the ER α receptor. However, there are studies showing no differences in GPER expression between endometrial cancer types I and II [102]. It should be noted that in endometrial cancer cells, the expression of GPER, as opposed to ER α [103], is three times lower compared to the normal endometrium. The signalling pathway of oestrogens in carcinogenesis is presented in Figure 2.




Figure 2. Interaction of oestrogens and leptin and their downstream signalling pathways in endometrial cancer. JAK—Janus kinase; STAT3—signal transducer and activator of transcription; PI3K/Akt—phosphatidylinositol 3-kinase/protein kinase B; Lep-R—leptin receptor; ER—oestrogen receptors; GPER—G-protein-blinded oestrogen receptor; EGFR—epidermal growth factor receptor; MMP—metalloproteinase; P—phosphorylation; IP3—inositol 1,4,5-trisphosphate; PLC—phospholipase C; Src—protein tyrosine kinase.

Epidemiological studies have shown obesity to be a major risk factor for the development of endometrial cancer in postmenopausal women [104], as well as being the main source of oestrogen and adipokines during this period of life. Studies carried out on an animal model confirm the mutual synergy of action between oestradiol and leptin in the

aetiology of endometrial cancer development, which is manifested by transactivation of the Ob-R receptor through mutual activation of the JAK-STAT pathway [105], and this is presented in Figure 2. Thus, oestradiol, directly on the endometrium and indirectly increasing the effect of leptin, may contribute to the development of endometrial cancer. However, this hypothesis requires further clinical research.

6. Summary

Many epidemiological studies indicate obesity as an independent risk factor for endometrial cancer in postmenopausal women. Moreover, endometrial cancer appears in about 4% of women at reproductive age [2], which can create difficulties in identifying adequate treatment and fertility preservation [106], which is why these patients very often need multidisciplinary care, such as oncological and psychological [107]. A better understanding of molecular biology, particularly regarding the genetics of tumours, can improve the selection of patients who may be eligible for fertility-sparing treatment [108,109]. Further studies of leptin and adiponectin signalling pathways and their “cross talk” with oestradiol may allow for the development of an effective targeted therapy in endometrial cancer.

Author Contributions: Conceptualization, A.S.-J.; data curation, A.S.-J., A.L. and M.J.-K.; writing—original draft preparation, A.S.-J., A.L. and M.J.-K.; writing—review and editing, A.S.-J., A.L., M.J.-K., A.W. and G.J.; visualization, A.S.-J.; supervision, A.W. and G.J.; project administration, A.S.-J. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Bray, F.; Ferlay, J.; Soerjomataram, I.; Siegel, R.L.; Torre, L.A.; Jemal, A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J. Clin.* **2018**, *68*, 394–424. [CrossRef] [PubMed]
2. World Health Organization; International Agency for Research on Cancer (IARC). Global Cancer Observatory. Available online: <https://gco.iarc.fr> (accessed on 20 February 2022).
3. Ali, A.T. Reproductive factors and risk of endometrial cancer. *Int. J. Gynecol. Cancer* **2014**, *21*, 384–393. [CrossRef] [PubMed]
4. Setiawan, V.W.; Yang, H.P.; Pike, M.C.; McCann, S.E.; Yu, H.; Xiang, Y.B.; Wolk, A.; Wentzensen, N.; Weiss, N.S.; Webb, P.M.; et al. Type I and II endometrial cancers: Have they different risk factors? *J. Clin. Oncol.* **2013**, *31*, 2607–2618. [CrossRef] [PubMed]
5. Perez-Hernandez, A.I.; Catalan, V.; Gomez-Ambrosi, J.; Rodríguez, A.; Frühbeck, G. Mechanisms linking excess adiposity and carcinogenesis promotion. *Front. Endocrinol.* **2014**, *5*, 65.
6. Lee, C.H.; Woo, Y.C.; Wang, Y.; Yeung, C.Y.; Xu, A.; Lam, K.S. Obesity, adipokines and cancer: An update. *Clin. Endocrinol. (Oxf)* **2015**, *83*, 147–156. [CrossRef]
7. Guadagni, F.; Roselli, M.; Martini, F.; Spila, A.; Riondino, S.; D’Alessandro, R.; Del Monte, G.; Formica, V.; Laudisi, A.; Portarena, I.; et al. Prognostic Significance of Serum Adipokine Levels in Colorectal Cancer Patients. *Anticancer Res.* **2009**, *29*, 3321–3327.
8. Chen, D.C.; Chung, Y.F.; Yeh, Y.T.; Chaung, H.C.; Kuo, F.C.; Fu, O.Y.; Chen, H.Y.; Hou, M.F.; Yuan, S.S.F. Serum adiponectin and leptin levels in Taiwanese breast cancer patients. *Cancer Lett.* **2006**, *237*, 109–114. [CrossRef]
9. Ellis, P.E.; Barron, G.A.; Bermano, G. Adipocytokines and their relationship to endometrial cancer risk: A systematic review and meta-analysis. *Gyn. Oncol.* **2020**, *158*, 507–516. [CrossRef]
10. Zhang, Y.; Proenca, R.; Maffei, M.; Barone, M.; Leopold, L.; Friedman, J.M. Positional cloning of the mouse obese gene and its human homologue. *Nature* **1994**, *372*, 425–432. [CrossRef]
11. Margetic, S.; Gazzola, C.; Pegg, G.G.; Hill, R.A. Leptin: A review of its peripheral actions and interactions. *Int. J. Obes. Relat. Metab. Disord.* **2002**, *26*, 1407–1433. [CrossRef]
12. Pérez-Pérez, A.; Sánchez-Jiménez, F.; Vilarinho-García, T.; Sánchez-Margalet, V. Role of Leptin in Inflammation and Vice Versa. *Int. J. Mol. Sci.* **2020**, *21*, 5887. [CrossRef] [PubMed]
13. Gruzdeva, O.; Borodkina, D.; Uchasova, E.; Dyleva, Y.; Barbarash, O. Leptin resistance: Underlying mechanisms and diagnosis. *Diabetes Metab. Syndr. Obes. Targets Ther.* **2019**, *12*, 191–198. [CrossRef] [PubMed]
14. Park, J.; Morley, T.S.; Kim, M.; Clegg, D.J.; Scherer, P.E. Obesity and cancer—Mechanisms underlying tumour progression and recurrence. *Nat. Rev. Endocrinol.* **2014**, *10*, 455–465. [CrossRef] [PubMed]

15. Boroń, D.; Nowakowski, R.; Grabarek, B.O.; Zmarzły, N.; Opławski, M. Expression Pattern of Leptin and Its Receptors in Endometrioid Endometrial Cancer. *J. Clin. Med.* **2021**, *10*, 2787. [[CrossRef](#)]
16. Ahima, R.S. Adipose tissue as an endocrine organ. *Obesity (Silver Spring)* **2006**, *14*, 242–249. [[CrossRef](#)]
17. Bjursell, M.; Ahnmark, A.; Bohlooly, Y.M.; William-Olsson, L.; Rhedin, M.; Peng, X.R.; Ploj, K.; Gerdin, A.K.; Arnerup, G.; Elmgren, A.; et al. Opposing Effects of Adiponectin Receptors 1 and 2 on Energy Metabolism. *Diabetes* **2007**, *56*, 583–593. [[CrossRef](#)]
18. Mao, X.; Kikani, C.K.; Riojas, R.A.; Langlais, P.; Wang, L.; Ramos, F.J.; Fang, Q.; Christ-Roberts, C.Y.; Hong, J.Y.; Kim, R.Y.; et al. APPL1 binds to adiponectin receptors and mediates adiponectin signalling and function. *Nat. Cell. Biol.* **2006**, *8*, 516–523. [[CrossRef](#)]
19. Chen, J.; Tan, B.; Karteris, E.; Zervou, S.; Digby, J.; Hillhouse, E.; Vatish, M.; Randevara, H.S. Secretion of adiponectin by human placenta: Differential modulation of adiponectin and its receptors by cytokines. *Diabetologia* **2006**, *49*, 1292–1302. [[CrossRef](#)]
20. Yamauchi, N.; Takazawa, Y.; Maeda, D.; Hibiya, T.; Tanaka, M.; Iwababu, M.; Okada-Iwababu, M.; Yamauchi, T.; Kadowaki, T.; Fukayama, M. Expression levels of adiponectin receptors are decreased in human endometrial adenocarcinoma tissues. *Int. J. Gynecol. Pathol.* **2012**, *31*, 352–357. [[CrossRef](#)]
21. Yamauchi, T.; Hara, K.; Kubota, N.; Terauchi, Y.; Tobe, K.; Froguel, P.; Nagai, R.; Kadowaki, T. Dual roles of adiponectin/Acrp30 in vivo as an anti-diabetic and anti-atherogenic adipokine. *Curr. Drug Targets-Immune Endocr. Metab. Disord.* **2003**, *3*, 243–254. [[CrossRef](#)]
22. Wang, Y.; Lam, K.S.; Xu, J.Y.; Lu, G.; Xu, L.Y.; Cooper, G.J.; Xu, A. Adiponectin inhibits cell proliferation by interacting with several growth factors in an oligomerization-dependent manner. *J. Biol. Chem.* **2005**, *280*, 18341–18347. [[CrossRef](#)] [[PubMed](#)]
23. Tilg, H.; Moschen, A.R. Adipocytokines: Mediators linking adipose tissue, inflammation and immunity. *Nat. Rev. Immunol.* **2006**, *6*, 772–783. [[CrossRef](#)] [[PubMed](#)]
24. Iwababu, M.; Yamauchi, T.; Okada-Iwababu, M.; Sato, K.; Nakagawa, T.; Funata, M.; Yamaguchi, M.; Namiki, S.; Nakayama, R.; Tabata, M. Adiponectin and AdipoR1 regulate PGC-1 α and mitochondria by Ca²⁺ and AMPK/SIRT1. *Nature* **2010**, *464*, 1313–1319. [[CrossRef](#)] [[PubMed](#)]
25. Miyazaki, T.; Bub, J.D.; Uzuki, M.; Iwamoto, Y. Adiponectin activates c-Jun NH2-terminal kinase and inhibits signal transducer and activator of transcription 3. *Biochem. Biophys. Res. Commun.* **2005**, *333*, 79–87. [[CrossRef](#)] [[PubMed](#)]
26. Pearson, G.; Robinson, F.; Beers Gibson, T.; Xu, B.E.; Karandikar, M.; Berman, K.; Cobb, M.H. Mitogen-activated protein (MAP) kinase pathways: Regulation and physiological functions. *Endocr. Rev.* **2001**, *22*, 153–183. [[PubMed](#)]
27. Kadowaki, T.; Yamauchi, T. Adiponectin and adiponectin receptors. *Endocr. Rev.* **2005**, *26*, 439–451. [[CrossRef](#)] [[PubMed](#)]
28. Inoki, K.; Zhu, T.; Guan, K.L. TSC2 mediates cellular energy response to control cell growth and survival. *Cell* **2003**, *115*, 577–590. [[CrossRef](#)]
29. Luo, Z.; Saha, A.K.; Xiang, X.; Ruderman, N.B. AMPK, the metabolic syndrome and cancer. *Trends Pharmacol. Sci.* **2005**, *26*, 69–76. [[CrossRef](#)]
30. Rattan, R.; Giri, S.; Singh, A.K.; Singh, I. 5-Aminoimidazole-4-carboxamide-1- β -D-ribofuranoside inhibits cancer cell proliferation in vitro and in vivo via AMP-activated protein kinase. *J. Biol. Chem.* **2005**, *280*, 582–593. [[CrossRef](#)]
31. Hardie, D.G.; Alessi, D.R. LKB1 and AMPK and the cancer-metabolism link—ten years after. *BMC Biol.* **2013**, *11*, 36. [[CrossRef](#)]
32. Gormand, A.; Henriksson, E.; Ström, K.; Jensen, T.E.; Sakamoto, K.; Göransson, O. Regulation of AMP-activated protein kinase by LKB1 and CaMKK in adipocytes. *J. Cell Biochem.* **2011**, *112*, 1364–1375. [[CrossRef](#)] [[PubMed](#)]
33. Hardie, D.G. The LKB1-AMPK pathway—friend or foe in cancer? *Cancer Cell* **2013**, *23*, 131–132. [[CrossRef](#)] [[PubMed](#)]
34. Goodwin, J.M.; Svensson, R.U.; Lou, H.J.; Winslow, M.M.; Turk, B.E.; Shaw, R.J. An AMPK-independent signaling pathway downstream of the LKB1 tumor suppressor controls Snail1 and metastatic potential. *Mol. Cell* **2014**, *55*, 436–450. [[CrossRef](#)] [[PubMed](#)]
35. Dieudonne, M.N.; Bussiere, M.; Dos Santos, E.; Leneuve, M.C.; Giudicelli, Y.; Pecquery, R. Adiponectin mediates antiproliferative and apoptotic responses in human MCF7 breast cancer cells. *Biochem. Biophys. Res. Commun.* **2006**, *345*, 271–279. [[CrossRef](#)]
36. Ouchi, N.; Kihara, S.; Arita, Y.; Okamoto, Y.; Maeda, K.; Kuriyama, H.; Hotta, K.; Nishida, M.; Takahashi, M.; Muraguchi, M. Adiponectin, an adipocyte-derived plasma protein, inhibits endothelial NF- κ B signaling through a cAMP-dependent pathway. *Circulation* **2000**, *102*, 1296–1301. [[CrossRef](#)]
37. Matsuzawa, Y. Establishment of a Concept of Visceral Fat Syndrome and Discovery of Adiponectin. *Proc. Jpn. Acad. Ser. B Phys. Biol. Sci.* **2010**, *86*, 131–140. [[CrossRef](#)]
38. Sivridis, E.; Giatromanolaki, A.; Gatter, K.C.; Harris, A.L.; Koukourakis, M.I. Association of hypoxia-inducible factors 1 α and 2 α with activated angiogenic pathways and prognosis in patients with endometrial carcinoma. *Cancer* **2002**, *95*, 1055–1063. [[CrossRef](#)]
39. Grosfeld, A.; Andre, J.; Hauguel-de Mouzon, S.; Berra, E.; Pouyssegur, J.; Guerre-Millo, M. Hypoxia-inducible factor 1 transactivates the human leptin gene promoter. *J. Biol. Chem.* **2002**, *277*, 42953–42957. [[CrossRef](#)]
40. Ambrosini, G.; Nath, A.K.; Sierra-Honigmann, M.R.; Flores-Riveros, J. Transcriptional activation of the human leptin gene in response to hypoxia: Involvement of hypoxia-inducible factor 1. *J. Biol. Chem.* **2002**, *277*, 34601–34609. [[CrossRef](#)]
41. Koda, M.; Sulkowska, M.; Wincewicz, A.; Kanczuga-Koda, L.; Musiatowicz, B.; Szymanska, M.; Sulkowski, S. Expression of leptin, leptin receptor, and hypoxia-inducible factor 1 α in human endometrial cancer. *Ann. N. Y. Acad. Sci.* **2007**, *1095*, 90–98. [[CrossRef](#)]

42. Wincewicz, A.; Koda, M.; Sulkowska, M.; Kanczuga-Koda, L.; Sulkowski, S. Comparison of STAT3 with HIF-1 α , Ob and ObR expressions in human endometrioid adenocarcinomas. *Tissue Cell* **2008**, *40*, 405–410. [[CrossRef](#)] [[PubMed](#)]
43. Deng, J.Y.; Sun, D.; Liu, X.Y.; Pan, Y.; Liang, H. STAT-3 correlates with lymph node metastasis and cell survival in gastric cancer. *World J. Gastroenterol.* **2010**, *16*, 5380–5387. [[CrossRef](#)] [[PubMed](#)]
44. Badr, G.; Mohany, M.; Abu-Tarboush, F. Thymoquinone decreases F-actin polymerization and the proliferation of human multiple myeloma cells by suppressing STAT3 phosphorylation and Bcl2/Bcl-XL expression. *Lipids Health Dis.* **2011**, *10*, 236. [[CrossRef](#)] [[PubMed](#)]
45. Deng, J.; Grande, F.; Neamati, N. Small molecule inhibitors of STAT3 signaling pathway. *Curr. Cancer Drug Targets* **2007**, *7*, 91–107. [[CrossRef](#)] [[PubMed](#)]
46. Gao, J.; Tian, J.; Lv, Y.; Shi, F.; Kong, F.; Shi, H.; Zhao, L. Leptin induces functional activation of cyclooxygenase-2 through JAK2/STAT3, MAPK/ERK, and PI3K/AKT pathways in human endometrial cancer cells. *Cancer Sci.* **2009**, *100*, 389–395. [[CrossRef](#)]
47. Chen, C.L.; Hsieh, F.C.; Lieblein, J.C.; Brown, J.; Chan, C.; Wallace, J.A.; Cheng, G.; Hall, B.M.; Lin, J. Stat3 activation in human endometrial and cervical cancers. *Br. J. Cancer* **2007**, *96*, 591–599. [[CrossRef](#)]
48. Zhang, Y.; Liu, L.; Li, C.; Ai, H. Correlation analysis between the expressions of leptin and its receptor (ObR) and clinicopathology in endometrial cancer. *Cancer Biomark.* **2014**, *14*, 353–359. [[CrossRef](#)]
49. Cymbaluk-Płoska, A.; Chudecka-Głaz, A.; Jagodzińska, A.; Pius-Sadowska, E.; Sompolska-Rzechuła, A.; Machaliński, B.; Menkiszak, J. Evaluation of biologically active substances promoting the development of or protecting against endometrial cancer. *Onco. Targets Ther.* **2018**, *11*, 1363–1372. [[CrossRef](#)]
50. Lipsey, C.C.; Harbuzariu, A.; Daley-Brown, D.; Gonzalez-Perez, R.R. Oncogenic role of leptin and Notch interleukin-1 leptin crosstalk outcome in cancer. *World J. Methodol.* **2016**, *6*, 43–55. [[CrossRef](#)]
51. Andersson, E.R.; Sandberg, R.; Lendahl, U. Notch signaling: Simplicity in design, versatility in function. *Development* **2011**, *138*, 3593–3612. [[CrossRef](#)]
52. Jonusiene, V.; Sasnauskiene, A. Notch and Endometrial Cancer. *Adv. Exp. Med. Biol.* **2021**, *1287*, 47–57. [[PubMed](#)]
53. Mitsuhashi, Y.; Horiuchi, A.; Miyamoto, T.; Kashima, H.; Suzuki, A.; Shiozawa, T. Prognostic significance of Notch signalling molecules and their involvement in the invasiveness of endometrial carcinoma cells. *Histopathology* **2012**, *60*, 826–837. [[CrossRef](#)] [[PubMed](#)]
54. Mikhailik, A.; Mazella, J.; Liang, S.; Tseng, L. Notch ligand-dependent gene expression in human endometrial stromal cells. *Biochem. Biophys. Res. Commun.* **2009**, *388*, 479–482. [[CrossRef](#)] [[PubMed](#)]
55. Cobellis, L.; Caprio, F.; Trabucco, E.; Mastrogiacomo, A.; Coppola, G.; Manente, L.; Colacurci, N.; De Falco, M.; De Luca, A. The pattern of expression of Notch protein members in normal and pathological endometrium. *J. Anat.* **2008**, *213*, 464–472. [[CrossRef](#)] [[PubMed](#)]
56. Suzuki, T.; Aoki, D.; Susumu, N.; Udagawa, Y.; Nozawa, S. Imbalanced expression of TAN-1 and human Notch4 in endometrial cancers. *Int. J. Oncol.* **2000**, *17*, 1131–1139. [[CrossRef](#)]
57. Mori, M.; Miyamoto, T.; Ohno, S.; Miyake, Y.; Sakaguchi, T.; Ohno, E. Diagnostic utility of notch-1 immuno-cytochemistry in endometrial cytology. *Acta Cytol.* **2012**, *56*, 166–170. [[CrossRef](#)]
58. Sasnauskiene, A.; Jonusiene, V.; Krikstaponiene, A.; Butkyte, S.; Dabkeviciene, D.; Kanopiene, D.; Kazbariene, B.; Didziapetriene, J. NOTCH1, NOTCH3, NOTCH4, and JAG2 protein levels in human endometrial cancer. *Medicina* **2014**, *50*, 14–18. [[CrossRef](#)]
59. Townsend, M.H.; Ence, Z.E.; Felsted, A.M.; Parker, A.C.; Piccolo, S.R.; Robison, R.A.; O'Neill, K.L. Potential new biomarkers for endometrial cancer. *Cancer Cell Int.* **2019**, *19*, 19. [[CrossRef](#)]
60. Jonusiene, V.; Sasnauskiene, A.; Lachej, N.; Kanopiene, D.; Dabkeviciene, D.; Sasnauskiene, S.; Kazbariene, B.; Didziapetriene, J. Down-regulated expression of Notch signaling molecules in human endometrial cancer. *Med. Oncol.* **2013**, *30*, 438. [[CrossRef](#)]
61. Williams, E.; Villar-Prados, A.; Bowser, J.; Broaddus, R.; Gladden, A.B. Loss of polarity alters proliferation and differentiation in low-grade endometrial cancers by disrupting Notch signaling. *PLoS ONE* **2017**, *12*, e0189081. [[CrossRef](#)]
62. Wang, J.; Zhang, K.; Liu, Z.; Wang, T.; Shi, F.; Zhang, Y.; Su, J.; Jia, Y. Upregulated delta-like protein 3 expression is a diagnostic and prognostic marker in endometrial cancer: A retrospective study. *Medicine* **2018**, *97*, e13442. [[CrossRef](#)] [[PubMed](#)]
63. Guo, S.; Gonzalez-Perez, R.R. Notch, IL-1 and Leptin Crosstalk Outcome (NILCO) Is Critical for Leptin- Induced Proliferation, Migration and VEGF/VEGFR-2 Expression in Breast Cancer. *PLoS ONE* **2011**, *6*, e21467. [[CrossRef](#)] [[PubMed](#)]
64. Harbuzariu, A.; Rampoldi, A.; Daley-Brown, D.S.; Candelaria, P.; Harmon, T.L.; Lipsey, C.C.; Beech, D.J.; Quarshie, A.; Ilies, G.O.; Gonzalez-Perez, R.R. Leptin-Notch signaling axis is involved in pancreatic cancer progression. *Oncotarget* **2017**, *8*, 7740–7752. [[CrossRef](#)] [[PubMed](#)]
65. Harbuzariu, A.; Gonzalez-Perez, R.R. Leptin-Notch axis impairs 5-fluorouracil effects on pancreatic cancer. *Oncotarget* **2018**, *9*, 18239–18253. [[CrossRef](#)]
66. Carino, C.; Olawaiye, A.B.; Cherfils, S.; Serikawa, T.; Lynch, M.P.; Rueda, B.R.; Gonzalez, R.R. Leptin regulation of proangiogenic molecules in benign and cancerous endometrial cells. *Int. J. Cancer* **2008**, *123*, 2782–2790. [[CrossRef](#)]
67. Battle, M.; Gillespie, C.; Quarshie, A.; Lanier, V.; Harmon, T.; Wilson, K.; Torroella-Kouri, M.; Gonzalez-Perez, R.R. Obesity Induced a Leptin-Notch Signaling Axis in Breast Cancer. *Int. J. Cancer* **2014**, *134*, 1605–1616. [[CrossRef](#)]

68. Daley-Brown, D.; Harbuzariu, A.; Kurian, A.A.; Oprea-Ilies, G.; Gonzalez-Perez, R.R. Leptin-induced Notch and IL-1 signaling crosstalk in endometrial adenocarcinoma is associated with invasiveness and chemoresistance. *World J. Clin. Oncol.* **2019**, *10*, 222–233. [[CrossRef](#)]
69. Yunusova, N.V.; Kondakova, I.V.; Kolomiets, L.A.; Afanas'ev, S.G.; Chernyshova, A.L.; Kudryavtsev, I.V.; Tsydenova, A.A. Molecular targets for the therapy of cancer associated with metabolic syndrome (transcription and growth factors). *Asia-Pac. J. Clin. Oncol.* **2018**, *14*, 134–140. [[CrossRef](#)]
70. Kelesidis, I.; Kelesidis, T.; Mantzoros, C.S. Adiponectin and cancer: A systematic review. *Br. J. Cancer* **2006**, *94*, 1221–1225. [[CrossRef](#)]
71. Cust, A.E.; Kaaks, R.; Friedenreich, C.; Bonnet, F.; Laville, M.; Lukanova, A.; Rinaldi, S.; Dossus, L.; Slimani, N.; Lundin, E.; et al. Plasma adiponectin levels and endometrial cancer risk in pre- and postmenopausal women. *J. Clin. Endocrinol. Metab.* **2007**, *92*, 255–263. [[CrossRef](#)]
72. Zhang, L.; Wen, K.; Han, X.; Liu, R.; Qu, Q. Adiponectin mediates antiproliferative and apoptotic responses in endometrial carcinoma by the AdipoRs/AMPK pathway. *Gynecol. Oncol.* **2015**, *137*, 311–320. [[CrossRef](#)] [[PubMed](#)]
73. Ashizawa, N.; Yahata, T.; Quan, J.; Adachi, S.; Yoshihara, K.; Tanaka, K. Serum leptin-adiponectin ratio and endometrial cancer risk in postmenopausal female subject. *Gynecol. Oncol.* **2010**, *119*, 65–69. [[CrossRef](#)] [[PubMed](#)]
74. Morielli, A.R.; Kokts-Porietis, R.L.; Benham, J.L. Associations of insulin resistance and inflammatory biomarkers with endometrial cancer survival: The Alberta endometrial cancer cohort study. *Cancer Med.* **2022**, *11*, 1701–1711. [[CrossRef](#)] [[PubMed](#)]
75. Yan, Y.; Shi, H.; Zhao, Z.; Wang, S.; Zhou, S.; Mu, Y.; Ding, N.; Lai, Y.; Zhao, A.Z.; Cheng, L.; et al. Adiponectin deficiency promotes endometrial carcinoma pathogenesis and development via activation of mitogen-activated protein kinase. *J. Pathol.* **2022**, *257*, 146–157. [[CrossRef](#)]
76. Taliaferro-Smith, L.; Nagalingam, A.; Zhong, D.; Zhou, W.; Saxena, N.K.; Sharma, D. LKB1 is required for adiponectin-mediated modulation of AMPK-S6K axis and inhibition of migration and invasion of breast cancer cells. *Oncogene* **2009**, *28*, 2621–2633. [[CrossRef](#)]
77. Moon, H.S.; Chamberland, J.P.; Aronis, K.; Tseleni-Balafouta, S.; Mantzoros, C.S. Direct role of adiponectin and adiponectin receptors in endometrial cancer: In vitro and ex vivo studies in humans. *Mol. Cancer Ther.* **2011**, *10*, 2234–2243. [[CrossRef](#)]
78. Barbe, A.; Bongrani, A.; Mellouk, N.; Estienne, A.; Kurowska, P.; Grandhaye, J.; Elfassy, Y.; Levy, R.; Rak, A.; Froment, P.; et al. Mechanisms of adiponectin action in fertility: An overview from gametogenesis to gestation in humans and animal models in normal and pathological conditions. *Int. J. Mol. Sci.* **2019**, *20*, 1526. [[CrossRef](#)]
79. Wu, X.; Yan, Q.; Zhang, Z.; Du, G.; Wan, X. Acrp30 inhibits leptin-induced metastasis by downregulating the JAK/STAT3 pathway via AMPK activation in aggressive SPEC-2 endometrial cancer cells. *Oncol. Rep.* **2012**, *27*, 1488–1496. [[CrossRef](#)]
80. Maybin, J.A.; Critchley, H.O.D. Menstrual physiology: Implications for endometrial pathology and beyond. *Hum. Reprod. Update* **2015**, *21*, 748–761. [[CrossRef](#)]
81. Williams, C.J.; Erickson, G.F. *Morphology and Physiology of the Ovary*; Endotext; MDText.com, Inc.: South Dartmouth, MA, USA, 2000.
82. Anckaert, E.; Jank, A.; Petzold, J.; Rohsmann, F.; Paris, R.; Renggli, M.; Schönfeld, K.; Schiettecatte, J.; Kriner, M. Extensive monitoring of the natural menstrual cycle using the serum biomarkers estradiol, luteinizing hormone and progesterone. *Pract. Lab. Med.* **2021**, *25*, e00211. [[CrossRef](#)]
83. Kershaw, E.E.; Flier, J.S. Adipose tissue as an endocrine organ. *J. Clin. Endocrinol. Metab.* **2004**, *89*, 2548–2556. [[CrossRef](#)] [[PubMed](#)]
84. Meseguer, A.; Puche, C.; Cabero, A. Sex steroid biosynthesis in white adipose tissue. *Horm. Metab. Res.* **2002**, *34*, 731–736. [[CrossRef](#)] [[PubMed](#)]
85. Dobbins, M.; Decorby, K.; Choi, B.C. The association between obesity and cancer risk: A meta-analysis of observational studies from 1985 to 2011. *Prev. Med.* **2013**, *2013*, 680536. [[CrossRef](#)] [[PubMed](#)]
86. Rodriguez, A.C.; Blanchard, Z.; Maurer, K.A.; Gertz, J. Estrogen signaling in endometrial cancer: A key oncogenic pathway with several open questions. *Horm. Cancer* **2019**, *10*, 51–63. [[CrossRef](#)] [[PubMed](#)]
87. Eyster, K.M. Estrogen receptors methods and protocols methods in molecular biology 1366. *Methods Mol. Biol.* **2016**, 18–19.
88. Pakdel, F. Molecular pathways of estrogen receptor action. *Int. J. Mol. Sci.* **2018**, *19*, 2591. [[CrossRef](#)]
89. Yu, K.; Huang, Z.Y.; Xu, X.L.; Li, J.; Fu, X.W.; Deng, S.L. Estrogen receptor function: Impact on the human endometrium. *Front. Endocrinol.* **2022**, *1*, 113. [[CrossRef](#)]
90. Samarthai, N.; Hall, K.; Yeh, I.T. Molecular profiling of endometrial malignancies. *Obstet. Gynecol. Int.* **2010**, *2010*, 162363. [[CrossRef](#)] [[PubMed](#)]
91. Sanderson, P.A.; Critchley, H.O.D.; Williams, A.R.W.; Arends, M.J.; Saunders, P.T.K. New concepts for an old problem: The diagnosis of endometrial hyperplasia. *Hum. Reprod. Update* **2017**, *23*, 232–254. [[CrossRef](#)]
92. Shen, F.; Gao, Y.; Ding, J.; Chen, Q. Is the positivity of estrogen receptor or progesterone receptor different between type 1 and type 2 endometrial cancer? *Oncotarget* **2017**, *8*, 506–511. [[CrossRef](#)]
93. Wan, J.; Gao, Y.; Zeng, K.; Yin, Y.; Zhao, M.; Wei, J.; Chen, Q. The levels of the sex hormones are not different between type 1 and type 2 endometrial cancer. *Sci. Rep.* **2016**, *6*, 39744. [[CrossRef](#)]
94. Zhang, W.; Saji, S.; Mäkinen, S.; Cheng, G.; Jensen, E.V.; Warner, M.; Gustafsson, J.Å. Estrogen receptor (ER) beta, a modulator of ER alpha in the uterus. *Proc. Natl. Acad. Sci. USA* **2000**, *97*, 5936–5941. [[CrossRef](#)]

95. Paul, M.; Cholewa, K.; Mazurek, U.; Witek, A.; Wilczok, T. Estrogen receptor beta delta 6 (ER Beta Delta 6) isoform in human endometrial hyperplasia and adenocarcinoma. *Cancer Investig.* **2004**, *22*, 211–218. [[CrossRef](#)]
96. Smuc, T.; Rizner, T.L. Aberrant pre-receptor regulation of estrogen and progesterone action in endometrial cancer. *Mol. Cell Endocrinol.* **2009**, *301*, 74–82. [[CrossRef](#)]
97. Skrzypczak, M.; Bieche, I.; Szymczak, S.; Tozlu, S.; Lewandowski, S.; Girault, I.; Radwanska, K.; Szczylik, C.; Jakowicki, J.A.; Lidereau, R.; et al. Evaluation of mRNA expression of estrogen receptor beta and its isoforms in human normal and neoplastic endometrium. *Int. J. Cancer* **2004**, *110*, 783–787. [[CrossRef](#)]
98. Häring, J.; Skrzypczak, M.; Stegerer, A.; Latratch, C.; Weber, F.; Goerse, R.; Ortmann, O.; Treeck, O. Estrogen receptor beta transcript variants associate with oncogene expression in endometrial cancer. *Int. J. Mol. Med.* **2012**, *29*, 1127–1136. [[CrossRef](#)]
99. Hewitt, S.C.; Korach, K.S. Estrogen receptors: Structure, mechanisms and function. *Rev. Endocr. Metab. Disord.* **2002**, *3*, 193–200. [[CrossRef](#)]
100. Tang, Z.R.; Zhang, R.; Lian, Z.X.; Deng, S.L.; Yu, K. Estrogen-Receptor Expression and Function in Female Reproductive Disease. *Cells* **2019**, *8*, 1123. [[CrossRef](#)]
101. Prossnitz, E.R.; Arterburn, J.B. International union of basic and clinical pharmacology. XCVII. G protein-coupled estrogen receptor and its pharmacologic modulators. *Pharmacol. Rev.* **2015**, *67*, 505–540. [[CrossRef](#)]
102. Wan, J.; Yin, Y.; Zhao, M.; Shen, F.; Chen, M.; Chen, Q. The positivity of G- protein-coupled receptor-30 (GPR 30), an alternative estrogen receptor is not different between type 1 and type 2 endometrial cancer. *Oncotarget* **2017**, *8*, 90897–90904. [[CrossRef](#)]
103. Carlson, M.J.; Thiel, K.W.; Leslie, K.K. Past, present, and future of hormonal therapy in recurrent endometrial cancer. *Int. J. Women's Health* **2014**, *6*, 429–435.
104. Onstad, M.A.; Schmandt, R.E.; Lu, K.H. Addressing the role of obesity in endometrial cancer risk, prevention, and treatment. *J. Clin. Oncol.* **2016**, *34*, 4225–4230. [[CrossRef](#)]
105. Shetty, A.; Venkatesh, T.; Tsutsumi, R.; Suresh, P.S. Gene expression changes and promoter methylation with the combined effects of estradiol and leptin in uterine tissue of the ovariectomized mice model of menopause. *Mol. Biol. Rep.* **2019**, *47*, 151–168. [[CrossRef](#)]
106. Tanos, P.; Dimitriou, S.; Gullo, G.; Tanos, V. Biomolecular and Genetic Prognostic Factors That Can Facilitate Fertility-Sparing Treatment (FST) Decision Making in Early Stage Endometrial Cancer (ES-EC): A Systematic Review. *Int. J. Mol. Sci.* **2022**, *23*, 2653. [[CrossRef](#)]
107. Rosa, V.L.; Valenti, G.; Sapia, F.; Gullo, G.; María, A.V.; Rapisarda, C. Psychological impact of gynecological diseases: The importance of a multidisciplinary approach. *It. J. Gynaecol. Obstet* **2018**, *30*, 2.
108. Cavaliere, A.F.; Perelli, F.; Zaami, S.; D'Indinosante, M.; Turrini, I.; Giusti, M.; Gullo, G.; Vizzielli, G.; Mattei, A.; Scambia, G.; et al. Fertility Sparing Treatments in Endometrial Cancer Patients: The Potential Role of the New Molecular Classification. *Int. J. Mol. Sci.* **2021**, *22*, 12248. [[CrossRef](#)]
109. Gullo, G.; Etrusco, A.; Cucinella, G.; Perino, A.; Chiantera, V.; Laganà, A.S.; Tomaiuolo, R.; Vitagliano, A.; Giampaolino, P.; Noventa, M.; et al. Fertility-Sparing Approach in Women Affected by Stage I and Low-Grade Endometrial Carcinoma: An Updated Overview. *Int. J. Mol. Sci.* **2021**, *22*, 11825. [[CrossRef](#)]