



Article

Tumor-Localized Administration of α -GalCer to Recruit Invariant Natural Killer T Cells and Enhance Their Antitumor Activity against Solid Tumors

Yan-Ruide Li, Yang Zhou, Matthew Wilson, Adam Kramer, Ryan Hon, Yichen Zhu, Ying Fang and Lili Yang

- ¹ Department of Microbiology, Immunology & Molecular Genetics, University of California, Los Angeles, CA 90095, USA; charlie.li@ucla.edu (Y.-R.L.); zzydcat@g.ucla.edu (Y.Z.); mwilson193@g.ucla.edu (M.W.); akramer931@g.ucla.edu (A.K.); ryanhonchenghao@gmail.com (R.H.); yichenzhujohn@g.ucla.edu (Y.Z.); yingfang1913@g.ucla.edu (Y.F.)
- ² Eli and Edythe Broad Center of Regenerative Medicine and Stem Cell Research, University of California, Los Angeles, CA 90095, USA
- ³ Jonsson Comprehensive Cancer Center, David Geffen School of Medicine, University of California, Los Angeles, CA 90095, USA
- ⁴ Molecular Biology Institute, University of California, Los Angeles, CA 90095, USA
- * Correspondence: liliyang@ucla.edu; Tel.: +13-10-825-8609

Citation: Li, Y.-R.; Zhou, Y.; Wilson, M.; Kramer, A.; Hon, R.; Zhu, Y.; Fang, Y. Tumor-Localized Administration of α -GalCer to Recruit Invariant Natural Killer T Cells and Enhance Their Antitumor Activity against Solid Tumors. *Int. J. Mol. Sci.* **2022**, *23*, 7547. <https://doi.org/10.3390/ijms23147547>

Supplementary Materials

Academic Editor: Walter Fiedler

Received: 2 June 2022

Accepted: 6 July 2022

Published: 7 July 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).




Figure S1. Tissue biodistribution of PBMC-iNKT cells in an A375-CD1d human melanoma xenograft NSG mouse model. Related to Figure 5. Representative images showing iNKT cell load in experimental mice over time. These mice are injected with A375-CD1d tumor cells, α GC and PBMC-iNKT-FG cells. Both ventral and left side views are shown.




Figure S2. Mouse serum cytokine changes in an A375-CD1d human melanoma xenograft NSG mouse model. Related to Figure 5. Levels of human IFN- γ (A) and TNF- α (B) were measured by ELISA on serum collected from mouse peripheral blood on different days after PBMC-iNKT-FG cell infusion. Representative of 2 experiments. Data are presented as the mean \pm SEM. ns, not significant, * $p < 0.05$ by Student's t test.