







**Supplement Table S1.** qPCR probe list.

probe	
<i>Tnfa</i>	Mm00443258_m1
<i>Tgfb1</i>	Mm01178820_m1
<i>Il1b</i>	Mm00434228_m1
<i>Il6</i>	Mm00446190_m1
<i>Il12b</i>	Mm01288989_m1
<i>Il23a</i>	Mm00518984_m1
<i>Il17a</i>	Mm00439618_m1
<i>Il17c</i>	Mm00521397_m1
<i>Il17f</i>	Mm00521423_m1
<i>Il22</i>	Mm01226722_m1
<i>Nlrp3</i>	Mm00840904_m1
<i>Ccl2</i>	Mm00441242_m1
<i>Krt6</i>	Mm00833464_g1
<i>Krt16</i>	Mm01306670_g1
<i>Krt17</i>	Mm00495207_m1
<i>B2m</i>	Mm00437762_m1
<i>Polr2a</i>	Mm00839493_m1

**A****B****C**

**Figure S1.** Body weight, Liver and spleen weights, and serum lipids profile on Day 0. Mice were fed a NASH or 7KC diet for three weeks, and samples were collected on Day 0. **(A)** Body weight. **(B)** Liver and spleen weights. **(C)** Serum lipid profile Results are presented as mean  $\pm$  SD, and *p*-values were calculated using Student's *t*-test. \* *p* < 0.05, Control vs. 7KC. *n* = 6, Control; *n* = 6, 7KC.



**Figure S2.** RNA sequence analysis of *Abcg1* in liver on Day 0.