

Anti-obesity and Anti-hyperglycemic Effects of *Meretrix lusoria* Protamex Hydrolysate in *ob/ob* mice

Min Ju Kim ^{1,†}, Ramakrishna Chilakala ^{1,†}, Hee Geun Jo ¹, Seung-Jae Lee ², Dong-Sung Lee ³, Sun Hee Cheong ^{1,*}

¹ Department of Marine Bio-Food Sciences, College of Fisheries and Ocean Sciences, Chonnam National University, Yeosu 59626, Korea.

modori96k@naver.com (M.J.K.); ramach2006@gmail.com (R.C.); altkwh@naver.com (H.G.J.)

² Immunoregulatory Material Research Center, Korea Research Institute of Bioscience and Biotechnology (KRIBB), Jeongeup 56212, Korea; seung99@kribb.re.kr

³ Department of Pharmacy, College of Pharmacy, Chosun University, Dong-gu, Gwangju, 61452, Korea
dslee2771@chosun.ac.kr

* Correspondence: sunny3843@jnu.ac.kr; Tel.: (Phone: +82-61-659-7215; Fax: +82-61-659-7219)

† These authors contributed equally to this study.




Figure S1. Effect of MLPH fraction on (A) hepatic triglycerides; (B) serum HDL cholesterol, and (C) serum LDL cholesterol. Each value represents the mean \pm SEM ($n=6$). ^{a-b} Values not sharing a common letter are significantly different at $p < 0.05$ by Tukey's multiple comparison test, ^{NS} : not significant.




Figure S2. Effects of MLPH fraction on (A) mouse body weight changes during 6 weeks of treatment; metabolic parameters: (B) serum protein, (C) serum albumin, (D) blood urea nitrogen (BUN), and (E) creatinine. Each value represents the mean \pm SEM (n=6). ^{a-c} Values not sharing a common letter are significantly different at $p < 0.05$ by Tukey's multiple comparison test, ^{NS} : not significant.