

Sirt6 Activation Ameliorates Inflammatory Bone Loss in Ligature-Induced Periodontitis in Mice

Myung Jin Lee ^{1,†}, Hyang Hwa Ryu ^{2,†}, Jae Won Hwang ², Jung Ryul Kim ³, Eui-Sic Cho ⁴, Jin Kyeong Choi ^{5,*} and Young Jae Moon ^{2,3,*}

1 Department of Chemistry, Case Western Reserve University, Cleveland, OH 44106, USA; jinalee939@gmail.com

2 Department of Biochemistry and Molecular Biology, Jeonbuk National University Medical School, Jeonju 54896, Republic of Korea; yy-ryu@hanmail.net (H.H.R.); toriake11@naver.com (J.W.H.)

3 Department of Orthopaedic Surgery, Jeonbuk National University Medical School and Hospital, Jeonju 54896, Republic of Korea; jrkeen@jbnu.ac.kr

4 Cluster for Craniofacial Development and Regeneration Research, Institute of Oral Biosciences, Jeonbuk National University School of Dentistry, Jeonju 54896, Republic of Korea; oasis@jbnu.ac.kr


5 Department of Immunology, Jeonbuk National University Medical School, Jeonju 54907, Republic of Korea

* Correspondence: jkchoi@jbnu.ac.kr (J.K.C.); yjmoonos@jbnu.ac.kr (Y.J.M.)

† These authors contributed equally to this work.

Keywords: alveolar bone loss; inflammation; MDL801; mouse models; osteoclast; periodontitis; Sirt6

1. Supplementary Figure



Supplement Figure S1. Systemic inflammatory cytokines did not change in the ligature-induced periodontitis model. ELISA analysis of IL-1 β , IL-6, RANKL, and OPG in serum of control sham, control ligature, Sirt6Tg sham, and Sirt6Tg ligature (n=3–5).