



Correction

Correction: Mesnard et al. Eliminating Synaptic Ribbons from Rods and Cones Halves the Releasable Vesicle Pool and Slows Down Replenishment. *Int. J. Mol. Sci.* 2022, 23, 6429

Chris S. Mesnard^{1,2}, Cody L. Barta¹, Asia L. Sladek¹, David Zenisek³ and Wallace B. Thoreson^{1,2,*}

¹ Department of Ophthalmology and Visual Sciences, Truhlsen Eye Institute, University of Nebraska Medical Center, Omaha, NE 68198, USA

² Pharmacology and Experimental Neuroscience, University of Nebraska Medical Center, Omaha, NE 68198, USA

³ Department of Cellular and Molecular Physiology, Yale University, New Haven, CT 06510, USA

* Correspondence: wbthoreson@unmc.edu; Tel.: +1-402-559-4076

The authors wish to make the following corrections to this paper [1]: In the original publication, there was a mistake in Figure 6 as published. We reversed the data in panels C and D, plotting B-wave latency data as A-wave amplitude in panel C and A-wave amplitude as B-wave latency in panel D. The corrected Figure 6 appears below.



Citation: Mesnard, C.S.; Barta, C.L.; Sladek, A.L.; Zenisek, D.; Thoreson, W.B. Correction: Mesnard et al. Eliminating Synaptic Ribbons from Rods and Cones Halves the Releasable Vesicle Pool and Slows Down Replenishment. *Int. J. Mol. Sci.* 2022, 23, 6429. *Int. J. Mol. Sci.* 2023, 24, 1561. <https://doi.org/10.3390/ijms24021561>

Received: 1 December 2022

Accepted: 5 December 2022

Published: 13 January 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).




Figure 6. Photopic ERG b-waves were reduced equally in the cone^{RibeyeCKO}, rod^{RibeyeCKO}, and Ribeye KI retinas. (A) Example waveforms. (B) B-wave amplitude as a function of flash intensity measured in the C57Bl6J ($n = 5$ mice), cone^{RibeyeCKO} ($n = 5$), rod^{RibeyeCKO} ($n = 5$), and Ribeye KI retinas ($n = 4$). (C) A-wave amplitude as a function of intensity. (D) B-wave latency vs. intensity.

The authors apologize for any inconvenience caused and state that the scientific conclusions are unaffected. This correction was approved by the Academic Editor. The original article has been updated.

Reference

1. Mesnard, C.S.; Barta, C.L.; Sladek, A.L.; Zenisek, D.; Thoreson, W.B. Eliminating Synaptic Ribbons from Rods and Cones Halves the Releasable Vesicle Pool and Slows Down Replenishment. *Int. J. Mol. Sci.* **2022**, *23*, 6429. [[CrossRef](#)] [[PubMed](#)]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.