

Supplementary material

Ellagic Acid from Hull Blackberries: Extraction, Purification, and Potential Anticancer Activity

Jialuan Wang ^{1,†}, Fengyi Zhao ^{1,†}, Wenlong Wu ¹, Lianfei Lyu ¹, Weilin Li ^{2,*}
and Chunhong Zhang ^{1,*}

¹ Jiangsu Key Laboratory for the Research and Utilization of Plant Resources, Institute of Botany, Jiangsu Province and Chinese Academy of Sciences (Nanjing Botanical Garden Mem. Sun Yat-Sen), Qian Hu Hou Cun No. 1, Nanjing 210014, China; wjl163youxiang@163.com (J.W.); zhaofengyi92@163.com (F.Z.); 1964wwl@163.com (W.W.); njbglq@163.com (L.L.)

² Co-Innovation Center for Sustainable Forestry in Southern China, College of Forestry, Nanjing Forestry University, 159 Longpan Road, Nanjing 210037, China

* Correspondence: wlli@njfu.edu.cn (W.L.); chzhang@cnbg.net (C.Z.)

[†] These authors contributed equally to this work.




Figure S1. Infrared spectra. Dark blue is the infrared spectrum of EA standard; light blue is the infrared spectrum of experimentally EA extract.




Figure S2. UPLC chromatogram.