

Supplemental Table S1. Patient's characteristics at diagnosis of *de novo* AML (n = 39)

Median age	58 [24 ; 88]
Sex ratio (M/W)	2
Leukocytosis	n (%)
< 4 10 ³ /μL	8 (20)
4-30 10 ³ /μL	17 (44)
30-100 10 ³ /μL	11 (28)
> 100 10 ³ /μL	3 (8)
AML subtype (FAB classification)	n (%)
M0	2 (5)
M1	9 (23)
M2	18 (46)
M4	6 (16)
M5	4 (10)
Bone marrow blasts	n (%)
20-40%	9 (23)
40-60%	11 (28)
60-80%	9 (23)
80-100%	10 (26)
Prognosis (ELN 2017)	n (%)
Favorable	21 (54)
Adverse	18 (46)
Molecular status	n (%)
<i>NPM1</i>	15 (38)
<i>FLT3-ITD</i>	11 (28)
<i>IDH</i>	9 (23)
<i>TET2</i>	4 (10)
<i>CEBPA</i>	4 (10)
<i>RUNX1</i>	1 (3)
<i>TP53</i>	1 (3)
Response to chemotherapy	n (%)
Chemosensitive	20 (51)
Chemoresistant	6 (16)
Non available	13 (33)