

**Supporting information**

**for**

**Synthesis of 3-(pyridin-2-yl)quinazolin-2,4(1*H*,3*H*)-diones via annulation of  
anthranilic esters with *N*-pyridyl ureas**

Svetlana O. Baykova, Kirill K. Geyl, Sergey V. Baykov, Vadim P. Boyarskiy \*

*Institute of Chemistry, Saint Petersburg State University, Universitetskaya Nab., 7/9, 199034  
Saint Petersburg, Russia*



\*Corresponding authors: v.boyarskii@spbu.ru.

## Table of Content


Copies of $^1\text{H}$ , $^{13}\text{C}$ , and $^{19}\text{F}$ NMR spectra.....	3
Table S1 Crystal data and structure refinement for <b>3a</b> . .....	42

## Copies of $^1\text{H}$ , $^{13}\text{C}$ , and $^{19}\text{F}$ NMR spectra


$^1\text{H}$  and  $^{13}\text{C}$  spectra of methyl 6-(3,3-dimethylureido)picolinate **1h**




<sup>1</sup>H and <sup>13</sup>C spectra of 1,1-dimethyl-3-(6-methylquinolin-2-yl)urea **1n**




<sup>1</sup>H and <sup>13</sup>C spectra of 1,1-dimethyl-3-(7-methylquinolin-2-yl)urea **10**




<sup>1</sup>H and <sup>13</sup>C spectra of 3-(6-methoxyquinolin-2-yl)-1,1-dimethylurea **1p**




<sup>1</sup>H and <sup>13</sup>C spectra of 3-(4-methylpyridin-2-yl)quinazoline-2,4(1*H*,3*H*)-dione **3a**




<sup>1</sup>H and <sup>13</sup>C spectra of 3-(pyridin-2-yl)quinazoline-2,4(1*H*,3*H*)-dione **3b**




<sup>1</sup>H and <sup>13</sup>C spectra of 3-(5-methylpyridin-2-yl)quinazoline-2,4(1*H*,3*H*)-dione **3c**




<sup>1</sup>H and <sup>13</sup>C spectra of 3-(6-methylpyridin-2-yl)quinazoline-2,4(1*H*,3*H*)-dione **3d**




<sup>1</sup>H and <sup>13</sup>C spectra of 3-(3,5-dimethylpyridin-2-yl)quinazoline-2,4(1*H*,3*H*)-dione **3e**




<sup>1</sup>H and <sup>13</sup>C spectra of 3-(4-methoxypyridin-2-yl)quinazoline-2,4(1*H*,3*H*)-dione **3f**




<sup>1</sup>H and <sup>13</sup>C spectra of 3-(6-phenylpyridin-2-yl)quinazoline-2,4(1*H*,3*H*)-dione **3g**




<sup>1</sup>H and <sup>13</sup>C spectra of methyl 6-(2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl)picolinate **3h**




<sup>1</sup>H and <sup>13</sup>C spectra of methyl 2-(2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl)isonicotinate **3i**



<sup>1</sup>H and <sup>13</sup>C spectra of 6-(2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl)picolinonitrile **3j**




<sup>1</sup>H and <sup>13</sup>C spectra of 2-(2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl)isonicotinonitrile **3k**




<sup>1</sup>H and <sup>13</sup>C spectra of 3-(quinolin-2-yl)quinazoline-2,4(1*H*,3*H*)-dione **3l**




<sup>1</sup>H and <sup>13</sup>C spectra of 3-(4-methoxyquinolin-2-yl)quinazoline-2,4(1*H*,3*H*)-dione **3m**




<sup>1</sup>H and <sup>13</sup>C spectra of 3-(6-methylquinolin-2-yl)quinazoline-2,4(1*H*,3*H*)-dione **3n**




<sup>1</sup>H and <sup>13</sup>C spectra of 3-(7-methylquinolin-2-yl)quinazoline-2,4(1*H*,3*H*)-dione **3o**




<sup>1</sup>H and <sup>13</sup>C spectra of 3-(6-methoxyquinolin-2-yl)quinazoline-2,4(1*H*,3*H*)-dione **3p**




<sup>1</sup>H and <sup>13</sup>C spectra of 3-(isoquinolin-1-yl)quinazoline-2,4(1*H*,3*H*)-dione **3q**




<sup>1</sup>H and <sup>13</sup>C spectra of 6-methyl-3-(4-methylpyridin-2-yl)quinazoline-2,4(1*H*,3*H*)-dione **3r**




<sup>1</sup>H and <sup>13</sup>C spectra of 8-methyl-3-(4-methylpyridin-2-yl)quinazoline-2,4(1*H*,3*H*)-dione **3s**




$^1\text{H}$ ,  $^{13}\text{C}$ , and  $^{19}\text{F}$  spectra of 6-fluoro-3-(4-methylpyridin-2-yl)quinazoline-2,4(1*H*,3*H*)-dione **3t**






376 MHz, DMSO-*d*<sub>6</sub>


-119.63




<sup>1</sup>H and <sup>13</sup>C spectra of 6-chloro-3-(4-methylpyridin-2-yl)quinazoline-2,4(1*H*,3*H*)-dione **3u**




<sup>1</sup>H and <sup>13</sup>C spectra of 7-chloro-3-(4-methylpyridin-2-yl)quinazoline-2,4(1*H*,3*H*)-dione **3v**




<sup>1</sup>H and <sup>13</sup>C spectra of 6-bromo-3-(4-methylpyridin-2-yl)quinazoline-2,4(1*H*,3*H*)-dione **3w**




<sup>1</sup>H and <sup>13</sup>C spectra of methyl 3-(4-methylpyridin-2-yl)-2,4-dioxo-1,2,3,4-tetrahydroquinazoline-7-carboxylate **3x**





<sup>1</sup>H and <sup>13</sup>C spectra of 1-methyl-3-(4-methylpyridin-2-yl)quinazoline-2,4(1*H*,3*H*)-dione **3y**





<sup>1</sup>H and <sup>13</sup>C spectra of 6,7-dimethoxy-3-(4-methoxypyridin-2-yl)quinazoline-2,4(1*H*,3*H*)-dione **3z**




<sup>1</sup>H and <sup>13</sup>C spectra of ethyl 2-(3-(5-cyanopyridin-2-yl)ureido)benzoate **4b**



<sup>1</sup>H and <sup>13</sup>C spectra of ethyl 2-(3-(4-methylpyridin-2-yl)ureido)-4,5,6,7-tetrahydrobenzo[b]thiophene-3-carboxylate **7a**




<sup>1</sup>H and <sup>13</sup>C spectra of 3-(4-methylpyridin-2-yl)-5,6,7,8-tetrahydrobenzo[4,5]thieno[2,3-d]pyrimidine-2,4(1H,3H)-dione **8a**




<sup>1</sup>H and <sup>13</sup>C spectra of 3-(pyridin-2-yl)-5,6,7,8-tetrahydrobenzo[4,5]thieno[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione **8b**




<sup>1</sup>H and <sup>13</sup>C spectra of 3-(4-methoxypyridin-2-yl)-5,6,7,8-tetrahydrobenzo[4,5]thieno[2,3-d]pyrimidine-2,4(1*H*,3*H*)-dione **8c**




<sup>1</sup>H and <sup>13</sup>C spectra of 3-(6-phenylpyridin-2-yl)-5,6,7,8-tetrahydrobenzo[4,5]thieno[2,3-d]pyrimidine-2,4(1*H*,3*H*)-dione **8d**



<sup>1</sup>H and <sup>13</sup>C spectra of 3-(quinolin-2-yl)-5,6,7,8-tetrahydrobenzo[4,5]thieno[2,3-d]pyrimidine-2,4(1*H*,3*H*)-dione **8e**



<sup>1</sup>H and <sup>13</sup>C spectra of 3-(4-methylpyridin-2-yl)-1,5,6,7-tetrahydro-2*H*-cyclopenta[4,5]thieno[2,3-*d*]pyrimidine-2,4(3*H*)-dione **8f**



**Table S1 Crystal data and structure refinement for 3a.**

Identification code	CJL-278
CCDC number	2249286
Empirical formula	C <sub>14</sub> H <sub>11</sub> N <sub>3</sub> O <sub>2</sub>
Formula weight	253.26
Temperature/K	100(2)
Crystal system	monoclinic
Space group	P2 <sub>1</sub> /c
a/Å	11.4062(3)
b/Å	14.6246(3)
c/Å	6.9952(2)
α/°	90
β/°	94.055(2)
γ/°	90
Volume/Å <sup>3</sup>	1163.96(5)
Z	4
ρ <sub>calcg/cm<sup>3</sup></sub>	1.445
μ/mm <sup>-1</sup>	0.822
F(000)	528.0
Crystal size/mm <sup>3</sup>	0.24 × 0.22 × 0.16
Radiation	Cu Kα ( $\lambda = 1.54184$ )
2Θ range for data collection/°	7.77 to 159.398
Index ranges	-14 ≤ h ≤ 10, -17 ≤ k ≤ 18, -8 ≤ l ≤ 8
Reflections collected	8636
Independent reflections	2434 [R <sub>int</sub> = 0.0380, R <sub>sigma</sub> = 0.0307]
Data/restraints/parameters	2434/0/173
Goodness-of-fit on F <sup>2</sup>	1.045
Final R indexes [I>=2σ (I)]	R <sub>1</sub> = 0.0432, wR <sub>2</sub> = 0.1137
Final R indexes [all data]	R <sub>1</sub> = 0.0469, wR <sub>2</sub> = 0.1167
Largest diff. peak/hole / e Å <sup>-3</sup>	0.24/-0.32