



Suppression PCR-based selective enrichment sequencing for pathogen and antimicrobial resistance detection on cell-free DNA in sepsis - A targeted, blood culture independent approach for rapid pathogen and resistance diagnostics in septic patients

Supplement



Supplement Figure S1. Identification of suitable HSV I primers. For the identification of most suitable SUPSETS target regions, bioinformatically identified primers were tested. Each lane represents a PCR product with the indicated HSV I primer. For all PCR experiments, the expected fragment length in bp is depicted and inside the gel image displayed with a red arrow. A: 40,000 GE of digested, adapter ligated HSV I DNA were tested in a one-step PCR as described in methods section. B: 40,000 GE digested, adapter ligated HSV I DNA were tested by the complete SUPSETS workflow. C: 40000 GE digested, adapter ligated HSV I DNA were spiked in 6,000 GE human digested, adapter ligated DNA and tested by the complete SUPSETS workflow. D: 4,000 GE digested, adapter ligated HSV I DNA were spiked in 6,000 GE human digested, adapter ligated DNA and tested by the complete SUPSETS workflow. NTC, non-template control; Hu, human only sample; Neg, HSV I DNA with human primer.



Supplement Figure S2. Antibiotic resistance plasmids. Antibiotic resistance plasmids. Resistance cassettes with relevant parts of AMR genes were cloned in the plasmid pUCIDT-AMP by Integrated DNA Technologies GmbH (IDT). A: Resistance plasmid A contains parts of the resistance genes *mecA*, *mecC* and *Oxa-48*. B: Resistance plasmid B contains parts of the resistance genes *ndm-1*, *bla-CTX-M-15*, *bla-KPC-2* and *catL*. C: Resistance plasmid C contains parts of the resistance genes *vanA*, *vanB*, *tetA* and *tetB*. D: Resistance plasmid D contains parts of the resistance genes *AAC(6')-Ib*, *strA-PH(3')-Ib*, *ermB* and *qnrA1*. All resistance cassettes are indicated in orange, while the resistance gene parts are depicted in blue. The selecting TEM116 ampicillin resistance is depicted in yellow. Furthermore, the used restriction enzymes AluI and HpyCH4V are shown. Images created in benchling.com.

Supplement table S1. Primer set used per experiment. Spike-In A: Spike-In *E. faecium*, *E. coli*, and HSV I. Spike-In B: Spike-In *B. fragilis*, *C. albicans*, *K. pneumoniae*, *P. aeruginosa* and *S. aureus*.

	Technical validation			HSV I testing		Clinical samples	
	Spike-In A	Spike-In B	Multiplex	1 Step	SUPSETS	Flongle	Flow Cell
<i>B. fragilis</i>	B. frag T1 & T2	B. frag T1 & T2	B. frag T1 & T2	-	-	B. frag T1 & T2	B. frag T1 & T2
<i>C. albicans</i>	C. alb T1 – T3	C. alb T1 – T3	C. alb T1 – T3	-	-	C. alb T1 & T3	C. alb T1 – T3
<i>E. faecium</i>	E. faec T1 & T2	E. faec T1 & T3	E. faec T1 & T2	-	-	E. faec T1 & T3	E. faec T1 & T2
<i>E. coli</i>	E. coli T1 & T2	-	E. coli T1 & T2	-	-	E. coli T1 & T2	E. coli T1 & T2

HSV I	HSV I T1 & T2	-	HSV I T1 & T2	HSV_01	HSV_01	-	HSV I T1 & T2
				HSV_02	HSV_02		
<i>K. pneumoniae</i>	K. pneum. T1 & T2	K. pneum. T1 & T2	K. pneum. T1 & T2	HSV_03	HSV I T1		
<i>P. aeruginosa</i>	P. aerug T1 & T2	P. aerug T1 & T2	P. aerug T1 & T2	HSV_04	HSV_06		
<i>S. aureus</i>	S. aureus T1 & T2	S. aureus T1 & T2	S. aureus T1 & T2	HSV I T1	HSV_07		
16S	16S	16S	16S	HSV_06	HSV_08		
AMR	-	-	cat1_V2 ermB mecA_V2 mecC qnrA1 strA-APH(3") tetA tetB vanA vanB AAC(6)-Ib TEM116 ndm1 OXA48 blaCTX-M-15_V2 blaKPC-2	-	cat1_V1 ermB mecA_V1 mecC qnrA1 strA-APH(3") tetA tetB vanA vanB AAC(6)-Ib TEM116 ndm1 OXA48 blaCTX-M-15_V1 blaKPC-2	cat1_V2 ermB mecA_V2 mecC qnrA1 tetA tetB vanA vanB AAC(6)-Ib TEM116 ndm1 OXA48 blaCTX-M-15_V2 blaKPC-2	

Supplement table S2. Raw sequencing and mapping data for technical validation. Mapping on full target genome.

Spike-In	Pathogen		Human	
	Mapped reads	portion all reads [%]	Mapped reads	portion all reads [%]
<i>S. aureus</i>	144,613	7.05375591	1,786,390	87.1343449
<i>P. aeruginosa</i>	830,584	25.4745282	2,297,695	70.4717356
<i>K. pneumoniae</i>	613,937	37.2932865	1,031,292	62.6453008
HSV I	18,424	16.3754333	91,532	81.3545463
<i>E. coli</i>	23,888	32.4283232	48,803	66.2508145
<i>E. faecium</i>	120,240	60.1795778	78,440	39.2588663
<i>C. albicans</i>	1,550	0.07287955	1,989,341	93.5369478
<i>B. fragilis</i>	235,297	16.6421593	1,099,192	55.2896548

Supplement table S3. Characteristics of used clinical samples. Timepoint reflects days after ICU admission. BC, blood culture.

Sample	Septic status	Timepoint [days]	Age [years]	Sex	Localization	BC result	Clinical Metagenomics [21]
FL_01	sepsis	14	57	male	Abdominal (peritonitis)	Negative (VRE OP smear)	B. fragilis, E. faecium, E. coli, H. influenzae, S. epidermidis
FL_02	sepsis	2	74	male	Abdominal (peritonitis)	Negative (VRE abdominal smear)	E. faecium
FL_03	sepsis	0	41	male	Abdominal (peritonitis)	E. coli	E. coli
FL_04	sepsis	1	75	male	Abdominal (peritonitis)	negative	B. thetaiotaomicron, B. animalis, E. faecium, E. coli, K. pneumoniae, L. fermentum, V. parvula
FC_01	sepsis	7	78	female	Abdominal (peritonitis)	negative	E. faecium, Human herpesvirus 1
FC_02	sepsis	28	66	male	Abdominal	negative	K. pneumoniae, P. mendocina, S. aureus
FC_03	OP control	1	NA	NA	-	-	-
FC_04	Healthy	-	NA	NA	-	-	-
FC_05	OP control	2	NA	NA	-	-	-
FC_06	sepsis	28	57	male	Abdominal (peritonitis)	Negative (VRE wound swab)	B. fragilis, B. thetaiotaomicron, E. faecium, H. influenza, Human herpesvirus 1

Supplement table S4. Raw sequencing and mapping data for selected clinical samples on MinION Flow Cell. Reads considered with mapq >4, all human reads accumulated, pathogen reads per target amplicons accumulated. FC, Flow Cell.

FC samples	FC_01	FC_02	FC_03	FC_04	FC_05	FC_06
<i>H. sapiens</i>	127,496	9,795	150,899	7,877	68,041	113,028
<i>B. fragilis</i>	0	0	0	0	0	2,262
<i>C. albicans</i>	0	0	0	0	0	0
<i>CARD</i>	0	0	0	0	0	832
<i>E. coli</i>	0	0	0	0	0	662
<i>E. faecium</i>	18	0	0	0	0	1
<i>HSV I</i>	101	0	51	0	0	520
<i>K. pneumoniae</i>	0	0	0	0	0	823
<i>P. aeruginosa</i>	0	0	0	0	0	0
<i>S. aureus</i>	0	0	0	0	0	0

Supplement table S5. Raw sequencing and mapping data for selected clinical samples on MinION Flongle. Reads considered with mapq >4, all human reads accumulated, pathogen reads per target amplicons accumulated. Igκ reads resulted from mapping on full target genome. FL, Flongle.

FL samples	FL_01	FL_02	FL_03	FL_04	FL_04-single
Igκ	6,474	760	8,104	9,64	8,692
<i>H. sapiens</i>	42,572	23,975	98,992	113,712	54,278
<i>B. fragilis</i>	1,603	1	1	1	0

<i>C. albicans</i>	0	0	0	0	0
CARD	1,562	12	1,187	2	54
<i>E. coli</i>	1	0	708	114	219
<i>E. faecium</i>	1,048	51	0	42	90
<i>K. pneumoniae</i>	0	0	0	0	0

Supplement table S6. Approximate duration of SUPSETS workflow. h, hours; min, minutes.

Step	Protocol
cfDNA isolation	2 h
End-prep and adapter ligation	3 h
Template generation	50 min
Purification	20 min
Amplification	1 h 50 min
Library preparation	3 - 4 h
Sequencing	5 min - 24 h
Total	11 - 12 h 5 min - 24 h sequencing

Supplement table S7. Approximate costs and price per sample of SUPSETS. For multiplexing on a MinION Flow Cell, 10 samples per run are calculated. For MinION Flongle experiments, costs for one single sample per analysis are depicted. State of prices: 12/2023.

Reagent	Selling price	Price per sample FC (10 samples)	Price per sample FL (1 sample)
cfDNA isolation kit e.g. EZ1&2 ccfDNA Kit (48)	1000 € / 48 reactions	20.83 €	20.83 €
AmpliTaq Gold 360 Polymerase	1,304 € / 1,000 U	5.22 €	5.22 €
Primer / Adapter	-	~ 0.1 €	~ 0.1 €
AMPure XP Beads	1,200 € / 60 mL	~ 2 - 3 €	~ 2 - 3 €
Ligation Sequencing Kit V14 SQK-LSK114	570 € / 6 reactions	-	95 €
Native Barcoding Kit 24 V14 SQK-NBD114.24	665 € / 6 reactions	11 € (for 6 x 10 samples)	-
MinION Flongle R.10.4.1	797 € / 12 Flongles	-	66.42 €
MinION Flow cell R.10.4.1	855 €	85 € (for 10 samples)	-
NEB Blunt/TA Ligase Master Mix	438 € / 250 reactions	1.75 €	1.75 €
dNTP Mix (10 mM)	100 € / mL	0.2 €	0.2 €
NEB Quick Blunting Kit	378 € / 100 reactions	3.78 €	3.78 €
NEBNext FFPE DNA Repair Mix	642 € / 96 reactions	6.69 €	6.69 €
NEBNext Ultra II End Repair/dA-Tailing Module	886 € / 96 reactions	9.23 €	9.23 €
NEBNext Quick Ligation Reaction Buffer (5x)	47 € / 2 mL	0.24 €	0.24 €
T4 Quick Ligase (400 U/μL)	288 € / 100,000 U	10 - 12 €	10 - 12 €
Total		~ 160 €	~ 225 €
Upfront costs:			

MinION Starter pack (1 Mk1B, 1 Flow cell, 1 sequencing kit, 1 washing kit)	950 €		
Optional: cfDNA isolation device e.g. EZ2 Connect	30,000 €		

Supplement table S8. Reference genomes and corresponding TaxIDs.

Reference	Corresponding ID
H. sapiens	GRCh38
B. fragilis	NZ_CP069563.1
C. albicans	C_albicans_SC5314
E. faecium	NZ_CP039729.1
E. coli	NC_000913
HSV I	NC_001806.2
K. pneumoniae	NC_016845.1
P. aeruginosa	NC_002516.2
S. aureus	NC_007795.1

Supplement table S9. List of primers and target sequence.

Primer name	Target Primer sequence (5' – 3')	Adressed Gen	Adressed genomic region
P. aerug T1	GATCGAGTCGTTCATCGG	PA2338	NC_002516.2:2,581,960-2,582,147
P. aerug T2	CAGGATATCGCTAACCG	PA1813	NC_002516.2:1,971,919-1,972,054
S. aureus T1	GTAGGTACATCATCGTTAAGGC	SAOUHSC_00060	NC_007795.1:64,011-64,883
S. aureus T2	GCGATTATTGTTGCTGAAGG	SAOUHSC_02279	NC_007795.1:2,110,362-2,110,725
C.alb T1	GGATTGCCTTGTCTAACCC	FGR14	Ca21chrR_C_albicans_SC5314:1,129,506-1,131,023
C.alb T2	GGTATTGGGTTGCTGGG	-	Ca21chr1_C_albicans_SC5314:173,093-173,226
C.alb T3	CTGCCTCCATACTTGTAGGC	SFI 1 2.0	Ca21chr7_C_albicans_SC5314:645,263-645,549
K.pneu T1	GTGGTGGAAACTCTACACCG	KPHS_05240	NC_016845.1:562,800-563,175
K.pneu T2	GGCTATGGGATGATATTGCC	KPHS_18590	NC_016845.1:1,929,825-1,930,019
B.frag T1	TCGGAAGGAGAGAACATCC	I6J55_RS14525	NZ_CP069563.1:3,565,788-3,566,174
B.frag T2	TCTATAGATGGCAACACACGG	I6J55_RS00215	NZ_CP069563.1:44,925-45,117
E.faec T1	AAGACCGACTAGCGAAAGC	sufD	NZ_CP039729.1:568,370-568,596
E.faec T2	GCAATTGTTAGAAGGGTGGC	pslx	NZ_CP039729.1:101,372-101,564
E.faec T3	GTCAATCTGGTCTGCCTCA	RecG	NZ_CP039729.1:101,067-101,280
E.coli T1	CAGAACCGCGCTAATT	cynR	NC_000913:358,005-358,231
E.coli T2	CTCGCGCGATTATC	yejB	NC_000913:2,275,142-2,275,233
HSV I T1	GGTCGAAAGCGTACTTGGC	UL12	NC_001806.2:25,762-25,934
HSV I T2	GCAACTCGTAGCAAATAGGC	UL16	NC_001806.2:30,224-30,342
HSV_01	CTGTTGGGTTAGAATTCCG	UL41	NC_001806.2:91,653-91,672
HSV_02	AACGTCTCCATCCATGCC	US8	NC_001806.2:141,973-141,990
HSV_03	AGTGATGAAATGGTGTCCG	UL4/5	NC_001806.2:12,022-12,041
HSV_04	GTGGTTCATGTGTGTGCC	-	NC_001806.2:94,673-94,690
HSV_06	AAAGCAGTAACCAGGTCCG	-	NC_001806.2:91,065-91,083
HSV_07	AAACCTAAGGACAGCGGC	UL15/17	NC_001806.2:31,426-31,443

HSV_08	CGATGTTAATGGTGAACGC	UL15/17	NC_001806.2:31,479-31,497
HSV_10	ACCACCAGGTGTTAACCC	UL44	NC_001806.2:97,225-97,243
16S	CCATGAAGTCGGAATCGC	16S	-
Igκ	AGAGCAGGACAGCAAGGAC	Igκ constant chain	NC_000002.12:88,857,361-88,857,683
cat1_V1	AAGCACAAGTTTATCCGGC	cat1	gb KC243783.1 + 1-2007 mecA
cat1_V2	AAGATGTGGCGTGTACGG	cat1	gb KC243783.1 + 1-2007 mecA
blaCTX-M-15_V1	TTGTTAGGAAGTGTGCCGC	blaCTX-M-15	gb AY044436.1 + 1436-2311 CTX-M-15
blaCTX-M-15_V2	GGAGACGAAACGTTCCG	blaCTX-M-15	gb AY044436.1 + 1436-2311 CTX-M-15
ndm1	ACATCGCTTGTTGGTGGC	ndm1	gb FN396876.1 - 2407-3219 NDM-1
TEM116	TGAGGCACCTATCTCAGCG	TEM116	gb U36911.1 + 1430-2290 TEM-116
mecA_V1	ACTACGGTAACATTGATCGC	mecA	gb KC243783.1 + 1-2007 mecA
mecA_V2	AATCGATGGTAAAGGTTGGC	mecA	gb KC243783.1 + 1-2007 mecA
mecC	CATCACCAAGGTTCAACCC	mecC	gb FR821779.1 - 35681-37678 mecC
strA-APH(3")	GCATTCTGACTGGTTGCC	strA-APH(3")	gb AF313472.2 + 15594-16397 APH(3")-Ib
ermB [46]	GAACATCTGTGGTATGGCG	ermB	gb AF242872.1 + 2132-2878 ErmB
OXA48	GCACGTATGAGCAAGATGC	OXA48	gb AY236073.2 + 2188-2985 OXA-48
blaKPC-2	TCTGACAACAGGCATGACG	blaKPC-2	gb AY034847.1 + 6-887 KPC-2
vanA	CCAATACCGCACAAACCGA	vanA	gb M97297.1 + 6980-8010 vanA
vanB	GAAGTGGATCAAATCCGGC	vanB	gb KF823969.1 + 5111-6139 vanB
qnr1	CAATGCCAATCTCAGCGG	qnr1	gb DQ831141.1 + 8922-9578 QnrA1
tetA	AGGCAAGCAGGATGTAGC	tetA	gb AF534183.1 + 2971-4245 tet(A)
tetB	CAAAGGCTTGBAATACTGAGTG	tetB	gb AB089595.1 + 1-1206 tet(B)
AAC(6)-Ib	TCATATCGTCGAGTGGTGG	AAC(6)-Ib	gb JQ808129.1 + 634-1188 AAC(6')-Ib