

International Journal of Molecular Sciences
Supplementary Materials

**Bioreduction of 4'-hydroxychalcone in deep eutectic solvents:
optimization and efficacy with various yeast strains**

Paweł Chlipała, Tomasz Janeczko, Marcelina Mazur*

Department of Food Chemistry and Biocatalysis, Wrocław University of Environmental and Life Sciences, Norwida 25, 50-375 Wrocław, Poland




Figure S1: ^1H -NMR spectrum of 4'-hydroxychalcone (*trans*-1).




Figure S2: ^{13}C -NMR spectrum of 4'-hydroxychalcone (*trans*-**1**).




Figure S3: COSY spectrum of 4'-hydroxychalcone (*trans*-**1**).




Figure S4: HSQC spectrum of 4'-hydroxychalcone (*trans*-1).




Figure S5: HMBC spectrum of 4'-hydroxychalcone (*trans*-1).




Figure S6: ^1H -NMR spectrum of 4'-hydroxychalcone (*cis*-1).




Figure S7: ^{13}C -NMR spectrum of 4'-hydroxychalcone (*cis*-1).




Figure S9: HSQC spectrum of 4'-hydroxychalcone (cis-1).




Figure S10: HMBC spectrum of 4'-hydroxychalcone (*cis*-1).



Figure S11: ^1H -NMR spectrum of 4'-hydroxy-dihydrochalcone (**2**).




Figure S12: ^{13}C -NMR spectrum of 4'-hydroxy-dihydrochalcone (**2**).




Figure S13: COSY spectrum of 4'-hydroxy-dihydrochalcone (**2**).




Figure S14: HSQC spectrum of 4'-hydroxy-dihydrochalcone (**2**).




Figure S15: HMBC spectrum of 4'-hydroxy-dihydrochalcone (**2**).