

## Short Communication: Novel Di- and Triselenoesters as Effective Therapeutic Agents Inhibiting Multidrug Resistance Proteins in Breast Cancer Cells

Dominika Radomska <sup>1\*</sup>, Robert Czarnomysy <sup>1\*</sup>, Krzysztof Marciniak <sup>2</sup>, Justyna Nowakowska <sup>2</sup>, Enrique Domínguez-Álvarez <sup>3</sup> and Krzysztof Bielawski <sup>1</sup>


<sup>1</sup> Department of Synthesis and Technology of Drugs, Medical University of Białystok, Kilinskiego 1, 15-089 Białystok, Poland; dominika.radomska@umb.edu.pl (D.R.); robert.czarnomysy@umb.edu.pl (R.C.)

<sup>2</sup> Department of Organic Chemistry, Medical University of Silesia, Jagiellonska 4, 41-200 Sosnowiec, Poland


<sup>3</sup> Instituto de Química Orgánica General (IQOG-CSIC), Consejo Superior de Investigaciones Científicas, Juan de la Cierva 3, 28006 Madrid, Spain

\* Correspondence: dominika.radomska@umb.edu.pl (D.R.); robert.czarnomysy@umb.edu.pl (R.C.); Tel.: +48-85-748-57-00; Fax: +48-85-879-57-00


This document contains representative histograms from the flow cytometric analysis, which show the level of measured autofluorescence of the culture medium and the tested cancer cells – their mean fluorescence intensity (MFI).



**Figure S1.** The original histogram from the cytometric analysis that was utilized to assess the level of autofluorescence in the culture medium used in the study (Dulbecco's Minimal Eagle Medium; DMEM). The mean values with standard deviation were reported based on data from three separate experiments ( $n = 3$ ) conducted in triplicate.



**Figure S2.** The original histogram from the cytometric analysis that was utilized to assess the level of autofluorescence in MCF-7 breast cancer cells. The mean values with standard deviation were reported based on data from three separate experiments (n = 3) conducted in triplicate.



**Figure S3.** The original histogram from the cytometric analysis that was utilized to assess the level of autofluorescence in MDA-MB-231 breast cancer cells. The mean values with standard deviation were reported based on data from three separate experiments (n = 3) conducted in triplicate.