

Rhodamine B Pentyl Ester and its Methyl, Ethyl, Propyl, and Butyl Homologues

Justin A. Ross,^{*,A} Benjamin P. Ross,^{B,C} Kelly L. Cosgrove,^B Halina Rubinsztein-Dunlop,^A and Ross P. McGearry^{B,C}

^ACentre for Biophotonics and Laser Science, School of Physical Sciences;

^BSchool of Molecular and Microbial Sciences;

^CSchool of Pharmacy;


The University of Queensland, Brisbane, Queensland 4072, Australia.

Tel.: +61 7 3366 9583; Fax: +61 7 3365 1242; E-mail: justinr@physics.uq.edu.au

*Author to whom correspondence should be addressed

Received: 22 June 2006 / Accepted: 11 August 2006 / Published: 1 December 2006

Keywords: rhodamine, esterification, acetyl chloride, lipophilicity.



The rhodamines are a highly fluorescent class of compound used in many different fields of research, from the lasing medium in dye lasers to biological stains and markers for cellular drug resistance [1-3]. We are interested in the influence of lipophilicity on the transport of esters of rhodamine in multidrug resistance transporter assays. Herein we describe the synthesis of the novel compound rhodamine B pentyl ester (**2**), and deposit the spectral data of the known methyl (**3**), ethyl (**4**), propyl (**5**), and butyl (**6**) homologues [4]. Ester **2** was prepared by a slightly modified facile method [5] that utilizes anhydrous hydrogen chloride generated *in situ* by the addition of AcCl to a mixture of the rhodamine B free acid (**1**) and *n*-pentanol.

Rhodamine B pentyl ester (**2**)

N-[6-(Diethylamino)-9-[2-(pentylloxycarbonyl)phenyl]-3*H*-xanthen-3-ylidene]-*N*-ethylethanaminium chloride

Acetyl chloride (1.25 mL, 17.6 mmol) was added dropwise to a stirred mixture of rhodamine B (**1**, 100 mg, 0.21 mmol) and *n*-pentanol (25 mL), and the reaction was heated at 50°C under an atmosphere of argon. After 2 days the reaction was ~ 90% complete by TLC, and additional acetyl chloride (250 µL, 3.52 mmol) was added to drive the reaction to completion. After a further 24 h TLC indicated no remaining rhodamine B (**1**) and the solution was evaporated *in vacuo* (water bath temperature < 50 °C to avoid degradation of the product) to afford rhodamine B pentyl ester (**2**), which was contaminated with a small amount of *n*-pentanol. The molar ratio of rhodamine B pentyl ester (**2**):*n*-pentanol was ~ 3:1 by ¹H NMR. To separate the ester (**2**) from residual alcohol, the crude product was dissolved in water (100 mL) and washed with ethyl acetate (3 × 50 mL). Acetonitrile (35 mL) was added to the aqueous layer and this solution was lyophilised to afford the title compound (**2**), a hygroscopic amorphous solid (85%).

TLC R_f 0.41 (silica gel 60 F₂₅₄ aluminum sheets, *n*-butanol:water:ethanol 9:2:1).

ES-MS, m/z : 513 [M+H]⁺.

¹H NMR (500 MHz, d₆-DMSO) δ 8.24 (1H, dd, $J = 7.9$ and 1.2 Hz), 7.91 (1H, dt, $J = 7.5$ and 1.3 Hz), 7.85 (1H, dt, $J = 7.7$ and 1.3 Hz), 7.51 (1H, dd, $J = 7.5$ and 1.1 Hz), 7.11 (2H, dd, $J = 9.6$ and 2.4 Hz), 7.02-6.99 (4H, m), 3.89 (2H, t, $J = 6.2$ Hz), 3.65 (8H, br q, $J = 7.1$ Hz), 1.20 (12H, br t, $J = 7.0$ Hz) overlapping with ~1.18-1.13 (2H, m), 1.06 (2H, sextet, $J = 7.4$ Hz), 0.88-0.82 (2H, m), 0.71 (3H, t, $J = 7.4$ Hz).

¹³C NMR (100 MHz [6], d₆-DMSO) δ 165.1, 157.6, 157.1, 155.1, 133.0, 132.7, 131.0, 130.9, 130.5, 130.4, 129.9, 114.6, 112.9, 95.9, 65.2, 45.3, 27.5, 27.4, 21.7, 13.6, 12.4.

UV-visible (H₂O): λ_{max} 559 nm.

HRMS calcd for [M+H]⁺ 513.3117, found 513.3122.

The methyl (**3**), ethyl (**4**), propyl (**5**) and butyl (**6**) homologues of **2** were synthesized by the same method except that evaporation of the solution in vacuo afforded the ester (**3-6**) pure, without the need for partitioning between water and ethyl acetate. Previously these compounds were isolated as the bromide, iodide or perchlorate salts, and their spectra were recorded in CDCl₃ [4]. Herein we deposit the spectra of the chloride salts (**3-6**) recorded in d₆-DMSO.

Rhodamine B methyl ester (**3**)

***N*-[6-(Diethylamino)-9-[2-(methoxycarbonyl)phenyl]-3*H*-xanthen-3-ylidene]-*N*-ethylethanaminium chloride**

TLC R_f 0.35 (silica gel 60 F₂₅₄ aluminum sheets, *n*-butanol:water:ethanol 9:2:1).

ES-MS, m/z : 457 [M+H]⁺.

¹H NMR (500 MHz, d₆-DMSO) δ 8.27 (1H, dd, $J = 7.9$ and 1.2 Hz), 7.92 (1H, dt, $J = 7.6$ and 1.3 Hz), 7.84 (1H, dt, $J = 7.7$ and 1.3 Hz), 7.51 (1H, dd, $J = 7.5$ and 1.1 Hz), 7.08 (2H, dd, $J = 9.6$ and 2.5 Hz), 6.99-6.97 (4H, m), 3.65 (8H, br q, $J = 7.02$ Hz), 3.60 (3H, s), 1.21 (12H, br t, $J = 7.0$ Hz).

¹³C NMR (75 MHz, d₆-DMSO) δ 165.0, 157.9, 157.1, 155.1, 133.4, 133.3, 130.8, 130.8, 130.5, 130.5, 129.2, 114.6, 112.8, 95.9, 52.5, 45.3, 12.5.

UV-visible (H₂O): λ_{max} 558 nm.

HRMS calcd for [M+H]⁺ 457.2491, found 457.2481.

Rhodamine B ethyl ester (**4**)

***N*-[6-(Diethylamino)-9-[2-(ethoxycarbonyl)phenyl]-3*H*-xanthen-3-ylidene]-*N*-ethylethanaminium chloride**

TLC R_f 0.37 (silica gel 60 F₂₅₄ aluminum sheets, *n*-butanol:water:ethanol 9:2:1).

ES-MS, m/z : 471 [M+H]⁺.

¹H NMR (500 MHz, d₆-DMSO) δ 8.25 (1H, dd, $J = 7.9$ and 1.1 Hz), 7.91 (1H, dt, $J = 7.6$ and 1.4 Hz),

7.84 (1H, dt, $J = 7.7$ and 1.2 Hz), 7.52 (1H, dd, $J = 7.6$ and 1.1 Hz), 7.09 (2H, dd, $J = 9.6$ and 2.5 Hz), 7.01-6.99 (4H, m), 3.98 (2H, q, $J = 7.1$ Hz), 3.65 (8H, br q, $J = 7.0$ Hz), 1.21 (12H, br t, $J = 7.0$ Hz), 0.90 (3H, t, $J = 7.1$ Hz).

^{13}C NMR (75 MHz, $\text{d}_6\text{-DMSO}$) δ 164.7, 157.7, 157.1, 155.1, 133.1, 133.0, 130.9, 130.7, 130.5, 130.4, 129.7, 114.6, 112.9, 95.9, 61.0, 45.4, 13.4, 12.4.

UV-visible (H_2O): λ_{max} 559 nm.

HRMS calcd for $[\text{M}+\text{H}]^+$ 471.2648, found 471.2643.

Rhodamine B propyl ester (5)

***N*-[6-(Diethylamino)-9-[2-(propoxycarbonyl)phenyl]-3*H*-xanthen-3-ylidene]-*N*-ethylethanaminium chloride**

TLC R_f 0.39 (silica gel 60 F₂₅₄ aluminum sheets, *n*-butanol:water:ethanol 9:2:1).

ES-MS, m/z : 485 $[\text{M}+\text{H}]^+$.

^1H NMR (500 MHz, $\text{d}_6\text{-DMSO}$) δ 8.25 (1H, dd, $J = 7.8$ and 1.2 Hz), 7.91 (1H, dt, $J = 7.6$ and 1.3 Hz), 7.85 (1H, dt, $J = 7.7$ and 1.2 Hz), 7.52 (1H, dd, $J = 7.5$ and 1.0 Hz), 7.09 (2H, dd, $J = 9.6$ and 2.5 Hz), 7.02-6.99 (4H, m), 3.89 (2H, t, $J = 6.4$ Hz), 3.65 (8H, br q, $J = 7.0$ Hz), 1.29 (2H, sextet, $J = 6.9$ Hz), 1.20 (12H, br t, $J = 7.0$ Hz), 0.62 (3H, t, $J = 7.4$ Hz).

^{13}C NMR (75 MHz, $\text{d}_6\text{-DMSO}$) δ 164.9, 157.7, 157.1, 155.1, 133.1, 132.9, 130.9, 130.8, 130.5, 130.4, 129.8, 114.6, 112.9, 95.9, 66.7, 45.4, 21.1, 12.4, 10.0.

UV-visible (H_2O): λ_{max} 559 nm.

HRMS calcd for $[\text{M}+\text{H}]^+$ 485.2804, found 485.2801.

Rhodamine B butyl ester (6)

***N*-[6-(Diethylamino)-9-[2-(butoxycarbonyl)phenyl]-3*H*-xanthen-3-ylidene]-*N*-ethylethanaminium chloride**

TLC R_f 0.40 (silica gel 60 F₂₅₄ aluminum sheets, *n*-butanol:water:ethanol 9:2:1).

ES-MS, m/z : 499 $[\text{M}+\text{H}]^+$.

^1H NMR (500 MHz, $\text{d}_6\text{-DMSO}$) δ 8.24 (1H, dd, $J = 7.9$ and 1.2 Hz), 7.91 (1H, dt, $J = 7.5$ and 1.4 Hz), 7.84 (1H, dt, $J = 7.7$ and 1.4 Hz), 7.51 (1H, dd, $J = 7.6$ and 1.1 Hz), 7.10 (2H, dd, $J = 9.6$ and 2.5 Hz), 7.02-7.00 (4H, m), 3.91 (2H, t, $J = 6.2$ Hz), 3.65 (8H, br q, $J = 7.2$ Hz), 1.21 (12H, br t, $J = 7.0$ Hz) overlapping with \sim 1.20-1.15 (2H, m), 0.94 (2H, sextet, $J = 7.6$ Hz), 0.68 (3H, t, $J = 7.4$ Hz).

^{13}C NMR (75 MHz, $\text{d}_6\text{-DMSO}$) δ 165.0, 157.5, 157.1, 155.1, 133.0, 132.7, 130.9, 130.7, 130.4, 130.3, 129.8, 114.6, 112.8, 95.8, 64.9, 45.3, 29.8, 18.4, 13.4, 12.4.

UV-visible (H_2O): λ_{max} 559 nm.

HRMS calcd for $[\text{M}+\text{H}]^+$ 499.2961, found 499.2973.

Acknowledgments

We thank Mr. Graham MacFarlane (School of Molecular and Microbial Sciences, The University of Queensland) for accurate mass measurements. This work was supported by funds from the Centre for Biophotonics and Laser Science, School of Physical Sciences, The University of Queensland.

References

1. Zollinger, H. *Color Chemistry : Syntheses, Properties, and Applications of Organic Dyes and Pigments*, 2nd rev. ed.; VCH: Weinheim, 1991.
2. Gordon, P. F.; Gregory, P. *Organic Chemistry in Colour*; Springer-Verlag: New York, 1983.
3. Abrahart, E. N.; *Dyes and Their Intermediates*, 2nd ed.; Edward Arnold: London, 1977.
4. Ramos, S. S.; Vilhena, A. F.; Santos, L.; Almeida, P. *Magn. Reson. Chem.* **2000**, *38*, 475-478.
5. Ross, J. A.; Ross, B. P.; Rubinsztein-Dunlop, H.; McGearry, R. P. *Synth. Commun.* **2006**, *36*, 1745-1750.
6. The ^{13}C NMR spectra of **3-6** were run at 75 MHz whereas the ^{13}C NMR spectrum of **2** was recorded at 100 MHz to enable resolution of the resonances which appeared at 27.46 and 27.43 ppm in the 100 MHz spectrum.

© 2006 [MDPI](http://www.mdpi.org). All rights reserved.