

Supplementary material

Methyl 2-((3-(3-methoxyphenyl)-4-oxo-3,4-dihydroquinazolin-2-yl)thio)acetate

Maja Molnar¹, Mario Komar¹, Igor Jerković^{2*}

¹ Josip Juraj Strossmayer University of Osijek, Faculty of Food Technology Osijek, Croatia

² Faculty of Chemistry and Technology, University of Split, R. Boškovića 35, 21000 Split, Croatia

* Correspondence: igor@ktf-split.hr

Figure S1. ¹H NMR spectra of 2-((3-(3-methoxyphenyl)-4-oxo-3,4-dihydroquinazolin-2-yl)thio)acetate.

Figure S2. Mass spectra of 2-((3-(3-methoxyphenyl)-4-oxo-3,4-dihydroquinazolin-2-yl)thio)acetate.

Figure S3. ¹³C NMR spectra of 2-((3-(3-methoxyphenyl)-4-oxo-3,4-dihydroquinazolin-2-yl)thio)acetate; Molfile of ¹H NMR spectra of 2-((3-(3-methoxyphenyl)-4-oxo-3,4-dihydroquinazolin-2-yl)thio)acetate.

Acquisition Time (sec)	2.6542	Comment	MK88 17.12.2021	Date	20 Dec 2021 05:20:16		
Date Stamp	20 Dec 2021 05:20:16	File Name	F:\nmr\prosinac_2021\molnar_mk881712\1\fid	Frequency (MHz)	300.13		
Nucleus	1H	Number of Transients	32	Origin	spect	Owner	nmr
Points Count	16384	Pulse Sequence	zq30	Receiver Gain	456.10	SW(cyclcal) (Hz)	6172.84
Spectrum Offset (Hz)	2701.1702	Spectrum Type	STANDARD	Sweep Width (Hz)	6172.46	Temperature (degree C)	25.000





