

## *Supplementary Materials*

### **(S)-1-Methyl-2-oxoimidazolidine-4-carboxylic acid**

Ashley L. Dey, Majid Motevalli, Isaac Abrahams and Peter B. Wyatt\*

Department of Chemistry, Joseph Priestley Building, Queen Mary University of London, Mile End Road, London UK E1 4NS


\*Correspondence: p.b.wyatt@qmul.ac.uk

#### **Contents**


Figure S1:  $^1\text{H}$  NMR spectrum of (S)-1-methyl-2-oxoimidazolidine-4-carboxylic acid **1** (400 MHz,  $\text{D}_2\text{O}$ )

Figure S2:  $^{13}\text{C}$  NMR spectrum of (S)-1-methyl-2-oxoimidazolidine-4-carboxylic acid **1** (100.6 MHz,  $\text{D}_2\text{O}$ )


Figure S3: ESI mass spectrum of (S)-1-methyl-2-oxoimidazolidine-4-carboxylic acid **1**



**Figure S1:**  $^1\text{H}$  NMR spectrum of (S)-1-methyl-2-oxoimidazolidine-4-carboxylic acid **1** (400 MHz,  $\text{D}_2\text{O}$ )



**Figure S2:**  $^{13}\text{C}$  NMR spectrum of (S)-1-methyl-2-oxoimidazolidine-4-carboxylic acid **1** (100.6 MHz,  $\text{D}_2\text{O}$ )



**Figure S3:** ESI mass spectrum of (S)-1-methyl-2-oxoimidazolidine-4-carboxylic acid **1**