

Article

# Two New Marine Free-Living Nematodes from Jeju Island Together with a Review of the Genus *Gammanema* Cobb 1920 (Nematoda, Chromadorida, Selachinematidae)

Alexei Tchesunov <sup>1,\*</sup>, Raehyuk Jeong <sup>2</sup> and Wonchoel Lee <sup>2</sup> <sup>1</sup> Department of Invertebrate Zoology, Faculty of Biology, Moscow State University, 119991 Moscow, Russia<sup>2</sup> Department of Life Science, Hanyang University, Seoul 04763, Korea; rhjeong88@gmail.com (R.J.); wlee@hanyang.ac.kr (W.L.)

\* Correspondence: AVTchesunov@yandex.ru

<http://zoobank.org/urn:lsid:zoobank.org:pub:4C6B07ED-52DB-42A1-80B7-2881F0559E56>

Received: 19 November 2019; Accepted: 18 December 2019; Published: 1 January 2020

**Abstract:** In the context of exploration of meiofauna in a sandy intertidal zone of Jeju Island (South Korea), over 70 nematode species are identified, some which have been proven to be new for science. Two new free-living marine nematode species of the family Selachinematidae (Chromadorida, Selachinematidae, Choniolaiminae) are described from the intertidal sandy sediments of Jeju Island (South Korea). *Gammanema okhlopkovi* sp. n. is closest to *Gammanema anthostoma* (Okhlopkov, 2002) and differs by having longer cephalic setae (8.5–19  $\mu\text{m}$  in *G. okhlopkovi* versus 6–7.5  $\mu\text{m}$  in *G. anthostoma*) and by the presence of precloacal supplementary organs. The genus diagnosis of *Gammanema* is updated. The genus includes fourteen valid species, while three species are considered species inquirendae due to incomplete diagnoses and illustrations impeding their correct recognition. An annotated list of valid and invalid *Gammanema* species is provided. A pictorial key for valid *Gammanema* species is constructed, which consists of two components: (1) simplified images of heads, and (2) a table summarizing most of the significant measured and numeric characters between species. *Latronema obscuramphis* sp. n. differs from its related species *Latronema aberrans* (Allgén 1934), *Latronema annulatum* (Gerlach, 1953), and *Latronema spinosum* (Andrássy, 1973) by body size, number of supplementary organs, tail shape, length of spicules, and cuticle ornamentation.

**Keywords:** free-living marine nematodes; pictorial key; taxonomy

## 1. Introduction


Data on Jeju marine free-living nematodes have been presented by Rho and Kim [1,2], Rho et al. [3], and Lim and Chang [4]. However, the exploration of the rich nematofauna of Jeju is still in the early stages. In connection with our project on Jeju intertidal meiofauna, we have recorded almost 70 nematode species at four points along a transect through the sandy intertidal zone at Shinyang, in the eastern corner of Jeju Island. A taxonomic survey of Jeju nematodes started with the family Thoracostomopsidae by Jeong et al. [5,6]. Now, we present a communication on the family Selachinematidae.

The selachinematid nematodes are usually characterized by a stout cylindrical body with a truncate anterior end. They have a complicated buccal armament of rhabdions, denticles, or heavy mandibles, enabling them to capture nematode prey. The selachinematids have been known for a long time as carnivores in the strict sense, ingesting other smaller nematodes [7,8]. The percentage of selachinematids and other predator taxa in nematode assemblages are usually low in mud and silty

sediment, but more abundant in sandy sediment with higher species diversity [9,10]. Particularly, most species of *Gammanema* and *Latronema* are confined to sandy sediments of upper shelf zones.

## 2. Materials and Methods

Quantitative samples were taken in June 2018, in the intertidal zone at Shinyang, close to the eastern corner of Jeju Island (Figure 1). Sediments were fixed in 5% neutralized formalin solution and brought back to the laboratory. Meiofauna were extracted using the Ludox method [11], and postfixed with 70% ethanol dyed with Rose bengal. Nematodes were counted and specimens were packed into a Syracuse glass filled with mixture of glycerin, 95% ethanol, and distilled water (1:29:70). The glass was placed in a drying oven set to 40 °C for 1–2 days until completely dehydrated, as in the glycerin-ethanol method [12]. Specimens were mounted in a drop of anhydrous glycerin within a wax-paraffin ring. Specimens were identified, measured, photographed, and drawn under a Leica DM5000 light microscope equipped with Leica Application Suite Version 3.8.0 software and a Leica DFC 425 C digital camera. All the measured sizes are given in  $\mu\text{m}$ . For scanning electron microscopy, specimens were dehydrated in a series: 40% ethanol, 70% ethanol, 95% ethanol, 95% ethanol + acetone (50:50), acetone I, and acetone II, and then critical point dried. Once dried, specimens were mounted on a stub to be coated with platinum–palladium alloy and examined with Cam Scan S-2.



**Figure 1.** Map of sampling locality. This map is made with QGIS software v.2.18.14, a free and open-source geographic information system (<https://qgis.org>).

Type specimens were deposited in National Institute of Biological Resources (South Korea).

## 3. Results—Systematics

### 3.1. Review of Genus *Gammanema*

Order Chromadorida Chitwood, 1933

Family Selachinematidae Cobb, 1915

Subfamily Choniolaiminae Schuurmans-Stekhoven and Adam, 1931

Genus *Gammanema* Cobb, 1920

## 3.1.1. Diagnosis

Updated after Leduc 2013 and Tchesunov 2014 [13,14].

Selachinematidae, Choniolaiminae. Cuticle with homogenous annulations, without longitudinal ridges or lateral differentiation. Six inner and six outer labial sensilla, either setose or papillose; four cephalic setae, often slender and longer; outer labial and cephalic sensilla combined in common circle of 10, with dorso- and ventrosublateral sensilla arranged in four pairs with cephalic sensilla. Amphideal fovea spiral or loop-shaped, usually noticeably larger in males than in females. Somatic setae in irregular longitudinal rows; anterior cervical setae may be as long as cephalic setae. Mouth opening surrounded by partly fused lips shaping a circumoral membrane with fine longitudinal striation. There are twelve projections, from small and inconspicuous to prominent and elaborate, at the rim of the mouth opening. Buccal cavity (pharyngostome) consists of two chambers, anterior cup-shaped and posterior cylindrical; walls of each chamber are strengthened with three cuticularized rhabdions; the rhabdions of the anterior chamber terminate posteriorly in minute denticles. Pharynx cylindroids, evenly muscular and devoid of a terminal bulb. Alimentary tract terminates by rectum and anus. Preloacal midventral supplementary organs sucker-like, cup-shaped, tubular, or absent. Tail short, conical, cuticle of its terminal cone levigated or smooth.

Type species: *Gammanema ferox* (Cobb, 1920).

Annotated species list (valid species **in bold**)

1. ***Gammanema agglutinans* Leduc, 2013**. Leduc 2013:21–25, Figures 11–13, Table 2 [13] (males, females); SW Pacific, region to the east of New Zealand, Chatham Rise, depths 350–1238 m.
2. ***Gammanema anthostoma* Okhlopkov, 2002**. Okhlopkov 2002:42–45, Figures 2 and 3 [15] (males and females); White Sea (Northern Russia), Kandalaksha Bay, depth 14 m.
3. ***Gammanema cancellatum* Gerlach, 1955**. Gerlach 1955:271–272, Abbildung. 6 [16] (female); San Salvador, coastal groundwater. **Remark:** the species is retained here as valid, despite the absence of males in diagnosis, since it can be distinguished sound from other congeners by peculiar cuticle ornamentation with rods.
4. ***Gammanema conicauda* Gerlach, 1953**. Gerlach 1953:553–555, Abbildungen 17 a-f [17] (males, females, juvenile); Tyrrhenian Sea, area of Naples, surf zone; North Sea coast.
5. ***Gammanema curvata* Gagarin et Klerman, 2007**. Gagarin and Klerman 2007:780, Figure 2, Table 2 [18] (male, females); Mediterranean coast of Israel, depth 50 m, sandy sediment.
6. ***Gammanema fennicum* (Gerlach, 1953) Gerlach, 1964**. Gerlach 1953:22–23, Abbildung 6 [19] (as *Halichoanolaimus fennicus*) (female, juvenile); east Baltic Sea. Gerlach 1964:37 [20] (as junior synonym of *Gammanema rapax*). Okhlopkov 2002:45–46, Figures 4 and 5 [15] (males, females, juvenile); White Sea (Northern Russia), Kandalaksha Bay, depth 16 m.
7. *Gammanema ferox* Cobb, 1920 **sp. inq.** Cobb 1920:291–293, Figure 74 [21] (male, female); New Hebrides, coral sand. Though the species is described for both genders, the description lacks some necessary measurements, such as anterior setae and spicules; amphids are not depicted and not mentioned in the text (amphideal fovea possibly inconspicuous or reduced); the only illustration presents a head without an indication of gender and position in slide (possibly sublateral). Restitution of valid state is possible after redescription of new specimens from the type area; the species can be recognized through the prominent anterior projections of rugae.
8. ***Gammanema kosswigi* Gerlach, 1964**. Gerlach 1964:38–39, Abbildungen 11 a–d [19] (male, juvenile); Maldive Islands, coastal groundwater, intertidal zone.
9. ***Gammanema magnum* Shi et Xu, 2018**. Shi and Xu 2018, 3–9, Figures 1–4, Table 1 [22] (males, females); East China Sea, intertidal sandy sediment.
10. ***Gammanema mediterraneum* Vitiello, 1970**. Vitiello 1970:491–493, planche XV, 30 a–e [23] (as *Gammanema mediterranea*) (females, juveniles); West Mediterranean, depths 310–650 m.

11. *Gammanema menzelii* (Ditlevsen, 1918) Gerlach, 1964 **sp. inq.** Ditlevsen 1918:172–173, Plate VI, Figure 2; Plate VII, Figures 1 and 8 [24] (as *Halichoanolaimus Menzelii*) (one male); Danish Belt Sea, depth 30 m, clean sand. Gerlach 1964:31 [19] (transfer to *Gammanema*).
12. *Gammanema okhlopkovi* Tchesunov, Jeong et Lee **sp. n.** Present paper.
13. *Gammanema polydonta* Murphy, 1965. Murphy 1965:176–179, Figures 3A–C and 4A–H [25] (males, females); Chilean coast, depth 45 m.
14. *Gammanema rapax* (Ssaweljev, 1912) Gerlach 1964 **sp. inq.** Ssaweljev 1912:122 [26] (as *Halichoanolaimus rapax*) (male, female); Barents Sea, Kolafjord, Palafjord, Mogilnojesee (lake with an intermediate layer of sea water on the Kildin Island in the Barents Sea), sand. Murphy (1965: 179 [25]) considered the species; dubious because the original description is not illustrated and type specimens are evidently lacking. Gerlach (1964 [20]) synonymized *Halichoanolaimus menzeli* Ditlevsen 1918, *Gammanema ferox* Cobb, 1920 and *Gammanema fennicum* (Gerlach, 1953) with *Gammanema rapax*.
15. *Gammanema smithi* Murphy, 1964. Murphy 1964: 194–198, Figure 3A,B,D,E, Figure 4, Figure 5B [27] (male); Puget Sound (NE Pacific), intertidal sand. **Remark:** there are some discrepancies in the original diagnosis. Caudal glands are described and depicted ([27], Figure 4) as incaudal (i.e., located entirely within tail), while the glands in Figure 5A are definitely excaudal (i.e., gland cell bodies protrude anteriorly and lie along the midgut). Supplementary organs are depicted from a medio-ventral view in Figure 3C, while the hind body in Figure 4 is shown laterally. Taking into account that the diagnosis is based on only one male (holotype), we consider Figures 3C and 5A as wrong and not pertaining to *G. smithi*.
16. *Gammanema tchesunovi* Gagarin et Klerman, 2007. Gagarin and Klerman 2007:782–785, Figure 3, Table 3 [18] (males, females); Mediterranean coast of Israel, depths 50–55 m, sand.
17. *Gammanema uniformis* (Cobb, 1920) Tchesunov et Okhlopkov, 2006. Cobb 1920:293–294, Figure 75 [21] (as *Trogolaimus uniformis*) (seemingly, one male); Atlantic coast of the United States, New Hampshire. Tchesunov and Okhlopkov 2006:40–43, Figure 12, Table 6 [28] (males, females); Atlantic coast of the United States, Maine.

### 3.1.2. Remarks on the Species List and Synonymy


Murphy (1965:179 [25]) considered the species *Gammanema rapax* (Ssaweljev, 1912) as dubious because the original description is not depicted, and type specimens are evidently lacking. Gerlach (1964 [20]) synonymized *Gammanema menzelii* (Ditlevsen, 1918), *Gammanema ferox* Cobb, 1920, and *Gammanema fennicum* (Gerlach, 1953) with *G. rapax*. We find this broad synonymization questionable. *G. rapax* is a large species, measuring 5800–6100  $\mu\text{m}$  in body length (Ssaweljev 1912 [26]), while all other species and specimens are at least half the size (except *G. menzelii*) and found far from the type locality. *G. menzelii* corresponds to *G. rapax* in body length (male 5900  $\mu\text{m}$ ) but differs in its higher number of preanal midventral supplementary organs (35 versus 20–22). Diagnosis of *G. menzelii* also lacks some important details, such as sizes of anterior sensilla, amphideal fovea, and spicules. Descriptions of *G. ferox* (New Hebrides [21]), *G. rapax* (North Sea [20]), and *G. rapax* (White Sea [15]) differ from the original diagnosis of Ssaweljev (1912 [26]) in their much smaller body size. Hence, we do not accept all these described specimens to be conspecific. Okhlopkov [15] redescribed females of *G. fennicum* and for the first time described males of *G. fennicum*. Though the specimens have not been sampled in the area of the type locality (Gulf of Finland, east Baltic Sea), but instead in the White Sea, the newly found females correspond to the original diagnosis in most details and morphometrics. Hence, *G. fennicum* is restored as a valid species.

### 3.1.3. Identification of Species of *Gammanema*

Since *Gammanema* species show evident sexual dimorphism in size and shape of amphids and other structures, it is highly desirable to have both sexes (or at least males) for species identification.

Due to deficiency of material, identification based on males is more reliable than identification solely based on females, since the males possess additional significant characteristics, such as supplementary organs and spicules.

For species identification of *Gammanema*, we propose a pictorial key constructed according to the principles of Platt [29], who first introduced pictorial keys in marine nematology. The pictorial key consists of two components, simplified images (caricatures) of heads (Figures 2 and 3), and Tables 1 and 2, with some metric and numeric characters being most important for species differentiation. Because of considerable sexual dimorphism, male and female head images are arranged in two separate plates, where the species are ordered in rows of gradual decrease of amphideal fovea (males) and cephalic setae (females), from top left to bottom right position.




**Figure 2.** Caricatured male heads of valid *Gammanema* species arranged from top left to bottom right along decrease of amphid size and cephalic setae in relation to head diameter.

**Table 1.** *Gammanema* valid species, comparison of males.

Species	Characters									
	Body Length	a	Inner Labial Setae Length	Outer Labial Setae Length	Cephalic Setae Length	Amphid Width	Supplements Shape	Number of Supplements	Spicule Sength	Some Particular Bold Features
<i>agglutinans</i>	544–696	10–11	2	2	10–11	10	tubular	≥6	34–36	Cuticle with minute spines, somatic setae leaf-shaped
<i>anthostoma</i>	1002–1950	22–29	5–6	5–8	6–8	14–26	absent	absent	62–87	Prominent circumoral projections each consisted of peduncle and triangular cap
<i>conicauda</i>	1985–2484	41–48	16	17	30–33	15	conical papillae	22	33–40	Prominent circumoral hair-like projections
<i>curvata</i>	3286	76	not indicated	17	13	8–8.5	cup-shaped	16	28	-
<i>fennicum</i>	2165–2600	28–49	17–24	8–11	34–50	28–31	sucker-like	13–21	62–64	-
<i>kosswigi</i>	1200	29	8	10–12	20	30	weak setose	4	39	Outer labial setae jointed
<i>magnum</i>	2900–4085	48–52	15–16	22–25	31–35	42–45	absent	absent	left 51–56, right 62–69	Spicules slightly unequal
<i>okhlopkovi</i>	961–1150	25–31	4–5	3–7	8–19	11–18	sucker-shaped	10–13	32–39	Prominent circumoral Projections peg-like
<i>polyodonta</i>	3200–4200	21–27	not indicated	papillae	16	42–45	cup-shaped	43–52	135	Very massive appearance; Anterior half of body opaque because of densely pigmented intestine
<i>smithi</i>	2210	41	not indicated	5	20	14	cup-shaped	20	78	-
<i>tchesunovi</i>	1603–1889	15–16	not indicated	5	8–8	15–18	cup-shaped	23–24	85–87	Tail rounded and conical, terminal cone not developed
<i>uniformis</i>	2700–2950	45–61	1–2	2–2	9–10	10–11	cup-shaped	19–22	82–87	Cephalic cuticle thickened and internally sclerotized around anterior chamber of pharyngostome

Table 2. *Gammanema* valid species, comparison of females.

Species	Characters						
	Body Length	a	Inner Labial Setae	Outer Labial Setae	Cephalic Setae	Amphid Width	Some Particular Bold Features
<i>agglutinans</i>	763–918	11–15	2	2	11–19	5	Cuticle with minute spines, somatic setae leaf-shaped
<i>anthostoma</i>	1487–1782	17–24	5–6.5	5–7.5	15–19	9.5–10.5	Prominent circumoral projections; each consisted of peduncle and triangular cap
<i>cancellatum</i>	2465	24	no data	40	15	8	Complex ornamentation of cuticle consists of dots and longitudinally arranged short rods
<i>conicauda</i>	2492–3724	30–35	16	12	30–33	8–9	Prominent circumoral hair-like projections
<i>curvata</i>	2392–2639	64–71	no data	15–17	12–13	8–8.5	-
<i>fennicum</i>	1672–2782	13–25	9–13	10–13	16–37	9.5–11	-
<i>magnum</i>	2540–3145	28–33	10–14	20–21	40–45	9–10	-
<i>mediterraneum</i>	660–745	18–20	papillae	papillae	7.5	5–6	-
<i>polyodonta</i>	3730–4050	23–31	papillae	papillae	16	20	Very massive appearance; anterior half of body opaque because of densely pigmented intestine
<i>tchesunovi</i>	1562–1682	11–12	papillae	5	7–8	16–18	Tail rounded and conical, terminal cone not developed
<i>uniformis</i>	2260–2790	40–57	1–2	2–2	9–12	9–10	Cephalic cuticle thickened and internally sclerotized around anterior chamber of pharyngostome



**Figure 3.** Caricatured female heads of valid *Gammanema* species arranged from top left to bottom right along decrease of cephalic setae in relation to head diameter.

### 3.2. Description of New Species of *Gammanema*

*Gammanema okhlopkovi* sp. n.


Figures 4–7, Table 3.

<http://zoobank.org/urn:lsid:zoobank.org:act:3B8336E6-9F74-4861-BCD3-94B27F52697B>

#### 3.2.1. Etymology

The species is named in honor of Yuri R. Okhlopkov, a PhD student who published a few papers on Selachinematidae, but unfortunately passed away in 2014 by mischance.






**Figure 4.** *Gammanema okhlopkovi* sp.n., entire view and anterior body: (A) holotype, male, entire view; (B) apical view, male, scanning electron micrograph; (C) subapical view, female, scanning electron micrograph; (D) head, lateral view, female, scanning electron micrograph; (E) head, sublateral view, male, scanning electron micrograph; (F) mouth opening, male, scanning electron micrograph. Scale bars: (A) 100 µm, (B) 10 µm, (C) 10 µm, (D) 10 µm, (E) 10 µm, (F) 3 µm.

### 3.2.2. Material Examined

One holotype and seven paratype males were deposited in the National Institute of Biological Resources (South Korea). Inventory numbers of the holotype and paratypes are NIBRIV0000861671 and NIBRIV000086172–NIBRIV0000861677, respectively.


Type locality: Intertidal zone at coast of Jeju Island, South Korea (33°26′05″ N, 126°55′15″ E), sandy beach, June 2018.



**Figure 5.** *Gammanema okhlopkovi* sp. n., head of the holotype male: (A) surface view; (B) optical section. Scale bars: 20  $\mu$ m.

### 3.2.3. Description


Body cylindrical, with truncate anterior end and short conical tail. Cuticle annulated and punctuated. There are ca. 13 annules within a 20  $\mu$ m area, everywhere along the body. Cuticle punctuation homogeneous, without lateral differentiation, the dots arranged in chessboard order. The dots are actually struts in the median zone of the cuticle; the struts are composed of two elements, as seen in sites of damaged cuticles.



**Figure 6.** *Gammanema okhlopkovi* sp. n., posterior body, males: (A) hind body; (B) spicule tips, precloacal seta, and supplementary organs, lateral view, scanning electron micrograph; (C) posterior-most supplementary organ, scanning electron micrograph; (D) cloacal opening and precloacal seta, subventral view, scanning electron micrograph; (E) tail with sensory papillae; (F) damaged cuticle near a supplement displays struts in the median zone, scanning electron micrograph. Scale bars: (A) 50  $\mu\text{m}$ , (B) 10  $\mu\text{m}$ , (C) 3  $\mu\text{m}$ , (D) 3  $\mu\text{m}$ , (E) 10  $\mu\text{m}$ , (F) 3  $\mu\text{m}$ .

Truncate anterior end bordered by a membranous fringe with indistinct sectioning in lips. The fringe is marked by longitudinal striations, with about 15 fine, short ribs or rugae between lateral outer labial sensilla and cephalic seta. There are twelve prominent peg-like projections around the rim of the anterior end.

Anterior sensilla arranged in 6 + 10 pattern. Six inner labial sensilla as pointed conical papillae situated just outside or the peg-like projections. Six outer labial sensilla situated at the base of the striated fringe; they are nearly equal in shape and size to the inner labial sensilla. Four cephalic setae at the same level with latero-median outer labial sensilla in tight pairs; within pair, the cephalic seta disposed laterally of the adjacent outer labial sensilla. Somatic sensilla represented by a few irregular postamphideal setae and two subsequent lateral papillae further posteriad. Somatic sensilla on the posterior body and tail region are represented by papillae sparsely arranged in latero-ventral, lateral, and dorso-lateral rows. Amphideal fovea large, transversally oval in outer outline, spirally coiled in three turns.



**Figure 7.** *Gammanema okhlopkovi* sp. n., posterior body of the holotype male. Scale bar: 20  $\mu$ m.

The mouth opening is situated on the bottom of a funnel-shaped depression on the truncate anterior end. Acute teeth arranged in three radiating groups are visible, even in a semi-open mouth. Cheilostome is narrow and indistinct. Pharyngostome consists of two chambers. Anterior chamber is cup-shaped. Its walls are strengthened by three sclerotized rhabdions terminating posteriorly, with a group of five small cusps or denticles projected freely into the buccal cavity. The anterior rhabdions spread widely, thus making the anterior chamber bowl-shaped. In each group, the median tooth is tridentate, while teeth of two lateral pairs are bidentate. Posterior chamber cylindrical and narrow; its walls are strengthened by three sclerotized rhabdions. Pharynx cylindrical and evenly strongly muscular along its entire length.

Reproductive system is diorchic. Anterior outstretched testis situated ventrally to the intestine; posterior reflexed testis situated to the right of the vas deferens and intestine. Spicules paired, arcuate, distally pointed, and proximally very slightly cephalated. Gubernaculum as paired, nearly straight bars directed dorsally. Midventral supplementary organs, ten to thirteen in number, shaped as a nail-head 5–6  $\mu$ m in diameter; the posteriormost organ situated at a distance 39  $\mu$ m from the cloacal opening. A short conical midventral precloacal seta is present.

Tail very short; its conical tip turned left in all the specimens.

**Table 3.** Morphometrics of males *Gammanema okhlopkovi* sp. n. (holotype and paratypes).

Character	Holotype	Total Number	Min.–Max.	Mean	SD	CV%
Body length	961	8	961–1150	1082	61.8	5.71
pharynx length	178	8	164–207	185	15.1	8.15
tail length	48	7	42–57.5	50.4	5.6	11.1
body diameter at level of cs	37	7	36.9–43.7	39.2	2.40	6.12
body diameter at level of nerve ring	35	8	33.1–40.0	36.8	2.32	6.30
body diameter at level of cardia	35	8	34.1–41.0	36.5	2.45	6.72
body diameter at level of midbody	37	8	37.0–42.0	38.8	2.54	6.65
body diameter at level of cloaca	36	8	36.0–42.5	38.2	2.54	6.65
a	26	8	25.6–31.1	27.8	2.17	7.80
b	5.4	8	5.40–6.56	5.88	0.41	6.97
c	20	7	18.5–24.6	21.6	2.32	10.7
c'	1.33	7	1.01–1.53	1.31	0.16	12.2
inner labial setae length	3.5–4	6	4.0–5.0	4.27	0.39	9.13
lateral outer labial setae length	3.7	6	3.0–7.2	5.22	1.63	31.2
latero-median outer labial setae length	5	5	5.0–7.0	5.88	0.76	12.9
cephalic setae length	12	6	8.5–19.0	14.1	4.01	28.5
anterior cervical setae length	7.6	7	7.6–15.6	10.8	2.56	23.8
amphid width	16	7	11.0–18.0	14.8	2.01	9.13
amphid width/corresponding body diameter	43	8	32.5–47.7	37.7	5.22	13.9
distance amphid—cephalic apex	8.6	8	7.3–11.3	8.95	1.37	15.3
anterior pharyngostome rhabdion length	15	8	13.0–16.9	14.9	1.49	10.0
posterior pharyngostome rhabdion length	14	8	11.2–16.0	13.5	1.40	10.3
spicule length (arc)	34	6	32.0–39.0	35.2	2.86	8.12
spicule 'length (chord)	28	6	29.0–36.0	32.9	2.4	7.29
number of supplements	12	6	10–13	11.5	1.05	9.13
tail terminal cone length	17	4	16.8–23.5	19.6	2.93	15.0

### 3.2.4. Diagnosis

*Gammanema* with short and stout body (L 961–1150  $\mu\text{m}$ , a 25–31). Twelve prominent peg-like projections protruded outward from inner side of the circumoral membranous fringe. Set of anterior sensilla composed of six inner labial setae 4–5  $\mu\text{m}$ , six outer labial setae 5–7  $\mu\text{m}$ , and cephalic setae 9–19  $\mu\text{m}$  long. Amphideal fovea spirally coiled in three turns, 32–48  $\mu\text{m}$  wide. Rhabdion of anterior pharyngostome chamber 13–17  $\mu\text{m}$ , rhabdion of posterior pharyngostome chamber 11–16  $\mu\text{m}$  long. Spicules 32–39  $\mu\text{m}$  long. Midventral preanal row of supplementary organs consists of precloacal seta and 10–13 nail-head organs. Tail short and conical, c' 1–1.6.

### 3.2.5. Differential Diagnosis

*G. okhlopkovi* sp. n. is closest to *G. anthostoma*, sharing similar body length and body proportions a, b, and c, and similar size of amphids and spicules. The most obvious differences between the two species are longer cephalic setae (8.5–19  $\mu\text{m}$  in *G. okhlopkovi* versus 6–7.5  $\mu\text{m}$  in *G. anthostoma*), as well as a lack of precloacal supplementary organs in *G. anthostoma* (present in *G. okhlopkovi*).

### 3.3. Description of New Species of *Latronema*

Genus *Latronema* (Wieser 1954)

Diagnosis of the genus has been recently updated by Leduc and Zhao [30]. A detailed review of the taxon is intended in forthcoming work on a peculiar diversity of *Latronema* species on an Atlantic seamount.

*Latronema obscuramphis* sp. n.

<http://zoobank.org/urn:lsid:zoobank.org:act:5E195F44-8FDA-48D4-9E71-00855406670D>

Figures 8 and 9, Table 4.

#### 3.3.1. Etymology

The species name reflects the indistinctness of the amphids (from Latin "obscurus", inconspicuous).


### 3.3.2. Material Examined

One holotype male, two paratype females, and four juveniles were deposited in National Institute of Biological Resources (South Korea). Inventory numbers of the holotype and paratypes are NIBRIV0000861678 and NIBRIV0000861679–NIBRIV0000861680, respectively.

### 3.3.3. Description

Body stout cylindrical, with truncate anterior end and short conical tail. Body cuticle finely cross-striated (16 annules in 20  $\mu\text{m}$  in light microscope, 12 annules in 10  $\mu\text{m}$ , from scanning electron micrographs) in the midbody laterally. The cuticle is not punctated under light microscope but marked with longitudinal wings. The wings number approximately five, visible on the lateral body side. Anteriorly, short and weak extra wings are added between the main wings, close to the head.


Labial region is presented by circular labial membrane with fine longitudinal ribbing (about ten ribs in between two adjacent setae of cephalic crown counted under light microscope, or seven to ten in SEM). Since the edges of the labial membrane are rolled inward, it is impossible to discern whether the membrane is divided into labial lobes. Labial region is encircled by a light cuticular ridge (collar).



**Figure 8.** *Latronema obscuramphis* sp. n., entire body of the holotype male and surface cuticular structure: (A) holotype male, entire view; (B) entire view of a female, scanning electron micrograph; (C) head of a female laterally, scanning electron micrograph; (D) amphideal fovea of a female, scanning electron micrograph; (E) body cuticle rings and alae laterally, scanning electron micrograph. Scale bars: (A) 50  $\mu\text{m}$ , (B) 30  $\mu\text{m}$ , (C,E) 10  $\mu\text{m}$ , (D) 3  $\mu\text{m}$ .

Six slender inner labial setae are situated at the edge of the circular labial membrane. Ten equal setae located at the edge of the collar; they are about equal in length to one another and equal to inner

labial setae, but wider basally; all the setae of the anterior circle look delicate and nearly transparent, and hence may be hardly visible against the labial cuticle. Fine ribs (striae) on the circumoral membrane may be arranged in fan-like bundles at sites opposite to cheilostomatal lobes. Quite possible the circle of ten equal setae comprises six outer labial and four cephalic setae, but all the setae are located at the same distance between them and do not make close pairs. Somatic setae comparable in length with anterior setae are distributed along the body in irregular longitudinal sublateral rows. Amphids in males are scarcely discernible under light microscope but detected in SEM as a small pit with simplified spiral structure. Amphids in females are not discernible at all under light microscope.



**Figure 9.** *Latronema obscuramphis* sp. n., male holotype structures: (A) surface view of the head; (B) sagittal optical section of the head; (C) posterior body. Scale bars: 20  $\mu$ m.

Buccal cavity consists of comparatively narrow cheilostome surrounded by labial membrane and voluminous pharyngostome composed of two compartments, with the anterior being cup-shaped and the posterior cylindrical. Cheilostome walls are differentiated interiorly into twelve rounded lobes that do not continue into anterior free projections, unlike that of *Gammanema*. Walls of the anterior pharyngostome are strengthened by three rhabdions—one dorsal and two lateroventral. The rhabdions terminate posteriorly with about seven acute denticles (one median denticle and two sets of three lateral denticles on both sides of it). Anterior pharyngostome is opened widely, and its rhabdions move

apart at obtuse angle. Posterior pharyngostome short and wide, its rhabdions slightly sclerotized, while intermediate cuticle between them thin and soft.

**Table 4.** *Latronema obscuramphis* sp. n., morphometry.

Character	Holotype Male	Paratype Female 1	Paratype Female 2
Body length	591	666	718
a	15.2	13.9	19.0
b	5.09	4.38	3.78
c	14.4	tail not measurable	tail not measurable
c'	1.60	tail not measurable	tail not measurable
V, %	-	57.4	70.0
body diameter at level of cephalic setae	30.0	38.0	33.1
body diameter at level of nerve ring	30.5	?	30.6
body diameter at level of cardia	35.0	41.5	33.8
body diameter at level of midbody	39.0	48.0	37.0
body diameter at level of cloaca/anus	25.6	30.0	?
inner labial setae length	8.7	7.7	6.0
latero-median outer labial setae length	11.6	not measurable	not measurable
lateral outer labial setae length	10.0	5.5	6.3
cephalic setae length	8.8	not measurable	not measurable
anterior cervical setae length	10.0	12.0	not measurable
anterior pharyngostome rhabdion length	12.0	12.0	12.4
posterior pharyngostome rhabdion length	8.0	7.5	10.7
spicules length (arc)	38.0	-	-
spicules length (chord)	33.0	-	-
gubernaculum length	12.0	-	-
tail terminal cone length	16	not measurable	not measurable

Pharynx cylindrical and strongly muscular. Nerve ring hardly discernible. Intestine is composed of convex enterocytes bulging in the gut lumen. Gut content in the holotype male is a long, fine-granular body, possibly a digested prey nematode in the posterior half of the intestine.

Female reproductive system is didelphic, both ovaries short and small, antidromously reflexed; in one female, anterior ovary to the right and posterior ovary to the left of the intestine, in other female anterior ovary to the left and posterior to the right of the intestine. Both spermathecas and uteri branch are filled with spermatozoa. Posterior ovary contains large oocyte with coarsely granulated cytoplasm and large nucleolated nucleus in the center. Somatic cuticle around vulva is thickened and distinctly set off as a vulvar plate.

Male reproductive system is diorchic. Both anterior outstretched and posterior reflexed testes situated ventrally and to the right of the intestine. Spicules equal, slightly arcuate, distally narrowed, proximally located, with very weakly developed knobs. Gubernaculum as a pair of short bars parallel to posterodistal edge of spicules. Preanal set of midventral supplementary organs composed of a midventral preanal seta and a row of about eight nail-head-shaped organs. The latter are not quite distinctly observed because of their position on the concave body side and covering by foreign particles.

Tail short and conical; its terminal cone covered with smooth cuticle and curved to the left.

### 3.3.4. Diagnosis

*Latronema* with small and stout body (L 591–718  $\mu\text{m}$ , a 13.9–19). All the anterior setae (i.e., inner labial setae, outer labial setae, and cephalic setae) are similar in length, ca 9–12  $\mu\text{m}$  in males, 6–6.5  $\mu\text{m}$  in females. Amphideal fovea small and obscure. Anterior pharyngostome rhabdion 12–12.5  $\mu\text{m}$  long, posterior rhabdions 7.5–11  $\mu\text{m}$ . Spicules arcuate, 38  $\mu\text{m}$  long, gubernaculum 12  $\mu\text{m}$ . Preanal midventral row of supplementary organs made up of preanal conical seta and twelve sucker-like organs. Tail very short (c 14.4, c' 1.6), with terminal cone ca 16  $\mu\text{m}$  long usually turned left.



### 3.3.5. Differential Diagnosis

*L. obscuramphis* sp. n. differs considerably from most *Latronema* species by combination of body size, anterior setae length, size and shape of amphideal fovea, size of spicules, number of preanal supplementary organs. *L. aberrans* (Allgén, 1934), *L. annulatum* (Gerlach, 1953), and *L. spinosum* (Andrássy, 1973) are closest to *L. obscuramphis* in characters cited above. *L. obscuramphis* differs from *L. aberrans* (the species known from the west Baltic Sea and White Sea) by smaller body length (male, L 591 versus 960  $\mu\text{m}$ ; females, L 666–718 versus 1140  $\mu\text{m}$ ) and lesser number of preanal supplementary organs (8 versus 13); from *L. annulatum* (Madagascar, littoral) by tail shape (in male  $c'$  1.64 versus 2.4–2.8), longer spicules (38  $\mu\text{m}$  versus 27  $\mu\text{m}$ ), and lesser number of supplements (8 versus 12); and from *L. spinosum* (Cuba, only female known) by no spinose body cuticular rings, smaller body (666–718 versus 850–1062  $\mu\text{m}$ ), and lesser index b (3.8–4.4 versus 4.5–5).

## 4. Conclusions

The nematofauna of the intertidal sandy beach of Jeju Island is very rich, including over 70 species representing nearly all the families of marine free-living nematodes. Even described species of Thoracostomopsidae [5,6] and Selachinematidae (present paper) constitute only about one-third of the actual diversity of those families from the studied Jeju beach. In forthcoming works, we will try to identify at least the most common nematode species, aiming in the future to reveal fauna composition and species distribution from upper to lower intertidal horizons, as well as vertical distribution in sediment columns.

**Author Contributions:** Conceptualization, A.T. and W.L.; investigation, A.T. and R.J.; data curation, R.J. and W.L.; writing—original draft preparation, A.T.; writing—review and editing, R.J. and W.L.; funding acquisition, W.L. and A.T. All authors have read and agreed to the published version of the manuscript.

**Funding:** The study was supported by a grant 18-504-51026 Russian Fund of Basic Research, a grant NRF-2017K2A9A1A06051528 from the Korea Research Foundation, a grant NIBR202000000 entitled “Study of Undiscovered Invertebrate Species from Korea” from the National Institute of Biological Resources (NIBR) funded by the Ministry of Environment (MOE) of the Republic of Korea, as well as by the Marine Biotechnology Program of the Korea Institute of Marine Science and Technology Promotion (KIMST) funded by the Ministry of Oceans and Fisheries (MOF) (No. 20170431).

**Acknowledgments:** The authors thank Vadim Mokievsky (P.P. Shirshov Institute of Oceanology, Moscow) and Jung-ho Hong (Biodiversity Institute, Marine Act Co., Seoul) for their help in the field work sampling in June 2018. The authors are grateful to both anonymous reviewers for their remarks and advice, which helped us amend the manuscript.

**Conflicts of Interest:** The authors declare no conflict of interest.

## Abbreviations

a	body length divided by body diameter at midbody
b	body length divided by pharynx length
c	body length divided by tail length
$c'$	tail length divided by anal body diameter
CV	coefficient of variation (SD divided by mean, in %)
SD	standard deviation
V	distance from anterior end to vulva in % of entire body length

## References

1. Rho, H.S.; Kim, W. *Paradraconema jejuense*, a new species of genus *Paradraconema* (Nematoda: Draconematidae) from Korea. *Korean J. Syst. Zool.* **2004**, *21*, 81–91.
2. Rho, H.S.; Kim, W. A new free-living marine nematode species of the genus *Dracogalerus* Allen and Noffsinger (Nematoda: Draconematidae) from a shallow subtidal zone of Jeju Island, Korea. *Integr. Biosci.* **2005**, *9*, 113–122. [[CrossRef](#)]

3. Rho, H.S.; Decraemer, W.; Sørensen, M.V.; Min, W.G.; Jung, J.; Kim, W. *Megadraconema cornutum*, a new genus and species from Korea, with a discussion of its classification and relationships within the family Draconematidae (Nematoda, Desmodorida) based on morphological and molecular characters. *Zool. Sci.* **2011**, *28*, 68–84. [[CrossRef](#)]
4. Lim, H.W.; Chang, C.Y. First record of *Desmoscolex* Nematoda (Desmoscolecida: Desmoscolecidae) from Korea. *Integr. Biosci.* **2006**, *10*, 219–225. [[CrossRef](#)]
5. Jeong, R.; Tchesunov, A.V.; Lee, W. Bibliographic revision of *Mesacanthion* Filipjev, 1927 (Nematoda: Thoracostomopsidae) with description of a new species from Jeju Island, South Korea. *PeerJ* **2019**, *7*, e8023. [[CrossRef](#)] [[PubMed](#)]
6. Jeong, R.; Tchesunov, A.V.; Lee, W. Two species of Thoracostomopsidae (Nematoda: Enoplida) from Jeju Island, South Korea. *PeerJ*, submitted, under review.
7. Okhlopkov, J.R. Feeding of free-living nematodes of the families Selachinematidae and Richtersiidae in the White Sea. *Proc. Pertsov White Sea Biol. Stn.* **2003**, *9*, 127–139. (In Russian)
8. Moens, T.; Braeckman, U.; Derycke, S.; Fonseca, F.; Galucci, F.; Gingold, R.; Guilini, K.; Ingels, J.; Leduc, D.; Vanaverbeke, J.; et al. Ecology of free-living marine nematodes. In *Handbook of Zoology*; Schmidt-Rhaesa, A., Ed.; De Gruyter: Berlin, Germany, 2014; Volume 2, pp. 109–152.
9. Warwick, R.M. Nematode associations in the Exe estuary. *J. Mar. Biol. Assoc. UK* **1971**, *51*, 439–454. [[CrossRef](#)]
10. Gallucci, F.; Steyaert, M.; Moens, T. Can field distributions of marine predacious nematodes be explained by sediment constraints on their foraging success? *Mar. Ecol. Prog. Ser.* **2005**, *304*, 167–178. [[CrossRef](#)]
11. Burgess, R. An improved protocol for separating meiofauna from sediments using colloidal silica soils. *Mar. Ecol. Prog. Ser.* **2001**, *214*, 161–165. [[CrossRef](#)]
12. Seinhorst, J. A rapid method for the transfer of nematodes from fixative to anhydrous glycerin. *Nematologica* **1959**, *4*, 67–69. [[CrossRef](#)]
13. Leduc, D. Two new genera and five new species of Selachinematidae (Nematoda, Chromadorida) from the continental slope of New Zealand. *Eur. J. Taxon.* **2013**, *63*, 1–32. [[CrossRef](#)]
14. Tchesunov, A.V. Order Chromadorida Chitwood, 1933. In *Handbook of Zoology*; Schmidt-Rhaesa, A., Ed.; Gastrotricha, Cycloneuralia and Gnathifera, Volume 2: Nematoda; De Gruyter: Berlin, Germany, 2014; pp. 373–398.
15. Okhlopkov, J.R. Free-living nematodes of the families Selachinematidae and Richtersiidae in the White Sea (Nematoda, Chromadoria). *Zoosyst. Ross.* **2002**, *11*, 41–55.
16. Gerlach, S.A. Zur Kenntnis der freilebenden marinen Nematoden von San Salvador. *Z. für Wiss. Zool.* **1955**, *158*, 249–303.
17. Gerlach, S.A. Die Nematodenbesiedlung des Sandstrandes und des Küstengrundwassers an der italienischen Küste. I. Systematischer Teil. *Arch. Zool. Ital.* **1953**, *37*, 517–640.
18. Gagarin, V.G.; Klerman, A.K. New species of predatory chromadorids (Nematoda, Chromadorida) from the Mediterranean Sea. *Zool. Zhurnal* **2007**, *86*, 778–786, (In Russian with English Abstract).
19. Gerlach, S.A. Die Nematodenfauna der Uferzonen und des Küstengrundwassers am finnischen Meeresbusen. *Acta Zool. Fenn.* **1953**, *73*, 1–32.
20. Gerlach, S.A. *Revision der Choniolaimidae und Selachinematidae (Freilebende Meeres-Nematoden)*; Kosswig-Festschrift; Mitteilungen aus dem Hamburgischen Zoologischen Museum und Institut: Hamburg, Germany, 1964; pp. 23–50.
21. Cobb, N.A. One hundred new nemas (type species of 100 new genera). *Contrib. Sci. Nematol.* **1920**, *9*, 217–343.
22. Shi, B.; Xu, K. Two new rapacious nematodes from intertidal sediments, *Gammanema magnum* sp. nov. and *Synonchium caudatubatum* sp. nov. (Nematoda, Selachinematidae). *Eur. J. Taxon.* **2018**, *405*, 1–17. [[CrossRef](#)]
23. Vitiello, P. Nématodes libres marins des vases profondes du Golfe du Lion. II. Chromadorida. *Tethys* **1970**, *2*, 449–500.
24. Ditlevsen, H. Marine freeliving nematodes from Danish waters. *Vidensk. Meddelelser Dansk Naturhistorisk Foren. I Kjobenhavn* **1918**, *70*, 147–214.
25. Murphy, D.G. Chilean marine nematodes. *Veröff. Inst. für Meeresforsch. Bremerhav.* **1965**, *9*, 173–203.
26. Ssaweljev, S. Zur Kenntnis der freilebenden Nematoden des Kolafjords und des Relictensee Mogilnoje. *Trav. Soc. Imp. Nat. St. Pétersbg.* **1912**, *43*, 108–126.
27. Murphy, D.G. Free-living marine nematodes, I. *Southerniella youngi*, *Dagda phinneyi*, and *Gammanema smithi*, new species. *Proc. Helminthol. Soc. Wash.* **1964**, *31*, 190–198.

28. Tchesunov, A.V.; Okhlopkov, Y.R. On some selachinematid nematodes (Chromadorida: Selachinematidae) deposited in the collection of the Smithsonian National Museum of Natural History. *Nematology* **2006**, *8*, 21–44.
29. Platt, H.M. Pictorial taxonomic keys: Their construction and use for the identification of free-living marine nematodes. *Cah. Biol. Mar.* **1984**, *25*, 83–91.
30. Leduc, D.; Zhao, Z. *Latronema whataitai* sp. n. (Nematoda: Selachinematidae) from intertidal sediments of New Zealand, with notes on relationships within the family based on preliminary 18S and D2-D3 phylogenetic analyses. *Nematology* **2015**, *17*, 941–952. [[CrossRef](#)]



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).